

**SOME INEQUALITIES FOR THE EXPECTATION AND  
VARIANCE OF A RANDOM VARIABLE WHOSE PDF IS  
 $n$ -TIME DIFFERENTIABLE**

N. S. BARNETT, P. CERONE, S.S. DRAGOMIR, AND J. ROUMELIOTIS

ABSTRACT. Some inequalities for the expectation and variance of a random variable whose p.d.f. is  $n$ -time differentiable are given.

1. INTRODUCTION

Let  $f : [a, b] \rightarrow \mathbb{R}_+$  be the p.d.f. of the random variable  $X$  and

$$E(X) := \int_a^b t f(t) dt$$

its *expectation* and

$$\begin{aligned} \sigma(X) &= \left[ \int_a^b (t - E(X))^2 f(t) dt \right]^{\frac{1}{2}} \\ &= \left[ \int_a^b t^2 f(t) dt - [E(X)]^2 \right]^{\frac{1}{2}} \end{aligned}$$

its dispersion or standard deviation.

In [1], using the identity

$$(1.1) \quad [x - E(X)]^2 + \sigma^2(X) = \int_a^b (x - t)^2 f(t) dt$$

and applying a variety of inequalities such as: Hölder's inequality, pre-Grüss, pre-Chebychev, pre-Lupaş, or Ostrowski type inequalities, a number of results concerning the expectation and variance of the random variable  $X$  were obtained.

For example,

$$(1.2) \quad \begin{aligned} &\sigma^2(X) + [x - E(X)]^2 \\ &\leq \begin{cases} (b-a) \left[ \frac{(b-a)^2}{12} + \left(x - \frac{a+b}{2}\right)^2 \right] \|f\|_\infty, & \text{if } f \in L_\infty[a, b]; \\ \left[ \frac{(b-x)^{2q+1} + (x-a)^{2q+1}}{2q+1} \right]^{\frac{1}{q}} \|f\|_p, & \text{if } f \in L_p[a, b], \\ & p > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \left( \frac{b-a}{2} + \left| x - \frac{a+b}{2} \right| \right)^2, & \end{cases} \end{aligned}$$

---

*Date:* January, 2000.

*1991 Mathematics Subject Classification.* Primary 60E15, 26D15.

*Key words and phrases.* Random Variable, Expectation, Variance, Dispersion.

for all  $x \in [a, b]$ , which imply, amongst other things, that

$$(1.3) \quad \begin{aligned} 0 &\leq \sigma(X) \\ &\leq \begin{cases} (b-a)^{\frac{1}{2}} \left[ \frac{(b-a)^2}{12} + [E(X) - \frac{a+b}{2}]^2 \right]^{\frac{1}{2}} \|f\|_{\infty}^{\frac{1}{2}}, & \text{if } f \in L_{\infty}[a, b]; \\ \left\{ \frac{[b-E(X)]^{2q+1} + [E(X)-a]^{2q+1}}{2q+1} \right\}^{\frac{1}{2q}} \|f\|_p^{\frac{1}{2}}, & \text{if } f \in L_p[a, b], \\ & p > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \frac{b-a}{2} + |E(X) - \frac{a+b}{2}|, & \end{cases} \end{aligned}$$

and

$$(1.4) \quad 0 \leq \sigma^2(X) \leq [b - E(X)][E(X) - a] \leq \frac{1}{4}(b-a)^2.$$

In this paper more accurate inequalities are obtained by assuming that the p.d.f. of  $X$  is  $n$ -time differentiable and that  $f^{(n)}$  is absolutely continuous on  $[a, b]$ . For other recent results on the application of Ostrowski type inequalities in Probability Theory, see [2]-[4].

## 2. SOME PRELIMINARY INTEGRAL IDENTITIES

The following lemma, which is interesting in itself, holds.

**Lemma 1.** *Let  $X$  be a random variable whose probability distribution function  $f : [a, b] \rightarrow \mathbb{R}_+$  is  $n$ -time differentiable and  $f^{(n)}$  is absolutely continuous on  $[a, b]$ . Then*

$$(2.1) \quad \begin{aligned} &\sigma^2(X) + [E(X) - x]^2 \\ &= \sum_{k=0}^n \frac{(b-x)^{k+3} + (-1)^k (x-a)^{k+3}}{(k+3)k!} f^{(k)}(x) \\ &\quad + \frac{1}{n!} \int_a^b (t-x)^2 \left( \int_x^t (t-s)^n f^{(n+1)}(s) ds \right) dt \end{aligned}$$

for all  $x \in [a, b]$ .

*Proof.* Is by Taylor's formula with integral remainder. Recall that

$$(2.2) \quad f(t) = \sum_{k=0}^n \frac{(t-x)^k}{k!} f^{(k)}(x) + \frac{1}{n!} \int_x^t (t-s)^n f^{(n+1)}(s) ds$$

for all  $t, x \in [a, b]$ .

Together with

$$(2.3) \quad \sigma^2(X) + [E(X) - x]^2 = \int_a^b (t-x)^2 f(t) dt,$$

where  $f$  is the p.d.f. of the random variable  $X$ , we obtain

$$(2.4) \quad \begin{aligned} &\sigma^2(X) + [E(X) - x]^2 \\ &= \int_a^b (t-x)^2 \left[ \sum_{k=0}^n \frac{(t-x)^k}{k!} f^{(k)}(x) + \frac{1}{n!} \int_x^t (t-s)^n f^{(n+1)}(s) ds \right] dt \\ &= \sum_{k=0}^n f^{(k)}(x) \int_a^b \frac{(t-x)^{k+2}}{k!} dt + \frac{1}{n!} \int_a^b (t-x)^2 \left( \int_x^t (t-s)^n f^{(n+1)}(s) ds \right) dt \end{aligned}$$

and since

$$\int_a^b \frac{(t-x)^{k+2}}{k!} dt = \frac{(b-x)^{k+3} + (-1)^k (x-a)^{k+3}}{(k+3)k!},$$

the identity (2.4) readily produces (2.1) ■

**Corollary 1.** *Under the above assumptions, we have*

$$(2.5) \quad \begin{aligned} & \sigma^2(X) + \left[ E(X) - \frac{a+b}{2} \right]^2 \\ &= \sum_{k=0}^n \frac{[1 + (-1)^k] (b-a)^{k+3}}{2^{k+3} (k+3) k!} f^{(k)}\left(\frac{a+b}{2}\right) \\ & \quad + \frac{1}{n!} \int_a^b \left(t - \frac{a+b}{2}\right)^2 \left( \int_{\frac{a+b}{2}}^t (t-s)^n f^{(n+1)}(s) ds \right) dt. \end{aligned}$$

The proof follows by using (2.4) with  $x = \frac{a+b}{2}$ .

**Corollary 2.** *Under the above assumptions,*

$$(2.6) \quad \begin{aligned} & \sigma^2(X) + \frac{1}{2} \left[ (E(X) - a)^2 + (E(X) - b)^2 \right] \\ &= \sum_{k=0}^n \frac{(b-a)^{k+3}}{(k+3)k!} \left[ \frac{f^{(k)}(a) + (-1)^k f^{(k)}(b)}{2} \right] \\ & \quad + \frac{1}{n!} \int_a^b \int_a^b K(t,s) (t-s)^n f^{(n+1)}(s) ds dt, \end{aligned}$$

where

$$K(t,s) := \begin{cases} \frac{(t-a)^2}{2} & \text{if } a \leq s \leq t \leq b, \\ -\frac{(t-b)^2}{2} & \text{if } a \leq t < s \leq b. \end{cases}$$

*Proof.* In (2.1), choose  $x = a$  and  $x = b$ , giving

$$(2.7) \quad \begin{aligned} & \sigma^2(X) + [E(X) - a]^2 \\ &= \sum_{k=0}^n \frac{(b-a)^{k+3}}{(k+3)k!} f^{(k)}(a) + \frac{1}{n!} \int_a^b (t-a)^2 \left( \int_a^t (t-s)^n f^{(n+1)}(s) ds \right) dt \end{aligned}$$

and

$$(2.8) \quad \begin{aligned} & \sigma^2(X) + [E(X) - b]^2 \\ &= \sum_{k=0}^n \frac{(-1)^k (b-a)^{k+3}}{(k+3)k!} f^{(k)}(b) + \frac{1}{n!} \int_a^b (t-b)^2 \left( \int_b^t (t-s)^n f^{(n+1)}(s) ds \right) dt. \end{aligned}$$

Adding these and dividing by 2 gives (2.6). ■

Taking into account that  $\mu = E(X) \in [a, b]$ , then we also obtain the following.

**Corollary 3.** *With the above assumptions,*

$$(2.9) \quad \sigma^2(X) = \sum_{k=0}^n \frac{(b-\mu)^{k+3} + (-1)^k (\mu-a)^{k+3}}{(k+3)k!} f^{(k)}(\mu) \\ + \frac{1}{n!} \int_a^b (t-\mu)^2 \left( \int_\mu^t (t-s)^n f^{(n+1)}(s) ds \right) dt.$$

*Proof.* The proof follows from (2.1) with  $x = \mu \in [a, b]$ . ■

**Lemma 2.** *Let the conditions of Lemma 1 relating to  $f$  hold. Then the following identity is valid.*

$$(2.10) \quad \sigma^2(X) + [E(X) - x]^2 \\ = \sum_{k=0}^n \frac{(b-x)^{k+3} + (-1)^k (x-a)^{k+3}}{k+3} \cdot \frac{f^{(k)}(x)}{k!} \\ + \frac{1}{n!} \int_a^b K_n(x, s) f^{(n+1)}(s) ds,$$

where

$$(2.11) \quad K(x, s) = \begin{cases} (-1)^{n+1} \psi_n(s-a, x-s), & a \leq s \leq x \\ \psi_n(b-s, s-x), & x < s \leq b \end{cases}$$

with

$$(2.12) \quad \psi_n(u, v) = \frac{u^{n+1}}{(n+3)(n+2)(n+1)} \cdot [(n+2)(n+1)u^2 \\ + 2(n+3)(n+1)uv + (n+3)(n+2)v^2].$$

*Proof.* From (2.1), an interchange of the order of integration gives

$$\frac{1}{n!} \int_a^b (t-x)^2 dt \int_x^t (t-s)^n f^{(n+1)}(s) ds \\ = \frac{1}{n!} \left\{ - \int_a^x \int_a^s (t-x)^2 (t-s)^n f^{(n+1)}(s) dt ds \right. \\ \left. + \int_x^b \int_s^b (t-x)^2 (t-s)^n f^{(n+1)}(s) dt ds \right\} \\ = \frac{1}{n!} \int_a^b \tilde{K}_n(x, s) f^{(n+1)}(s) ds,$$

where

$$\tilde{K}_n(x, s) = \begin{cases} p_n(x, s) = - \int_a^s (t-x)^2 (t-s)^n dt, & a \leq s \leq x \\ q_n(x, s) = \int_s^b (t-x)^2 (t-s)^n dt, & x < s < b. \end{cases}$$

To prove the lemma it is sufficient to show that  $K \equiv \tilde{K}$ .

Now,

$$\begin{aligned}
\tilde{p}_n(x, s) &= - \int_a^s (t-x)^2 (t-s)^n dt \\
&= (-1)^{n+1} \int_0^{s-a} (u+x-s)^2 u^n du \\
&= (-1)^{n+1} \int_0^{s-a} [u^2 + 2(x-s)u + (x-s)^2] u^n du \\
&= (-1)^{n+1} \psi_n(s-a, x-s),
\end{aligned}$$

where  $\psi(\cdot, \cdot)$  is as given by (2.12).

Further,

$$\begin{aligned}
\tilde{q}_n(x, s) &= \int_s^b (t-x)^2 (t-s)^n dt \\
&= \int_0^{b-s} [u+(s-x)]^2 u^n du \\
&= \psi_n(b-s, s-x),
\end{aligned}$$

where, again,  $\psi(\cdot, \cdot)$  is as given by (2.12). Hence  $K \equiv \tilde{K}$  and the lemma is proved. ■

### 3. SOME INEQUALITIES

We are now able to obtain the following inequalities.

**Theorem 1.** *Let  $X$  be a random variable whose probability density function  $f : [a, b] \rightarrow \mathbb{R}_+$  is  $n$ -time differentiable and  $f^{(n)}$  is absolutely continuous on  $[a, b]$ , then*

$$(3.1) \quad \left| \sigma^2(X) + [E(X) - x]^2 - \sum_{k=0}^n \frac{(b-x)^{k+3} + (-1)^k (x-a)^{k+3}}{(k+3)k!} f^{(k)}(x) \right|$$

$$\leq \begin{cases} \frac{\|f^{(n+1)}\|_\infty}{(n+1)!(n+4)} [(x-a)^{n+4} + (b-x)^{n+4}], & \text{if } f^{(n+1)} \in L_\infty[a, b]; \\ \frac{\|f^{(n+1)}\|_p}{n!(n+3+\frac{1}{q})} \frac{[(x-a)^{n+3+\frac{1}{q}} + (b-x)^{n+3+\frac{1}{q}}]}{(nq+1)^{\frac{1}{q}}}, & \text{if } f^{(n+1)} \in L_p[a, b], \\ & p > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \frac{\|f^{(n+1)}\|_1}{n!(n+3)} [(x-a)^{n+3} + (b-x)^{n+3}], & \text{if } f^{(n+1)} \in L_1[a, b] \end{cases}$$

for all  $x \in [a, b]$ , where  $\|\cdot\|_p$  ( $1 \leq p \leq \infty$ ) are the usual Lebesgue norms on  $[a, b]$ , i.e.,

$$\|g\|_\infty := \text{ess sup}_{t \in [a, b]} |g(t)| \quad \text{and} \quad \|g\|_p := \left( \int_a^b |g(t)|^p dt \right)^{\frac{1}{p}}, \quad p \geq 1.$$

*Proof.* By Lemma 1,

$$\begin{aligned}
(3.2) \quad & \sigma^2(X) + [E(X) - x]^2 - \sum_{k=0}^n \frac{(b-x)^{k+3} + (-1)^k (x-a)^{k+3}}{k!(k+3)} f^{(k)}(x) \\
&= \frac{1}{n!} \int_a^b (t-x)^2 \left( \int_x^t (t-s)^n f^{(n+1)}(s) ds \right) dt := M(a, b; x).
\end{aligned}$$

Clearly,

$$\begin{aligned}
|M(a, b; x)| &\leq \frac{1}{n!} \int_a^b (t-x)^2 \left| \int_x^t (t-s)^n f^{(n+1)}(s) ds \right| dt \\
&\leq \frac{1}{n!} \int_a^b (t-x)^2 \left[ \sup_{s \in [x, t]} |f^{(n+1)}(s)| \left| \int_x^t |t-s|^n ds \right| \right] dt \\
&\leq \frac{\|f^{(n+1)}\|_\infty}{n!} \int_a^b \frac{(t-x)^2 |t-x|^{n+1}}{n+1} dt \\
&= \frac{\|f^{(n+1)}\|_\infty}{(n+1)!} \int_a^b |t-x|^{n+3} dt \\
&= \frac{\|f^{(n+1)}\|_\infty}{(n+1)!} \left[ \int_a^x (x-t)^{n+3} dt + \int_x^b (t-x)^{n+3} dt \right] \\
&= \frac{\|f^{(n+1)}\|_\infty}{(n+1)!} \frac{[(x-a)^{n+4} + (b-x)^{n+4}]}{(n+4)}
\end{aligned}$$

and the first inequality in (3.1) is obtained.

For the second, we use Hölder's integral inequality to obtain

$$\begin{aligned}
|M(a, b; x)| &\leq \frac{1}{n!} \int_a^b (t-x)^2 \left| \int_x^t |t-s|^{nq} ds \right|^{\frac{1}{q}} \left| \int_x^t |f^{(n+1)}(s)|^p ds \right|^{\frac{1}{p}} dt \\
&\leq \frac{1}{n!} \left( \int_a^b |f^{(n+1)}(s)|^p ds \right)^{\frac{1}{p}} \int_a^b (t-x)^2 |t-x|^{\frac{nq+1}{q}} dt \\
&= \frac{1}{n!} \frac{\|f^{(n+1)}\|_p}{(nq+1)^{\frac{1}{q}}} \int_a^b |t-x|^{n+2+\frac{1}{q}} dt \\
&= \frac{1}{n!} \frac{\|f^{(n+1)}\|_p}{(nq+1)^{\frac{1}{q}}} \left[ \frac{(b-x)^{n+3+\frac{1}{q}} + (x-a)^{n+3+\frac{1}{q}}}{n+3+\frac{1}{q}} \right].
\end{aligned}$$

Finally, note that

$$\begin{aligned}
|M(a, b; x)| &\leq \frac{1}{n!} \int_a^b (t-x)^2 |t-x|^n \left| \int_x^t |f^{(n+1)}(s)| ds \right| dt \\
&\leq \frac{\|f^{(n+1)}\|_1}{n!} \int_a^b |t-x|^{n+2} dt \\
&= \frac{\|f^{(n+1)}\|_1}{n!} \left[ \frac{(x-a)^{n+3} + (b-x)^{n+3}}{n+3} \right]
\end{aligned}$$

and the third part of (3.1) is obtained. ■

It is obvious that the best inequality in (3.1) is when  $x = \frac{a+b}{2}$ , giving Corollary 4.

**Corollary 4.** *With the above assumptions on  $X$  and  $f$ ,*

$$(3.3) \quad \left| \sigma^2(X) + \left[ E(X) - \frac{a+b}{2} \right]^2 - \sum_{k=0}^n \frac{[1 + (-1)^k] (b-a)^{k+3}}{2^{k+3} (k+3) k!} f^{(k)} \left( \frac{a+b}{2} \right) \right|$$

$$\leq \begin{cases} \frac{\|f^{(n+1)}\|_\infty}{2^{n+3} (n+1)! (n+4)} (b-a)^{n+4}, & \text{if } f^{(n+1)} \in L_\infty[a, b]; \\ \frac{\|f^{(n+1)}\|_p}{2^{n+2+\frac{1}{q}} n! (n+3+\frac{1}{q})} \frac{(b-a)^{n+3+\frac{1}{q}}}{(nq+1)^{\frac{1}{q}}}, & \text{if } f^{(n+1)} \in L_p[a, b], p > 1, \\ & \frac{1}{p} + \frac{1}{q} = 1; \\ \frac{\|f^{(n+1)}\|_1}{2^{n+2} n! (n+3)} (b-a)^{n+3}, & \text{if } f^{(n+1)} \in L_1[a, b]. \end{cases}$$

The following corollary is interesting as it provides the opportunity to approximate the variance when the values of  $f^{(k)}(\mu)$  are known,  $k = 0, \dots, n$ .

**Corollary 5.** *With the above assumptions and  $\mu = \frac{a+b}{2}$ , we have*

$$(3.4) \quad \left| \sigma^2(X) - \sum_{k=0}^n \frac{(b-\mu)^{k+3} + (-1)^k (\mu-a)^{k+3}}{(k+3) k!} f^{(k)}(\mu) \right|$$

$$\leq \begin{cases} \frac{\|f^{(n+1)}\|_\infty}{(n+1)! (n+4)} [(\mu-a)^{n+4} + (b-\mu)^{n+4}], & \text{if } f^{(n+1)} \in L_\infty[a, b]; \\ \frac{\|f^{(n+1)}\|_p}{n! (n+3+\frac{1}{q})} \frac{[(\mu-a)^{n+3+\frac{1}{q}} + (b-\mu)^{n+3+\frac{1}{q}}]}{(nq+1)^{\frac{1}{q}}}, & \text{if } f^{(n+1)} \in L_p[a, b], \\ & p > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \frac{\|f^{(n+1)}\|_1}{n! (n+3)} [(\mu-a)^{n+3} + (b-\mu)^{n+3}], & \text{if } f^{(n+1)} \in L_1[a, b]. \end{cases}$$

The following result also holds.

**Theorem 2.** *Let  $X$  be a random variable whose probability density function  $f : [a, b] \rightarrow \mathbb{R}_+$  is  $n$ -time differentiable and  $f^{(n)}$  is absolutely continuous on  $[a, b]$ , then*

$$(3.5) \quad \left| \sigma^2(X) + \frac{1}{2} \left[ (E(X) - a)^2 + (E(X) - b)^2 \right] - \sum_{k=0}^n \frac{(b-a)^{k+3}}{(k+3) k!} \left[ \frac{f^{(k)}(a) + (-1)^k f^{(k)}(b)}{2} \right] \right|$$

$$\leq \begin{cases} \frac{1}{(n+4)(n+1)!} \|f^{(n+1)}\|_\infty (b-a)^{n+4}, \\ \text{if } f^{(n+1)} \in L_\infty[a, b]; \\ \frac{1}{n! (qn+1)^{\frac{1}{q}} [(n+2)q+2]^{\frac{1}{q}}} \|f^{(n+1)}\|_p \frac{(b-a)^{n+3+\frac{1}{q}}}{(nq+1)^{\frac{1}{q}}}, \\ \text{if } f^{(n+1)} \in L_p[a, b], \text{ where } p > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \frac{1}{2n!} \|f^{(n+1)}\|_1 (b-a)^{n+3}, \end{cases}$$

where  $\|\cdot\|_p$  ( $1 \leq p \leq \infty$ ) are the usual Lebesgue  $p$ -norms.

*Proof.* Using Corollary 2,

$$\begin{aligned} & \left| \sigma^2(X) + \frac{1}{2} \left[ (E(X) - a)^2 + (E(X) - b)^2 \right] \right. \\ & \quad \left. - \sum_{k=0}^n \frac{(b-a)^{k+3}}{(k+3)k!} \left[ \frac{f^{(k)}(a) + (-1)^k f^{(k)}(b)}{2} \right] \right| \\ & \leq \frac{1}{n!} \int_a^b \int_a^b |K(t,s)| |t-s|^n |f^{(n+1)}(s)| ds dt =: N(a,b). \end{aligned}$$

It is obvious that

$$\begin{aligned} & N(a,b) \\ & \leq \|f^{(n+1)}\|_\infty \frac{1}{n!} \int_a^b \int_a^b |K(t,s)| |t-s|^n ds dt \\ & = \|f^{(n+1)}\|_\infty \frac{1}{n!} \int_a^b \left( \int_a^t |K(t,s)| |t-s|^n ds + \int_t^b |K(t,s)| |t-s|^n ds \right) dt \\ & = \frac{1}{n!} \|f^{(n+1)}\|_\infty \int_a^b \left[ \frac{(t-a)^2}{2} \cdot \frac{(t-a)^{n+1}}{n+1} + \frac{(t-b)^2}{2} \cdot \frac{(b-t)^{n+1}}{n+1} \right] dt \\ & = \frac{1}{2(n+1)!} \|f^{(n+1)}\|_\infty \int_a^b \left[ (t-a)^{n+3} + (b-t)^{n+3} \right] dt \\ & = \frac{1}{2(n+1)!} \|f^{(n+1)}\|_\infty \left[ \frac{(b-a)^{n+4}}{n+4} + \frac{(b-a)^{n+4}}{n+4} \right] \\ & = \frac{\|f^{(n+1)}\|_\infty}{(n+4)(n+1)!} (b-a)^{n+4} \end{aligned}$$

so the first part of (3.5) is proved.

Using Hölder's integral inequality for double integrals,

$$\begin{aligned} & N(a,b) \\ & \leq \frac{1}{n!} \left( \int_a^b \int_a^b |f^{(n+1)}(s)|^p ds dt \right)^{\frac{1}{p}} \times \left( \int_a^b \int_a^b |K(t,s)|^q |t-s|^{qn} ds dt \right)^{\frac{1}{q}} \\ & = \frac{(b-a)^{\frac{1}{p}} \|f^{(n+1)}\|_p}{n!} \times \\ & \quad \left[ \int_a^b \left( \int_a^t |K(t,s)|^q |t-s|^{qn} ds + \int_t^b |K(t,s)|^q |t-s|^{qn} ds \right) dt \right]^{\frac{1}{q}} \\ & = \frac{(b-a)^{\frac{1}{p}} \|f^{(n+1)}\|_p}{n!} \\ & \quad \times \left[ \int_a^b \left[ \frac{(t-a)^{2q}}{2^q} \int_a^t |t-s|^{qn} ds + \frac{(t-b)^{2q}}{2^q} \int_t^b |t-s|^{qn} ds \right] dt \right]^{\frac{1}{q}} \end{aligned}$$

$$\begin{aligned}
&= \frac{(b-a)^{\frac{1}{p}} \|f^{(n+1)}\|_p}{n!} \\
&\quad \times \left[ \int_a^b \left[ \frac{(t-a)^{2q} (t-a)^{qn+1}}{2^q (qn+1)} + \frac{(t-b)^{2q} (b-t)^{qn+1}}{2^q (qn+1)} \right] dt \right]^{\frac{1}{q}} \\
&= \frac{(b-a)^{\frac{1}{p}} \|f^{(n+1)}\|_p}{n!} \cdot \left[ \frac{1}{2^q (qn+1)} \right]^{\frac{1}{q}} \\
&\quad \times \left[ \int_a^b (t-a)^{(n+2)q+1} dt + \int_a^b (b-t)^{(n+2)q+1} dt \right]^{\frac{1}{q}} \\
&= \frac{(b-a)^{\frac{1}{p}} \|f^{(n+1)}\|_p}{n!} \cdot \left[ \frac{1}{2^q (qn+1)} \right]^{\frac{1}{q}} \left[ \frac{(b-a)^{(n+2)q+2}}{(n+2)q+2} + \frac{(b-a)^{(n+2)q+2}}{(n+2)q+2} \right]^{\frac{1}{q}} \\
&= \frac{2 \|f^{(n+1)}\|_p (b-a)^{n+2+\frac{1}{p}+\frac{2}{q}}}{n! 2^q (qn+1)^{\frac{1}{q}} ((n+2)q+2)^{\frac{1}{q}}} \\
&= \frac{\|f^{(n+1)}\|_p \left[ (b-a)^{n+3+\frac{1}{q}} \right]}{n! (qn+1)^{\frac{1}{q}} [(n+2)q+2]^{\frac{1}{q}}}
\end{aligned}$$

and the second part of (3.5) is proved.

Finally, we observe that

$$\begin{aligned}
N(a, b) &\leq \frac{1}{n!} \sup_{(t,s) \in [a,b]^2} |K(t,s)| |t-s|^n \int_a^b \int_a^b |f^{(n+1)}(s)| ds dt \\
&= \frac{1}{n!} \frac{(b-a)^2}{2} \cdot (b-a)^n (b-a) \int_a^b |f^{(n+1)}(s)| ds \\
&= \frac{1}{2n!} (b-a)^{n+3} \|f^{(n+1)}\|_1,
\end{aligned}$$

which is the final result of (3.5). ■

The following particular cases can be useful in practical applications.

1. For  $n = 0$ , (3.1) becomes

$$\begin{aligned}
(3.6) \quad &\left| \sigma^2(X) + [E(X) - x]^2 - (b-a) \left[ \left( x - \frac{a+b}{2} \right)^2 + \frac{(b-a)^2}{12} \right] f(x) \right| \\
&\leq \begin{cases} \frac{\|f'\|_\infty}{4} [(x-a)^4 + (b-x)^4], & \text{if } f' \in L_\infty[a, b]; \\ \frac{q\|f'\|_p}{3q+1} [(x-a)^{3+\frac{1}{q}} + (b-x)^{3+\frac{1}{q}}], & \text{if } f' \in L_p[a, b], \\ & p > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \|f'\|_1 \left[ \frac{(b-a)^2}{12} + \left( x - \frac{a+b}{2} \right)^2 \right], & \text{if } f' \in L_1[a, b], \end{cases}
\end{aligned}$$

for all  $x \in [a, b]$ .

In particular, for  $x = \frac{a+b}{2}$ ,

$$(3.7) \quad \left| \sigma^2(X) + \left[ E(X) - \frac{a+b}{2} \right]^2 - \frac{(b-a)^3}{12} f\left(\frac{a+b}{2}\right) \right| \leq \begin{cases} \frac{\|f'\|_\infty}{32} (b-a)^4, & \text{if } f' \in L_\infty[a, b]; \\ \frac{q\|f'\|_p (b-a)^{3+\frac{1}{q}}}{2^{2+\frac{1}{q}}(3q+1)}, & \text{if } f' \in L_p[a, b], \\ & p > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \frac{\|f'\|_1}{12} (b-a)^3, & \end{cases}$$

which is, in a sense, the best inequality that can be obtained from (3.6). If in (3.6)  $x = \mu = E(X)$ , then

$$(3.8) \quad \left| \sigma^2(X) - (b-a) \left[ \left( E(X) - \frac{a+b}{2} \right)^2 + \frac{(b-a)^2}{12} \right] f(E(X)) \right| \leq \begin{cases} \frac{\|f'\|_\infty}{4} \left[ (E(X) - a)^4 + (b - E(X))^4 \right], & \text{if } f' \in L_\infty[a, b]; \\ \frac{\|f'\|_p}{(3+\frac{1}{q})} \left[ (E(X) - a)^4 + (b - E(X))^4 \right], & \text{if } f' \in L_p[a, b], p > 1, \\ \|f'\|_1 \left[ \frac{(b-a)^2}{12} + \left( E(X) - \frac{a+b}{2} \right)^2 \right], & \text{if } f' \in L_1[a, b]. \end{cases}$$

In addition, from (3.5),

$$(3.9) \quad \left| \sigma^2(X) + \frac{1}{2} \left[ (E(X) - a)^2 + (E(X) - b)^2 \right] - \frac{(b-a)^3}{3} \left[ \frac{f(a) + f(b)}{2} \right] \right| \leq \begin{cases} \frac{1}{4} \|f'\|_\infty (b-a)^4, & \text{if } f' \in L_\infty[a, b]; \\ \frac{1}{n!2^{\frac{1}{q}}(q+1)^{\frac{1}{q}}} \|f'\|_p (b-a)^{3+\frac{1}{q}}, & \text{if } f' \in L_p[a, b], p > 1, \\ \frac{1}{2} \|f'\|_1 (b-a)^3, & \end{cases}$$

which provides an approximation for the variance in terms of the expectation and the values of  $f$  at the end points  $a$  and  $b$ .

**Theorem 3.** Let  $X$  be a random variable whose p.d.f.  $f : [a, b] \rightarrow \mathbb{R}_+$  is  $n$ -time differentiable and  $f^{(n)}$  is absolutely continuous on  $[a, b]$ . Then

$$(3.10) \quad \left| \sigma^2(X) + (E(X) - x)^2 - \sum_{k=0}^n \frac{(b-x)^{k+3} + (-1)^k (x-a)^{k+3}}{k+3} \cdot \frac{f^{(k)}(x)}{k!} \right|$$

$$\leq \begin{cases} \left[ (x-a)^{n+4} + (b-x)^{n+4} \right] \frac{\|f^{(n+1)}\|_\infty}{(n+1)!(n+4)}, \\ C^{\frac{1}{q}} \left[ (x-a)^{(n+3)q+1} + (b-x)^{(n+3)q+1} \right]^{\frac{1}{q}} \frac{\|f^{(n+1)}\|_p}{n!}, \\ \left[ \frac{b-a}{2} + \left| x - \frac{a+b}{2} \right| \right]^{n+3} \cdot \frac{\|f^{(n+1)}\|_1}{n!(n+3)}, \end{cases}$$

where

$$(3.11) \quad C = \int_0^1 \left[ \frac{u^{n+3}}{n+3} + 2(1-u) \frac{u^{n+2}}{n+2} + (1-u)^2 \frac{u^{n+1}}{n+1} \right]^q du.$$

*Proof.* From (2.10),

$$(3.12) \quad \left| \sigma^2(X) + (E(X) - x)^2 - \sum_{k=0}^n \frac{(b-x)^{k+3} + (-1)^k (x-a)^{k+3}}{k+3} \cdot \frac{f^{(k)}(x)}{k!} \right|$$

$$= \left| \frac{1}{n!} \int_a^b K_n(x, s) f^{(n+1)}(s) ds \right|.$$

Now, on using the fact that from (2.11), (2.12),  $\psi_n(u, v) \geq 0$  for  $u, v \geq 0$ ,

$$(3.13) \quad \left| \frac{1}{n!} \int_a^b K_n(x, s) f^{(n+1)}(s) ds \right|$$

$$\leq \frac{\|f^{(n+1)}\|_\infty}{n!} \left\{ \int_a^x \psi_n(s-a, x-s) ds + \int_x^b \psi_n(b-s, s-x) ds \right\}.$$

Further,

$$(3.14) \quad \psi_n(u, v) = \frac{u^{n+3}}{n+3} + 2v \frac{u^{n+2}}{n+2} + v^2 \frac{u^{n+1}}{n+1}$$

and so

$$(3.15) \quad \int_a^x \psi_n(s-a, x-s) ds$$

$$= \int_a^x \left[ \frac{(s-a)^{n+3}}{n+3} + 2(x-s) \frac{(s-a)^{n+2}}{n+2} + (x-s)^2 \frac{(s-a)^{n+1}}{n+1} \right] ds$$

$$= (x-a)^{n+4} \int_0^1 \left[ \frac{\lambda^{n+3}}{n+3} + 2(1-\lambda) \frac{\lambda^{n+2}}{n+2} + (1-\lambda)^2 \frac{\lambda^{n+1}}{n+1} \right] d\lambda,$$

where we have made the substitution  $\lambda = \frac{s-a}{x-a}$ .

Collecting powers of  $\lambda$  gives

$$\lambda^{n+3} \left[ \frac{1}{n+3} - \frac{2}{n+2} + \frac{1}{n+1} \right] - \frac{2\lambda^{n+2}}{(n+2)(n+1)} + \frac{\lambda^{n+1}}{n+1}$$

and so, from (3.15),

$$\begin{aligned}
(3.16) \quad & \int_a^x \psi_n(s-a, x-s) ds \\
&= (x-a)^{n+4} \left\{ \frac{1}{n+4} \left[ \frac{1}{n+3} - \frac{2}{n+2} + \frac{1}{n+1} \right] \right. \\
&\quad \left. - \frac{2}{(n+3)(n+2)(n+1)} + \frac{1}{(n+2)(n+1)} \right\} \\
&= \frac{(x-a)^{n+4}}{(n+4)(n+1)}.
\end{aligned}$$

Similarly, on using (3.14),

$$\begin{aligned}
& \int_x^b \psi_n(b-s, s-x) ds \\
&= \int_x^b \left[ \frac{(b-s)^{n+3}}{n+3} + 2(s-x) \frac{(b-s)^{n+2}}{n+2} + (s-x)^2 \frac{(b-s)^{n+1}}{n+1} \right] ds
\end{aligned}$$

and making the substitution  $\nu = \frac{b-s}{b-x}$  gives

$$\begin{aligned}
(3.17) \quad & \int_x^b \psi_n(b-s, s-x) ds \\
&= (b-x)^{n+4} \int_0^1 \left[ \frac{\nu^{n+3}}{n+3} + 2(1-\nu) \frac{\nu^{n+2}}{n+2} + (1-\nu)^2 \frac{\nu^{n+1}}{n+1} \right] d\nu \\
&= \frac{(b-x)^{n+4}}{(n+4)(n+1)},
\end{aligned}$$

where we have used (3.15) and (3.16). Combining (3.16) and (3.17) gives the first inequality in (3.10).

For the second inequality in (3.10), we use Hölder's integral inequality to obtain

$$(3.18) \quad \left| \frac{1}{n!} \int_a^b K_n(x, s) f^{(n+1)}(s) ds \right| \leq \frac{\|f^{(n+1)}(s)\|_p}{n!} \left( \int_a^b |K_n(x, s)|^q ds \right)^{\frac{1}{q}}.$$

Now, from (2.11) and (3.14)

$$\begin{aligned}
\int_a^b |K_n(x, s)|^q ds &= \int_a^x \psi^q(s-a, x-s) ds + \int_x^b \psi^q(b-s, s-x) ds \\
&= C \left[ (x-a)^{(n+3)q+1} + (b-x)^{(n+3)q+1} \right],
\end{aligned}$$

where  $C$  is as defined in (3.11) and we have used (3.15) and (3.16). Substitution into (3.18) gives the second inequality in (3.10).

Finally, for the third inequality in (3.10). From (3.12),

$$\begin{aligned}
(3.19) \quad & \left| \frac{1}{n!} \int_a^b K_n(x, s) f^{(n+1)}(s) ds \right| \\
& \leq \frac{1}{n!} \left\{ \int_a^x \psi_n(s-a, x-s) \left| f^{(n+1)}(s) \right| ds \right. \\
& \quad \left. + \int_x^b \psi_n(b-s, s-x) \left| f^{(n+1)}(s) \right| ds \right\} \\
& \leq \frac{1}{n!} \left\{ \psi_n(x-a, 0) \int_a^x \left| f^{(n+1)}(s) \right| ds + \psi_n(b-x, 0) \int_x^b \left| f^{(n+1)}(s) \right| ds \right\},
\end{aligned}$$

where, from (3.14),

$$(3.20) \quad \psi_n(u, 0) = \frac{u^{n+3}}{n+3}.$$

Hence, from (3.19) and (3.20)

$$\begin{aligned}
& \left| \frac{1}{n!} \int_a^b K_n(x, s) f^{(n+1)}(s) ds \right| \\
& \leq \frac{1}{n!} \max \left\{ \frac{(x-a)^{n+3}}{n+3}, \frac{(b-x)^{n+3}}{n+3} \right\} \left\| f^{(n+1)}(\cdot) \right\|_1 \\
& = \frac{1}{n!(n+3)} [\max\{x-a, b-x\}]^{n+3} \left\| f^{(n+1)}(\cdot) \right\|_1,
\end{aligned}$$

which, on using the fact that for  $X, Y \in \mathbb{R}$

$$\max\{X, Y\} = \frac{X+Y}{2} + \left| \frac{X-Y}{2} \right|$$

gives, from (3.12), the third inequality in (3.10). The theorem is now completely proved. ■

**Remark 1.** *The results of Theorem 3 may be compared with those of Theorem 1. Theorem 3 is based on the single integral identity developed in Lemma 2, while Theorem 1 is based on the double integral identity representation for the bound. It may be noticed from (3.1) and (3.10) that the bounds are the same for  $f^{(n+1)} \in L_\infty[a, b]$ , that for  $f^{(n+1)} \in L_1[a, b]$  the bound obtained in (3.1) is better and for  $f^{(n+1)} \in L_p[a, b]$ ,  $p > 1$ , the result is inconclusive.*

#### REFERENCES

- [1] N.S. BARNETT, P. CERONE, S.S. DRAGOMIR and J. ROUMELIOTIS, Some inequalities for the dispersion of a random variable whose p.d.f. is defined on a finite interval, *submitted for publication*.
- [2] N.S. BARNETT and S.S. DRAGOMIR, An inequality of Ostrowski type for cumulative distribution functions, Accepted for publication in *Kyungpook Math. J.*, 1999.  
[ONLINE] <http://rgmia.vu.edu.au/v1n1.html>
- [3] N.S. BARNETT and S.S. DRAGOMIR, An Ostrowski type inequality for a random variable whose probability density function belongs to  $L_\infty[a, b]$ . *Submitted for publication*.  
[ONLINE] <http://rgmia.vu.edu.au/v1n1.html>

- [4] S.S. DRAGOMIR, N.S. BARNETT and S. WANG, An Ostrowski type inequality for a random variable whose probability density function belongs to  $L_p[a, b]$ ,  $p > 1$ , *Mathematical Inequalities & Applications*, **2** (1999), No. 4, 501-508.

SCHOOL OF COMMUNICATIONS AND INFORMATICS, VICTORIA UNIVERSITY OF TECHNOLOGY, PO  
BOX 14428, MELBOURNE CITY MC, VICTORIA, 8001. AUSTRALIA  
*E-mail address:* {neil, pc, sever, johnr}@matilda.vu.edu.au