



VICTORIA UNIVERSITY
MELBOURNE AUSTRALIA

On the Ostrowski's Integral Inequality for Mappings with Bounded Variation and Applications

This is the Published version of the following publication

Dragomir, Sever S (1999) On the Ostrowski's Integral Inequality for Mappings with Bounded Variation and Applications. RGMIA research report collection, 2 (1).

The publisher's official version can be found at

Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/17133/>

ON THE OSTROWSKI'S INTEGRAL INEQUALITY FOR MAPPINGS WITH BOUNDED VARIATION AND APPLICATIONS

S.S. Dragomir

ABSTRACT. A generalization of Ostrowski's inequality for mappings with bounded variation and applications in Numerical Analysis for Euler's Beta function is given.

1 INTRODUCTION

The following theorem contains the integral inequality which is known in the literature as Ostrowski's inequality [2, p. 469].

Theorem 1.1. *Let $f : [a, b] \rightarrow \mathbf{R}$ be a differentiable mapping on (a, b) whose derivative is bounded on (a, b) and denote $\|f'\|_\infty = \sup_{t \in (a, b)} |f'(t)| < \infty$. Then for all $x \in [a, b]$ we have the inequality*

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left[\frac{1}{4} + \frac{\left(x - \frac{a+b}{2}\right)^2}{(b-a)^2} \right] (b-a) \|f'\|_\infty.$$

The constant $\frac{1}{4}$ is sharp in the sense that it can not be replaced by a smaller one.

In this paper we prove an Ostrowski's type inequality for mappings with bounded variation and apply it in obtaining a Riemann's type quadrature formula for this class of mappings. Applications for Euler's Beta function are also given.

2 OSTROWSKI'S INEQUALITY FOR MAPPINGS WITH BOUNDED VARIATION

The following inequality for mappings with bounded variation holds:

Theorem 2.1. *Let $u : [a, b] \rightarrow \mathbf{R}$ be mapping with bounded variation on $[a, b]$. Then for all $x \in [a, b]$, we have the inequality*

$$(2.1) \quad \left| \int_a^b u(t) dt - u(x)(b-a) \right| \leq \left[\frac{1}{2}(b-a) + \left| x - \frac{a+b}{2} \right| \right] \bigvee_a^b(u).$$

where $\bigvee_a^b(u)$ denotes the total variation of u .
 The constant $\frac{1}{2}$ is the best possible one.

Proof. Using the integration by parts formula for Riemann-Stieltjes integral we have

$$\int_a^x (t-a) du(t) = u(x)(x-a) - \int_a^x u(t) dt$$

Date. January, 1999

1991 Mathematics Subject Classification. Primary 26 D 15; Secondary 41 A 55.

Key words and phrases. Ostrowski's Inequality, Numerical Integration, Beta Mapping.

and

$$\int_x^b (t-b) du(t) = u(x)(b-x) - \int_x^b u(t) dt.$$

If we add the above two equalities, we get

$$(2.2) \quad u(x)(b-a) - \int_a^b u(t) dt = \int_a^b p(x, t) du(t)$$

where

$$p(x, t) := \begin{cases} t-a & \text{if } t \in [a, x) \\ t-b & \text{if } x \in [x, b], \end{cases}$$

for all $x, t \in [a, b]$.

Now, assume that $\Delta_n : a = x_0^{(n)} < x_1^{(n)} < \dots < x_{n-1}^{(n)} < x_n^{(n)} = b$ is a sequence of divisions with $\nu(\Delta_n) \rightarrow 0$ as $n \rightarrow \infty$, where $\nu(\Delta_n) := \max_{i \in \{0, \dots, n-1\}} (x_{i+1}^{(n)} - x_i^{(n)})$ and $\xi_i^{(n)} \in [x_i^{(n)}, x_{i+1}^{(n)}]$.

If $p : [a, b] \rightarrow \mathbf{R}$ is continuous on $[a, b]$ and $v : [a, b] \rightarrow \mathbf{R}$ is with bounded variation on $[a, b]$, then

$$(2.3) \quad \left| \int_a^b p(x) dv(x) \right| = \left| \lim_{\nu(\Delta_n) \rightarrow 0} \sum_{i=0}^{n-1} p(\xi_i^{(n)}) [v(x_{i+1}^{(n)}) - v(x_i^{(n)})] \right|$$

$$\leq \lim_{\nu(\Delta_n) \rightarrow 0} \sum_{i=0}^{n-1} |p(\xi_i^{(n)})| |v(x_{i+1}^{(n)}) - v(x_i^{(n)})|$$

$$\leq \sup_{x \in [a, b]} |p(x)| \sup_{\Delta_n} \sum_{i=0}^{n-1} |v(x_{i+1}^{(n)}) - v(x_i^{(n)})| = \sup_{x \in [a, b]} |p(x)| \bigvee_a^b(v).$$

Applying the inequality (2.3) for $p(x, t)$ as above and $v(x) = u(x)$, $x \in [a, b]$, we get

$$(2.4) \quad \left| \int_a^b p(x, t) du(t) \right| \leq \sup_{t \in [a, b]} |p(x, t)| \bigvee_a^b(u)$$

$$= \max\{x-a, b-x\} \bigvee_a^b(u) = \left[\frac{b-a}{2} + \left| x - \frac{a+b}{2} \right| \right] \bigvee_a^b(u)$$

and then by (2.4), via the identity (2.2), we deduce the desired inequality (2.1).

Now, assume that the inequality (2.1) holds with a constant $C > 0$, i.e.,

$$(2.5) \quad \left| \int_a^b u(t) dt - u(x)(b-a) \right| \leq \left[C(b-a) + \left| x - \frac{a+b}{2} \right| \right] \bigvee_a^b(u).$$

for all $x \in [a, b]$.

Consider the mapping $u : [a, b] \rightarrow \mathbf{R}$, given by

$$u(x) = \begin{cases} 0 & \text{if } x \in [a, b] \setminus \left\{ \frac{a+b}{2} \right\} \\ 1 & \text{if } x = \frac{a+b}{2} \end{cases}$$

in (2.5). Then u is with bounded variation on $[a, b]$, and

$$\bigvee_a^b(u) = 2, \quad \int_a^b u(t) dt = 0$$

and for $x = \frac{a+b}{2}$, we get in (2.5)

$$1 \leq 2C$$

which implies that $C \geq \frac{1}{2}$ and the theorem is completely proved.

The following corollary holds:

Corollary 2.2. *Let $u : [a, b] \rightarrow \mathbf{R}$ be a monotonous mapping on $[a, b]$. Then we have the inequality*

$$\left| \int_a^b u(t) dt - u(x)(b-a) \right| \leq \left[\frac{1}{2}(b-a) + \left| x - \frac{a+b}{2} \right| \right] |f(b) - f(a)|.$$

The case of lipschitzian mappings is embodied in the following corollary.

Corollary 2.3. *Let $u : [a, b] \rightarrow \mathbf{R}$ be an L -lipschitzian mapping on $[a, b]$, i.e., we recall*

$$|u(x) - u(y)| \leq L |x - y| \quad \text{for all } x, y \in [a, b].$$

Then we have the inequality

$$\left| \int_a^b u(t) dt - u(x)(b-a) \right| \leq L \left[\frac{1}{2}(b-a) + \left| x - \frac{a+b}{2} \right| \right] (b-a).$$

The best inequality we can get from (2.1) is that one for which $x = \frac{a+b}{2}$ obtaining

Corollary 2.4. *Let $u : [a, b] \rightarrow \mathbf{R}$ be as above. Then we have the inequality:*

$$(2.6) \quad \left| \int_a^b u(t) dx - u\left(\frac{a+b}{2}\right)(b-a) \right| \leq \frac{1}{2}(b-a) \bigvee_a^b(u).$$

Similar inequalities can be found if we assume that u is monotonous or lipschitzian on $[a, b]$. We shall omit the details.

Remark 2.1. *If we assume that u is continuous differentiable on (a, b) and u' is integrable on (a, b) , then by (2.1) we get*

$$\left| \int_a^b u(t) dx - u\left(\frac{a+b}{2}\right)(b-a) \right| \leq \left[\frac{1}{2}(b-a) + \left| x - \frac{a+b}{2} \right| \right] \|u'\|_1$$

which is the inequality obtained by Dragomir and Wang in the recent paper [1].

Remark 2.2. *It is well known that if $f : [a, b] \rightarrow \mathbf{R}$ is a convex mapping on $[a, b]$, then Hermite-Hadamard's inequality holds*

$$(2.7) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2}.$$

Now, if we assume that $f : I \subset \mathbf{R} \rightarrow \mathbf{R}$ is convex on I and $a, b \in \text{Int}(I)$, $a < b$; then f'_+ is monotonous nondecreasing on $[a, b]$ and by Corollary 2.4 we get

$$(2.8) \quad 0 \leq \frac{1}{b-a} \int_a^b f(x) dx - f\left(\frac{a+b}{2}\right) \leq \frac{1}{2} \|f'_+\|_1$$

which gives a counterpart for the first membership of Hadamard's inequality.

Similar results can be obtained if we assume that f is convex and monotonous or convex and lipschitzian on $[a, b]$.

3 A QUADRATURE FORMULA OF RIEMANN TYPE

Let $I_n : a = x_0 < x_1 < \dots < x_{n-1} < x_n = b$ be a division of the interval $[a, b]$ and $\xi_i \in [x_i, x_{i+1}]$ ($i = 0, \dots, n-1$) a sequence of intermediate points for I_n . Construct the Riemann sums

$$R_n(f, I_n, \xi) = \sum_{i=0}^{n-1} f(\xi_i) h_i$$

where $h_i := x_{i+1} - x_i$.

We have the following quadrature formula

Theorem 3.1. *Let $f : [a, b] \rightarrow \mathbf{R}$ be a mapping with bounded variation on $[a, b]$ and I_n, ξ_i ($i = 0, \dots, n-1$) be as above. Then we have the Riemann quadrature formula*

$$(3.1) \quad \int_a^b f(x) dx = R_n(f, I_n, \xi) + W_n(f, I_n, \xi) \quad (3.1)$$

where the remainder satisfies the estimation

$$(3.2) \quad |W_n(f, I_n, \xi)| \leq \sup_{i=0, \dots, n} \left[\frac{1}{2} h_i + \left| \xi_i - \frac{x_i + x_{i+1}}{2} \right| \right] \bigvee_a^b(f)$$

$$\leq \left[\frac{1}{2} \nu(h) + \sup_{i=0, \dots, n} \left| \xi_i - \frac{x_i + x_{i+1}}{2} \right| \right] \bigvee_a^b(f) \leq \nu(h) \bigvee_a^b(f) \quad (3.2)$$

for all ξ_i ($i = 0, \dots, n-1$) as above, where $\nu(h) := \max_{i=0, \dots, n} h_i$.

The constant $\frac{1}{2}$ is sharp in (3.2).

Proof. Apply Theorem 2.1 on the interval $[x_i, x_{i+1}]$ to get

$$(3.3) \quad \left| \int_{x_i}^{x_{i+1}} f(x) dx - f(\xi_i) h_i \right| \leq \left[\frac{1}{2} h_i + \left| \xi_i - \frac{x_i + x_{i+1}}{2} \right| \right] \bigvee_{x_i}^{x_{i+1}}(f).$$

Summing over i from 0 to $n-1$ and using the generalized triangle inequality we get

$$\begin{aligned} |W_n(f, I_n, \xi)| &\leq \sum_{i=0}^{n-1} \left| \int_{x_i}^{x_{i+1}} f(x) dx - f(\xi_i) h_i \right| \\ &\leq \sum_{i=0}^{n-1} \left[\frac{1}{2} h_i + \left| \xi_i - \frac{x_i + x_{i+1}}{2} \right| \right] \bigvee_{x_i}^{x_{i+1}}(f) \\ &\leq \sup_{i=0, \dots, n} \left[\frac{1}{2} h_i + \left| \xi_i - \frac{x_i + x_{i+1}}{2} \right| \right] \sum_{i=0}^{n-1} \bigvee_{x_i}^{x_{i+1}}(f). \\ &= \sup_{i=0, \dots, n} \left[\frac{1}{2} h_i + \left| \xi_i - \frac{x_i + x_{i+1}}{2} \right| \right] \bigvee_a^b(f) \end{aligned}$$

The second inequality follows by the properties of $\sup(\cdot)$.

Now, as

$$\left| \xi_i - \frac{x_i + x_{i+1}}{2} \right| \leq \frac{1}{2} h_i$$

for all $\xi_i \in [x_i, x_{i+1}]$ ($i = 0, \dots, n-1$) the last part of (3.2) is also proved.

Corollary 3.2. Let $u : [a, b] \rightarrow \mathbf{R}$ be a monotonous mapping on $[a, b]$ and I_n, ξ_i ($i = 0, \dots, n-1$) be as above. Then we have the Riemann quadrature formula (3.1) and the remainder satisfies the estimation

$$\begin{aligned} |W_n(f, I_n, \xi)| &\leq \sup_{i=0, \dots, n} \left[\frac{1}{2} h_i + \left| \xi_i - \frac{x_i + x_{i+1}}{2} \right| \right] |f(b) - f(a)| \\ &\leq \left[\frac{1}{2} \nu(h) + \sup_{i=0, \dots, n} \left| \xi_i - \frac{x_i + x_{i+1}}{2} \right| \right] |f(b) - f(a)| \leq \nu(h) |f(b) - f(a)| \end{aligned}$$

for all ξ_i ($i = 0, \dots, n-1$) as above.

The case of lipschitzian mappings is embodied into the following corollary.

Corollary 3.3. Let $u : [a, b] \rightarrow \mathbf{R}$ be an L -lipschitzian mapping on $[a, b]$ and I_n, ξ_i ($i = 0, \dots, n-1$) be as above. Then we have the Riemann quadrature formula (3.1) and the remainder satisfies the estimation

$$\begin{aligned} |W_n(f, I_n, \xi)| &\leq L \sum_{i=0}^{n-1} \left[\frac{1}{2} h_i + \left| \xi_i - \frac{x_i + x_{i+1}}{2} \right| \right] h_i \\ &\leq L \sum_{i=0}^{n-1} h_i^2 \end{aligned}$$

The proof is obvious by Corollary 2.3 applied on the intervals $[x_i, x_{i+1}]$ and summing the obtained inequalities.

We shall omit the details.

Note that the best estimation we can get from (3.2) is that one for which $\xi_i = \frac{x_i + x_{i+1}}{2}$ obtaining the following midpoint formula:

Corollary 3.4. Let f, I_n be as Theorem 3.1. Then we have the midpoint rule

$$\int_a^b f(x) dx = M_n(f, I_n) + S_n(f, I_n)$$

where

$$M_n(f, I_n) = \sum_{i=0}^{n-1} f\left(\frac{x_i + x_{i+1}}{2}\right) h_i$$

and the remainder $S_n(f, I_n)$ satisfies the estimation

$$|S_n(f, I_n)| \leq \frac{1}{2} \nu(h) \bigvee_a^b(f).$$

Similar results can be obtained from Corollaries 3.2 and 3.3.

Remark 3.1. If we assume that $f : [a, b] \rightarrow \mathbf{R}$ is differentiable on (a, b) and whose derivative f' is integrable on (a, b) we can put instead of $\bigvee_a^b(f)$ the L_1 -norm $\|f'\|_1$ obtaining the estimation due to Dragomir-Wang from the paper [1].

4 APPLICATIONS FOR EULER'S BETA MAPPING

Consider the mapping Beta for real numbers

$$B(p, q) := \int_0^1 t^{p-1} (1-t)^{q-1} dt, \quad p, q > 0$$

and the mapping $e_{p,q}(t) := t^{p-1} (1-t)^{q-1}$, $t \in [0, 1]$.

We have for $p, q > 1$ that

$$e'_{p,q}(t) = e_{p-1,q-1}(t)[p-1-(p+q-2)t]$$

and as

$$|p-1-(p+q-2)t| \leq \max\{p-1, q-1\}$$

for all $t \in [0, 1]$, then

$$\begin{aligned} (4.1) \quad \|e'_{p,q}\|_1 &\leq \max\{p-1, q-1\} \|e_{p-2,q-2}\|_1 \\ &= \max\{p-1, q-1\} B(p-1, q-1); \quad p, q > 1. \end{aligned}$$

The following inequality for Beta mapping holds

Proposition 4.1. *Let $p, q > 1$ and $x \in [0, 1]$. Then we have the inequality*

$$\begin{aligned} (4.2) \quad &|B(p, q) - x^{p-1} (1-x)^{q-1}| \\ &\leq \max\{p-1, q-1\} B(p-1, q-1) \left[\frac{1}{2} \left| x - \frac{1}{2} \right| \right]. \end{aligned}$$

The proof follows by Theorem 2.1 applied for the mapping $e_{p,q}$ and taking into account that $\|e'_{p,q}\|_1$ satisfies the inequality (4.1).

Corollary 4.2. *Let $p, q > 1$. Then we have the inequality*

$$\left| B(p, q) - \frac{1}{2^{p+q-2}} \right| \leq \frac{1}{2} \max\{p-1, q-1\} B(p-1, q-1).$$

Now, if we apply Theorem 3.1 for the mapping $e_{p,q}$ we get the following approximation of Beta mapping in terms of Riemann sums.

Proposition 4.3. *Let $I_n : a = x_0 < x_1 < \dots < x_{n-1} < x_n = b$ be a division of the interval $[a, b]$, $\xi_i \in [x_i, x_{i+1}]$ ($i = 0, \dots, n-1$) a sequence of intermediate points for I_n and $p, q > 1$. Then we have the formula*

$$B(p, q) = \sum_{i=0}^{n-1} \xi_i^{p-1} (1-\xi_i)^{q-1} h_i + T_n(p, q)$$

where the remainder $T_n(p, q)$ satisfies the estimation

$$\begin{aligned} &|T_n(p, q)| \\ &\leq \max\{p-1, q-1\} \left[\frac{1}{2} \nu(h) + \sup_{i=0, \dots, n} \left| \xi_i - \frac{x_i + x_{i+1}}{2} \right| \right] B(p-1, q-1) \\ &\leq \max\{p-1, q-1\} \nu(h) B(p-1, q-1). \end{aligned}$$

Particularly, if we choose above $\xi_i = \frac{x_i + x_{i+1}}{2}$ ($i = 0, \dots, n-1$) then we get the approximation

$$B(p, q) = \frac{1}{2^{p+q-2}} \sum_{i=0}^{n-1} (x_i + x_{i+1})^{p-1} (2 - x_i - x_{i+1})^{q-1} + V_n(p, q)$$

where

$$|V_n(p, q)| \leq \frac{1}{2} \max\{p-1, q-1\} \nu(h) B(p-1, q-1).$$

REFERENCES

- [1] S.S. DRAGOMIR and S. WANG, A new inequality of Ostrowski's type in L_1 norm and applications to some special means and to some numerical quadrature rules, *Tamkang J. of Math.*, **28**(1997), 239-244.
- [2] D.S. MITRINOVIĆ, J.E. PEČARIĆ, and A.M. FINK, *Inequalities for Functions and their Integrals and Derivatives*, Kluwer Academic Publishers, 1994.

SCHOOL OF COMMUNICATIONS AND INFORMATICS, VICTORIA UNIVERSITY OF TECHNOLOGY, PO Box 14428,
MELBOURNE CITY MC, VICTORIA 8001, AUSTRALIA.

E-mail address: `sever@matilda.vut.edu.au`

