



VICTORIA UNIVERSITY
MELBOURNE AUSTRALIA

A Variational Characterization of Reflexivity and Strict Convexity

This is the Published version of the following publication

Dragomir, Sever S (1999) A Variational Characterization of Reflexivity and Strict Convexity. RGMIA research report collection, 2 (6).

The publisher's official version can be found at

Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/17252/>

A VARIATIONAL CHARACTERIZATION OF REFLEXIVITY AND STRICT CONVEXITY

S. S. DRAGOMIR

ABSTRACT. In this paper we give a variational characterization of reflexivity and strict convexity which is related to James and Krein theorems in Geometry of Banach spaces.

1. INTRODUCTION

Let $(X, \|\cdot\|)$ be a real normed space and consider the norm derivatives

$$(x, y)_{i(s)} = \lim_{t \rightarrow -(+)0} \frac{(\|y + tx\|^2 - \|y\|^2)}{2t}.$$

Note that these mappings are well defined on $X \times X$ and the following properties are valid (see also [1]-[5]):

- (i) $(x, y)_i = -(-x, y)_s$ if x, y are in X ;
- (ii) $(x, x)_p = \|x\|^2$ for all x in X ;
- (iii) $(\alpha x, \beta y)_p = \alpha\beta (x, y)_p$ for all x, y in X and $\alpha\beta \geq 0$;
- (iv) $(\alpha x + y, x)_p = \alpha\|x\|^2 + (y, x)_p$ for all x, y in X and α a real number;
- (v) $(x + y, z)_p \leq \|x\| \cdot \|z\| + (y, z)_p$ for all x, y, z in X ;
- (vi) The element x in X is Birkhoff orthogonal over y in X (we denote this by $x \perp y$), i.e., $\|x + ty\| \geq \|x\|$ for all t a real number iff $(y, x)_i \leq 0 \leq (y, x)_s$;
- (vii) The space X is smooth iff $(y, x)_i = (y, x)_s$ for all x, y in X iff $(\cdot, \cdot)_p$ is linear in the first variable;

where $p = s$ or $p = i$.

The following characterization of reflexivity is well known (see [6]):

Theorem 1. (James). *The Banach space X is reflexive iff for any continuous linear functional $f : X \rightarrow \mathbb{R}$ there exists an element u_f in X such that $f(u_f) = \|f\| \cdot \|u_f\|$, i.e., u_f is a maximal element for f .*

The following characterization of strict convexity in terms of maximal elements is well known and is due to M. G. Krein ([7, p. 27]):

Theorem 2. (Krein). *The real Banach space X is strictly convex iff any nonzero continuous linear functional defined on it has at most one maximal element having a norm equal to one.*

Date: June, 1999.

1991 Mathematics Subject Classification. Primary 46B20; Secondary 41Xxx.

Key words and phrases. James theorem, Krein's theorem, Reflexivity, Strict convexity.

2. THE RESULTS

We give here a variational characterization of reflexivity and strict convexity as follows.

Theorem 3. *Let $(X, \|\cdot\|)$ be a real Banach space. The following statements are equivalent:*

- (i) X is reflexive [strictly convex (reflexive and strictly convex)];
- (ii) For any nonzero continuous linear functional $f : X \rightarrow \mathbb{R}$ there exists at least one [at most one (a unique)] vector $u_f \in X$, $\|u_f\| = 1$ which minimizes the quadratic functional $F_f : X \rightarrow \mathbb{R}$, $F_f(x) = \|x\|^2 - \frac{2f(x)}{\|f\|}$.

Proof. “(i) \Rightarrow (ii)” a). Assume that X is reflexive and let $f \in X^* \setminus \{0\}$. Then by James’ theorem there exists a vector $u_f \in X$, $\|u_f\| = 1$ such that $f(u_f) = \|f\|$. However,

$$\|u_f\| = 1 = \frac{f(u_f)}{\|f\|} = \frac{f(u_f + \lambda u)}{\|f\|} \leq \|u_f + \lambda u\|$$

for all $\lambda \in \mathbb{R}$ and $u \in \text{Ker}(f)$, which gives us that $u_f \perp \text{Ker}(f)$.

Let $x \in X$ be arbitrary but fixed and define $y := f(x)u_f - f(u_f)x$. As $f(y) = 0$, we get that $y \in \text{Ker}(f)$ and then $u_f \perp y$ in Birkhoff’s sense. By the property (vi) we get that

$$(2.1) \quad (y, x)_i \leq 0 \leq (y, x)_s$$

which is equivalent with

$$(f(x)u_f - f(u_f)x, u_f)_i \leq 0 \leq (f(x)u_f - f(u_f)x, u_f)_s \text{ for all } x \in X.$$

Using the properties of semi-inner products we get

$$(f(x)u_f - f(u_f)x, u_f)_i = f(x) - \|f\|(x, u_f)_s$$

and

$$(f(x)u_f - f(u_f)x, u_f)_s = f(x) - \|f\|(x, u_f)_i$$

for all $x \in X$, and then by (2.1) we get that

$$(2.2) \quad \|f\|(x, u_f)_i \leq f(x) \leq \|f\|(x, u_f)_s \text{ for all } x \in X.$$

We shall prove now that u_f minimizes the quadratic functional F_f .

Let $u \in X$. Then, as $f(u_f) = \|f\|$ and $\|u_f\| = 1$, we get that

$$\begin{aligned} F_f(u) - F_f(u_f) &= \|u\|^2 - \frac{2f(u)}{\|f\|} - \|u_f\|^2 + \frac{2f(u_f)}{\|f\|} \\ &= \|u\|^2 - 2\frac{f(u)}{\|f\|} + \|u_f\|^2. \end{aligned}$$

By (2.2) we have that

$$\frac{-2f(u)}{\|f\|} \geq -2(x, u_f)_s$$

and then

$$\begin{aligned} F_f(u) - F_f(u_f) &\geq \|u\|^2 - 2(x, u_f)_s + \|u_f\|^2 \\ &\geq \|u\|^2 - 2\|u\| \cdot \|u_f\| + \|u_f\|^2 \\ &= (\|u\| - \|u_f\|)^2 \geq 0 \end{aligned}$$

which shows that u_f minimizes F_f .

“(ii) \Rightarrow (i)” a). Let $f \in X^* \setminus \{0\}$ and u_f be an element minimizing F_f . Then, for all $\lambda \in \mathbb{R}$ and $u \in X$, we have:

$$(2.3) \quad F_f(u + \lambda u_f) \geq F_f(u_f).$$

However,

$$\begin{aligned} F_f(u + \lambda u_f) - F_f(u_f) &= \|u + \lambda u_f\|^2 - \frac{2f(u + \lambda u_f)}{\|f\|} - \|u_f\|^2 + \frac{2f(u_f)}{\|f\|} \\ &= \|u + \lambda u_f\|^2 - \|u_f\|^2 - \frac{2\lambda f(u)}{\|f\|} \end{aligned}$$

and then (2.3) is equivalent to

$$\frac{2\lambda f(u)}{\|f\|} \leq \|u + \lambda u_f\|^2 - \|u_f\|^2 \text{ for all } \lambda \in \mathbb{R} \text{ and } u \in X.$$

Assume that $\lambda > 0$. Then

$$f(u) \leq \left[\frac{(\|u + \lambda u_f\|^2 - \|u_f\|^2)}{2\lambda} \right] \cdot \|f\|.$$

Letting $\lambda \rightarrow 0+$, we get

$$f(u) \leq \|f\| (u, u_f)_s$$

for all $u \in X$. Now, changing u with $(-u)$, we get from the previous inequality that

$$f(u) \geq -\|f\| (-u, u_f)_s = \|f\| (u, u_f)_i$$

and then we get the estimation

$$\|f\| (u, u_f)_i \leq f(u) \leq \|f\| (u, u_f)_s \text{ for all } u \in X.$$

Choosing $u = u_f$ we get $f(u_f) = \|f\|$ and by James' theorem it follows that $(X, \|\cdot\|)$ is reflexive.

“(i) \Rightarrow (ii)” b). Assume that there exists a nonzero functional $f_0 \in X^*$ for which we can find at least two distinct vectors

$$u_{f_0}^i (i = 1, 2), \|u_{f_0}^i\| = 1$$

which minimize F_{f_0} . As above (see “(ii) \Rightarrow (i)” a.), we get that $f_0(u_{f_0}^i) = \|f_0\|$, which, by Krein's theorem contradicts the strict convexity of X .

“(ii) \Rightarrow (i)” b). Assume that X is not reflexive. Thus, by Krein's theorem, there exists a continuous linear functional $f_0 \neq 0$ and at least two distinct elements

$$u_{f_0}^i (i = 1, 2), \|u_{f_0}^i\| = 1$$

such that $f_0(u_{f_0}^i) = \|f_0\|$. Now, by a similar argument as in “(i) \Rightarrow (ii)” a.), we deduce that $u_{f_0}^i (i = 1, 2)$ will minimize the quadratic functional F_{f_0} , which is a contradiction. ■

REFERENCES

- [1] J. Diestel: *Geometry of Banach Spaces - Selected Topics*, Springer Verlag, Berlin, Heidelberg, New York, 485, 1975.
- [2] S. S. Dragomir: On continuous sublinear functionals in reflexive Banach spaces and applications. *Riv. Mat. Univ. Parma*, **16**, 239-250 (1990).
- [3] S.S. Dragomir: Approximation of continuous linear functionals in real normed spaces. *Rendiconti di Matematica, Serie VII, Roma*, **12**, 357-364 (1992).
- [4] S. S. Dragomir: The approximation of continuous linear functionals, *Extracta Math.*, **6**, 50-51(1991).
- [5] S. S. Dragomir: A characterization of reflexivity, *Proc. Japan Acad.*, *72, Ser A.* 152-153(1996).
- [6] R.C. James: Characterizations of reflexivity, *Studia Math.*, **23**(1964), 205-216
- [7] V.I. Istratescu: *Strict Convexity and Complex Strict Convexity*, Marcel Dekker Inc., New York and Basel, 1984.

SCHOOL OF COMMUNICATIONS AND INFORMATICS, VICTORIA UNIVERSITY OF TECHNOLOGY, PO BOX 14428, MELBOURNE CITY, MC 8001, VICTORIA, AUSTRALIA

E-mail address: `sever@matilda.edu.au`

URL: `http://rgmia.vu.edu.au`