



VICTORIA UNIVERSITY
MELBOURNE AUSTRALIA

Characterizations of Stability for Strongly Continuous Semigroups by Convolutions

This is the Published version of the following publication

Buse, Constantin, Dragomir, Sever S and Lupulescu, V (1999)
Characterizations of Stability for Strongly Continuous Semigroups by
Convolutions. RGMIA research report collection, 2 (7).

The publisher's official version can be found at

Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/17277/>

CHARACTERIZATIONS OF STABILITY FOR STRONGLY CONTINUOUS SEMIGROUPS BY CONVOLUTIONS

C. Buşe, S. S. Dragomir and V. Lupulescu

Abstract

Let $\mathbf{T} = \{T(t)\}_{t \geq 0}$ be a strongly continuous semigroup of bounded linear operators acting on a Banach space X . We prove that if the convolution $\mathbf{T} * (e^{-i\mu(\cdot)} f)$ is bounded for every continuous and 1-periodic function which is null in $t = 0$ and some $\mu \in \mathbf{R}$, then $T(1)$ is power bounded and $e^{i\mu} \in \rho(T(1))$. Applications to questions of exponential stability are also presented.

1. Introduction A well-known result of M.G. Krein, see [K] or [DK], says:

Let X be a Banach space and A be a linear and bounded operator acting on X . If for all $\mu \in \mathbf{R}$ and every $x_0 \in X$ the solution of the Cauchy problem

$$\dot{x}(t) = Ax(t) + e^{i\mu t} x_0, \quad x(0) = 0, \quad t \geq 0$$

is bounded, then there exists constants $N > 0$ and $\nu > 0$ such that

$$\|e^{tA}\| \leq Ne^{-\nu t}, \quad \forall t \geq 0.$$

A proof of this classic result can be found in [Ba]. The above result cannot be extended for the case when A is the infinitesimal generator of a strong semigroup, cf [RB, Example 3.1]. However, weakly related results, hold. For example, in [VuS] (Corollary 4.5 and its reformulation), it has been proved that if the Cauchy problem

$$\dot{x}(t) = Ax(t) + f(t), \quad t \geq 0, \quad x(0) = 0 \quad (A, f, 0)$$

has a bounded solution for every $f \in \mathcal{P}(\omega)$ (that is, f is a continuous and ω -periodic function) then $1 \in \rho(T(\omega))$. Moreover the semigroup \mathbf{T} is uniformly exponentially stable if and only if for every $f \in BUC(\mathbf{R}_+, X)$ (or $f \in AP(\mathbf{R}_+, X)$) the solution of the problem $(A, f, 0)$, is bounded. A short history about the problematic exposed previously and other references can

be found in [VuS]. We present in the following simple generalizations of the above results. Our proofs are elementary and only use first principles.

2. Preliminary results. Let X be a real or complex Banach space and $L(X)$ the Banach algebra of all linear and bounded operators acting on X . We denote by $\|\cdot\|$, the norms of vectors and operators. Let $T \in L(X)$. We will denote by $\sigma(T)$ the spectrum of T and with $r(T)$ we denote the spectral radius of T . We recall that

$$r(T) = \sup\{|z| : z \in \sigma(T)\}. \quad (1)$$

The resolvent set of T is $\rho(T) := \mathbf{C} \setminus \sigma(T)$, i.e. the set of all complex scalar λ such that $\lambda Id - T$ is an invertible operator. Id denotes here the identity operator in $L(X)$. We recall that an operator $T \in L(X)$ is power bounded if there exists an $M > 0$ such that

$$\|T^n\| \leq M, \quad \forall n \in \mathbf{N} = \{0, 1, 2, \dots\}.$$

We shall prove several lemmas which would be used later.

Lemma 1 *Let $T \in L(X)$. If there exist $M > 0$ such that*

$$\sup\{\|Id + T + \dots + T^n\| : n \in \{1, 2, \dots\}\} = M < \infty \quad (2)$$

then T is power bounded and $1 \in \rho(T)$.

Proof. The first assertion follows from (2) and the identity

$$T^{n+1} = Id + (T - Id)(Id + T + \dots + T^n).$$

Suppose that $1 \in \sigma(T)$. Then there exists a sequence $(x_m)_{m \in \mathbf{N}}$ with $x_m \in X$, $\|x_m\| = 1$ and $(Id - T)x_m \rightarrow 0$ as $m \rightarrow \infty$, (see [Na, Proposition 2.2, p. 64]). However, T is power bounded, and hence $T^k(Id - T)x_m \rightarrow 0$ as $m \rightarrow \infty$, uniformly for $k \in \mathbf{N}$. Let $N \in \mathbf{N}$, $N > 2M$ and $m \in \mathbf{N}$ such that

$$\|T^k(Id - T)x_m\| \leq \frac{1}{2N}, \quad k = 0, 1, \dots, N.$$

Then

$$\begin{aligned} M &\geq \|x_m + \sum_{k=1}^N (x_m + \sum_{j=0}^{k-1} T^j(T - Id)x_m)\| \\ &= \|(N+1)x_m + \sum_{k=1}^N \sum_{j=0}^{k-1} T^j(T - Id)x_m\| \\ &\geq (N+1) - \frac{N(N+1)}{4N} > \frac{N}{2} > M. \end{aligned}$$

This is a contradiction and thus $1 \notin \sigma(T)$.

Lemma 2 *Let $U \in L(X)$ and $\mu \in \mathbf{R}$. If*

$$\sup_{n=1,2,\dots} \left\{ \left\| \sum_{k=0}^n e^{i\mu k} U^k \right\| \right\} = M_\mu < \infty \quad (3)$$

then U is power bounded and $e^{-i\mu} \in \rho(U)$.

Proof. It follows from Lemma 1 for $T = e^{i\mu}U$.

Lemma 3 *Let $U \in L(X)$. If the condition (3) holds for all $\mu \in \mathbf{R}$, then $r(T) < 1$.*

Proof. It follows by (1), Lemma 2 and the fact that $\sigma(T)$ is a compact set.

3. Exponential stability and convolutions

We recall that a strongly continuous semigroup is a family $\mathbf{T} = \{T(t)\}_{t \geq 0}$ of bounded linear operators acting on the Banach space X which satisfies the following conditions:

- (i) $T(t+s) = T(t)T(s)$ for all $t, s \in \mathbf{R}_+ := [0, \infty)$;
- (ii) $T(0) = Id$;
- (iii) the function $t \mapsto T(t)x : \mathbf{R}_+ \rightarrow X$ is continuous on \mathbf{R}_+ for all $x \in X$.

The semigroups theory is developed in the books [Pa], [vC], [Na], [Ne] and others.

Let $P_1^0(\mathbf{R}_+, X)$ be the set of all continuous X -valued functions such that $f(t+1) = f(t)$ for all $t \geq 0$ and $f(0) = 0$.

Proposition 4 *Let $\mathbf{T} = \{T(t)\}_{t \geq 0}$ be a strongly continuous semigroup on X and $\mu \in \mathbf{R}$. If*

$$\sup_{t>0} \left\| \int_0^t e^{i\mu\xi} T(t-\xi) f(\xi) d\xi \right\| < \infty,$$

for all $f \in P_1^0(\mathbf{R}_+, X)$, (4)

then $T(1)$ is power bounded and $e^{i\mu} \in \rho(T(1))$.

Proof. Let $U = T(1)$, $x \in X$ and $f_1 \in P_1^0(\mathbf{R}_+, X)$, the function defined by

$$f_1(\xi) = \xi(1 - \xi)T(\xi)x,$$

for all $\xi \in [0, 1]$. From (4), it follows that

$$\sup_{n \in \{1, 2, \dots\}} \left\| \sum_{k=0}^n \int_k^{k+1} T(n+1-\xi)e^{-i\mu\xi} f_1(\xi) d\xi \right\| = M(\mu, f_1) < \infty. \quad (5)$$

Simple calculus gives

$$\begin{aligned} & \int_k^{(k+1)} T(n+1-\xi)e^{-i\mu\xi} f_1(\xi) d\xi \\ &= e^{i\mu(n+1)} \left(\int_0^1 e^{-i\mu\xi} \xi(1-\xi) d\xi \right) e^{-i\mu(n-k+1)} T(n-k+1)x. \end{aligned}$$

Substituting this into (4), we obtain:

$$\sup_{n \in \mathbf{N}} \left\| \sum_{j=1}^{n+1} e^{-\mu j} U^j \right\| < \infty.$$

Now, from Lemma 2, it follows that $T(1)$ is power bounded and $e^{i\mu} \in \rho(T(1))$.

Corollary 5 *Let $\mathbf{T} = \{T(t)\}_{t \geq 0}$ be a strongly continuous semigroup on the Banach space X . If the condition (4) holds for all $\mu \in \mathbf{R}$ and every $f \in P_1^0(\mathbf{R}_+, X)$, then $r(T(1)) < 1$ and \mathbf{T} is uniformly exponentially stable.*

Proof. We recall that a strongly continuous semigroup on X is uniformly exponentially stable if its growth bound $\omega_0(\mathbf{T})$ is negative (or, equivalently) if there exists the constants $N > 0$ and $\nu > 0$ such that

$$\|T(t)\| \leq N e^{-\nu t} \quad \forall t \geq 0.$$

The above assertion follows from Proposition 4, Lemma 3 and the fact that $r(T(1)) = e^{\omega_0(\mathbf{T})}$, cf. [Ne, Proposition 1.2.2].

Let $BUC(\mathbf{R}_+, X)$ the Banach space of all X -valued, bounded and uniformly continuous functions on \mathbf{R}_+ , endowed with the sup-norm and $AP(\mathbf{R}_+, X)$

the space of almost periodic functions in the sense of Bohr, i.e. $AP(\mathbf{R}_+, X)$ is the linear closed hull in $BUC(\mathbf{R}_+, X)$ of the set of all functions:

$$\{e^{i\mu(\cdot)}x : \mu \in \mathbf{R} \quad x \in X\}.$$

Let $AP_0(\mathbf{R}_+, X)$ be the set of all $f \in AP(\mathbf{R}_+, X)$ such that $f(0) = 0$. It is clear that $AP_0(\mathbf{R}_+, X)$ is a closed subspace of either $AP(\mathbf{R}_+, X)$ or $BUC(\mathbf{R}_+, X)$.

Corollary 6 *Let $\mathbf{T} = \{T(t)\}_{t \geq 0}$ be a strongly continuous semigroup on X . If*

$$\sup_{t \geq 0} \left\| \int_0^t T(\xi) f(t - \xi) d\xi \right\| < \infty, \quad \$$$

for all $f \in AP_0(\mathbf{R}_+, X)$ then \mathbf{T} is uniformly exponentially stable.

References

- [Ba] Balint, S., *On the Perron-Bellman theorem for systems with constants coefficients*, Ann. Univ. Timișoara, **21**, fasc. 1-2(1983), 3-8.
- [vC] J. Van Casteren, *Generators of Strongly Continuous Semigroups*, Pitman, Boston-London-Melbourne, 1985.
- [DK] Daletckii Ju.L. and Krein M.G., *Stability of Solutions of Differential Equations in Banach Spaces*, Amm. Math. Soc. Providence , RI, 1974
- [K] M. G. Krein, *On some questions related to the ideas of Lyapunov in the theory of stability*, Uspekhi Mat. Nauk **3**, 1948, 166-169, [in Russian].
- [Na] Nagel R. (ed.), *One-parameter semigroups of positive operators*, Lect. Notes in Math. 1184, Springer- Verlag, Berlin-New York, 1986.
- [Ne] J.M.A.M. van Neerven, *The Asymptotic Behaviour of Semigroups of Linear Operators*, Birkhäuser-Verlag, 1996
- [Pa] A. Pazy, *Semigroups of Linear Operators and Applications to Partial Differential Equations*, Springer- Verlag, 1983.

- [VuS] Vũ Quốc Phóng and E. Schüler, *The operator equation $AX - XB = C$, admissibility and asymptotic behaviour of differential equations*, J. Diff. Equations, **145** (1998), 394-419.
- [RB] Rehiş M., and Buşe C., *On the Perron-Bellman theorem for C_0 -semigroups and periodic evolutionary processes in Banach spaces*, Italian Journal of Pure and Applied Mathematics, **4**(1998), 155-166.