



VICTORIA UNIVERSITY
MELBOURNE AUSTRALIA

Exponential Stability and Bounded Convolutions

This is the Published version of the following publication

Barbu, Dorel, Buse, Constantin and Dragomir, Sever S (2000) Exponential Stability and Bounded Convolutions. RGMIA research report collection, 3 (3).

The publisher's official version can be found at

Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/17321/>

EXPONENTIAL STABILITY AND BOUNDED CONVOLUTIONS

Dorel Barbu, Constantin Buse, Sever S. Dragomir

Abstract

We consider a mild solution u_f of a well-posed inhomogeneous, Cauchy problem

$$\dot{u}(t) = A(t)u(t) + f(t), \quad u(0) = 0$$

on a Banach space X , where $A(\cdot)$ is periodic. We prove that if for every almost periodic X -valued functions f , with $f(0) = 0$, the solution u_f is almost periodic, then the solution of the well-posed Cauchy problem

$$\dot{u}(t) = A(t)u(t), \quad u(0) = x \in X,$$

is uniformly exponentially stable.

1991 *Mathematics Subject Classification*: 47D06, 34G10

Keywords and phrases: evolution semigroup, exponential stability, periodic evolution families, almost periodic function.

1. Introduction

Let X be a complex Banach space and $\mathcal{L}(X)$ the space of all bounded and linear operators on X . We denote by $\|\cdot\|$ the norms of vectors and operators on X . Let $BUC(\mathbf{R}_+, X)$ the Banach space of all X -valued bounded and uniformly continuous functions on \mathbf{R}_+ endowed with sup-norm and $AP(\mathbf{R}_+, X)$ the space of almost periodic function in the sense of Bohr, i.e. the linear closed hull in $BUC(\mathbf{R}_+, X)$ of the set of all functions

$$\{e^{i\mu \cdot} x : \mu \in \mathbf{R}, x \in X\}.$$

Let $AP_0(\mathbf{R}_+, X)$ the set of all functions $f \in AP(\mathbf{R}_+, X)$ such that $f(0) = 0$. It is clear that $AP_0(\mathbf{R}_+, X)$ is a closed subspace of $AP(\mathbf{R}_+, X)$ or of $BUC(\mathbf{R}_+, X)$. We recall that a strongly continuous semigroup on X is a family $\mathbf{T} = \{T(t)\}_{t \geq 0}$ of bounded linear operators acting on the Banach space X which satisfies the following conditions:

- **(i)** $T(t + s) = T(t)T(s)$ for all $t, s \in \mathbf{R}_+ := [0, \infty)$;
- **(ii)** $T(0) = Id$, Id is the identity operator on $\mathcal{L}(X)$;
- **(iii)** the function $t \mapsto T(t)x : \mathbf{R}_+ \rightarrow X$ is continuous on \mathbf{R}_+ for all $x \in X$ (or, equivalently this function is continuous in $t = 0$).

Let \mathbf{T} be a strongly continuous semigroup on X and A its infinitesimal generator. It is well known that in this case the Cauchy problem

$$\dot{x}(t) = Ax(t), \quad x(0) = x \in X \quad (1.1)$$

is well-posed and the mild solution of (1.1) is defined by

$$x(t) = T(t)x \quad (t \geq 0).$$

For a locally integrable function $f : \mathbf{R}_+ \rightarrow X$, a mild solution of the inhomogeneous Cauchy problem

$$\dot{u}(t) = Au(t) + f(t) \quad (t \geq 0), \quad u(0) = 0$$

is defined by

$$u_f(t) = \int_0^t T(t - \xi)f(\xi)d\xi, \quad t \geq 0.$$

For a well-posed Cauchy problem

$$\dot{x}(t) = A(t)x(t) \quad (t \geq 0), \quad x(0) = x \in X \quad (1.2)$$

with (unbounded) linear operators $A(t)$ the solution lead to an evolution family $\mathcal{U} = \{U(t, s) : t \geq s \geq 0\}$ in the space $\mathcal{L}(X)$, that is

- **(e₁)** $U(t, t) = Id, U(t, \tau)U(\tau, s) = U(t, s)$ for $t \geq \tau \geq s \geq 0$;
- **(e₂)** the map $(t, s) \mapsto U(t, s)x$ is continuous for every $x \in X$.

When the Cauchy problem (1.2) is periodic, i.e. there exists $q > 0$ such that $A(t + q) = A(t)$ for all $t \in \mathbf{R}_+$, the corresponding evolution family \mathcal{U} on X has exponential growth, i.e. there exist $\omega \in \mathbf{R}$ and $M > 0$ such that

$$\|U(t, s)\| \leq Me^{\omega(t-s)} \quad \forall t \geq s \geq 0, \quad (1.3)$$

see [BP, Lemma 4.1] or [DK, Theorem 6.6]. We recall that an evolution family \mathcal{U} , as above, is called uniformly exponentially stable if there are $\omega < 0$ and $M > 0$ such that (1.3) holds. For a locally integrable function $f : \mathbf{R}_+ \rightarrow X$ a mild solution of the well-posed inhomogeneous Cauchy problem

$$\dot{u}(t) = A(t)u(t) + f(t), \quad (t \geq 0), \quad u(0) = 0$$

is defined by

$$u_f(t) = \int_0^t U(t, \tau) f(\tau) d\tau \quad (t \geq 0).$$

We shall prove the following two theorems.

THEOREM 1. Let $\mathbf{T} = \{T(t)\}_{t \geq 0}$ be a strongly continuous semigroup on X . The following statements are equivalent:

- (1) \mathbf{T} is uniformly exponentially stable, i.e. its growth bound

$$\omega_0(\mathbf{T}) := \lim_{t \rightarrow \infty} \frac{\ln \|T(t)\|}{t}$$

is negative;

- (2) the function $t \mapsto \int_0^t T(\xi) f(t - \xi) d\xi : \mathbf{R}_+ \rightarrow X$ belongs to $AP_0(\mathbf{R}_+, X)$ for all $f \in AP_0(\mathbf{R}_+, X)$;
- (3) $\sup_{t \geq 0} \|\int_0^t T(\xi) f(t - \xi) d\xi\| = M_f < \infty, \forall f \in AP_0(\mathbf{R}_+, X)$.

THEOREM 2. Let $\mathcal{U} = \{U(t, s) : t \geq s \geq 0\}$ be a q -periodic evolution family on X . The following statements are equivalent:

- (i) \mathcal{U} is uniformly exponentially stable;
- (ii) the function $t \mapsto u_f(t) = \int_0^t U(t, \xi) f(\xi) d\xi : \mathbf{R}_+ \rightarrow X$ belongs to $AP_0(\mathbf{R}_+, X)$ for all $f \in AP_0(\mathbf{R}_+, X)$;
- (iii) $\sup_{t \geq 0} \|\int_0^t U(t, \xi) f(\xi) d\xi\| = K_f < \infty, \text{ for all } f \in AP_0(\mathbf{R}_+, X)$.

2. Proofs of the theorems

Proof of Theorem 1. The proof of implications **(1)** \Rightarrow **(3)** and **(2)** \Rightarrow **(3)** are obvious and we omit the details. The proof of **(3)** \Rightarrow **(1)** is based on the following result which has been proved in [BDL, Proposition 4], see also [VS, Corollary 4.5 and its Reformulation] for a related result:

If $\sup_{t>0} \|\int_0^t e^{-i\mu\xi} T(t-\xi)g(\xi)d\xi\| < \infty$ for every $g \in P_q^0(\mathbf{R}_+, X)$ and some $\mu \in \mathbf{R}$ then $T(q)$ is power bounded and $e^{i\mu} \in \rho(T(q))$. Here $P_q^0(\mathbf{R}_+, X)$ is the set of all X -valued continuous functions such that $f(t+q) = f(t)$ for any $t \geq 0$ and $f(0) = 0$.

Now we prove that **(1)** implies **(2)**. Let $\mathcal{T} = \{\mathcal{T}^t\}_{t \geq 0}$ the evolution semigroup associated of \mathbf{T} on the space $AP_0(\mathbf{R}_+, X)$, i.e.,

$$(\mathcal{T}^t f)(s) = \begin{cases} T(t)f(s-t), & s \geq t \\ 0, & 0 \leq s \leq t \end{cases}$$

for every $f \in AP_0(\mathbf{R}_+, X)$. It is easy to see that \mathcal{T}^t acts on $AP_0(\mathbf{R}_+, X)$ for all $t \geq 0$ and, in addition, \mathcal{T} is strongly continuous, see [NM, Lemma 2]. Let $(G, D(G))$ the infinitesimal generator of \mathcal{T} and $u, f \in AP_0(\mathbf{R}_+, X)$. As in [MRS, Lemma 1.1] it is can be proves that $u \in D(G)$ and $Gu = -f$ if and only if $u = u_f$. Moreover, if \mathbf{T} is uniformly exponentially stable then the growth bound of \mathcal{T} is negative, hence G is invertible. It follows that $u_f \in D(G) \subset AP_0(\mathbf{R}_+, X)$ and the proof of Theorem 1 is finished.

Proof of Theorem 2. The proof of **(iii)** \Rightarrow **(i)** follows from the fact that if $u_{e^{-i\mu \cdot} g(\cdot)}$ is bounded for every $g \in P_q^0(\mathbf{R}_+, X)$ and some $\mu \in \mathbf{R}$ then the monodromy operator $V := U(q, 0)$ is power bounded and $e^{i\mu} \in \rho(V)$, see [B, Proof of Theorem 4]. Here $\rho(V)$ is the resolvent set of V . The proofs of **(i)** \Rightarrow **(iii)** and **(ii)** \Rightarrow **(iii)** are obvious and the proof of **(i)** \Rightarrow **(ii)** follows along the lines of **(1)** \Rightarrow **(2)** from Theorem 1. Another proof for the implication **(ii)** \Rightarrow **(i)** we will give here. This proof is based on a method indicated in [CLMR, Theorem 2.5]. Let $h : \mathbf{R} \rightarrow AP_0(\mathbf{R}_+, X)$, $(G, D(G))$ the infinitesimal generator of the evolutionary semigroup E associated to \mathcal{U} on $AP_0(\mathbf{R}_+, X)$, and

$$[(\tilde{G}h)(\theta)](t) := [Gh(\theta)](t) = \int_0^t U(t, t-\xi)(h(\theta))(t-\xi)d\xi, \quad \theta \in \mathbf{R}, t \geq 0.$$

It is easy to see that the function

$$\theta \mapsto \int_0^\cdot U(\cdot, \cdot - \xi)h(\theta)(\cdot - \xi)d\xi \text{ belongs to } AP(\mathbf{R}, AP_0(\mathbf{R}_+, X))$$

for all $h \in AP(\mathbf{R}, AP_0(\mathbf{R}_+, X))$, i.e. \tilde{G} is a linear operator on $AP(\mathbf{R}, AP_0(\mathbf{R}_+, X))$. Moreover \tilde{G} is bounded on $AP(\mathbf{R}, AP_0(\mathbf{R}_+, X))$, because

$$\begin{aligned} \|\tilde{G}h\|_{AP(\mathbf{R}, AP_0(\mathbf{R}_+, X))} &= \sup_{\theta \in \mathbf{R}} \|Gh(\theta)\|_{AP_0(\mathbf{R}_+, X)} \\ &\leq \|G\|_{\mathcal{L}(AP_0(\mathbf{R}_+, X))} \|h\|_{AP(\mathbf{R}, AP_0(\mathbf{R}_+, X))}. \end{aligned}$$

For the isometry J defined on the space $AP(\mathbf{R}, AP_0(\mathbf{R}_+, X))$ by

$$[(Jh)(\theta)](t) = (h(\theta + t))(t),$$

we have

$$[(J^{-1}\tilde{G}Jh)(\theta)](t) = \int_0^t U(t, t - \xi)(h(\theta - \xi))(t - \xi)d\xi$$

Let $E = \{E^t\}_{t \geq 0}$ be the evolution semigroup on $AP_0(\mathbf{R}_+, X)$, defined by

$$(E^t f)(\xi) = \begin{cases} U(t, t - \xi)f(t - \xi), & t \geq \xi \\ 0, & 0 \leq t \leq \xi \end{cases}$$

and

$$(G_*h)(\theta) := \int_0^\infty E^\tau h(\theta - \tau)d\tau \quad \theta \in \mathbf{R}, \quad h \in AP(\mathbf{R}, AP_0(\mathbf{R}_+, X)).$$

A simple calculus show that $G_* = J^{-1}\tilde{G}J$, therefore G_* is a bounded operator on $AP(\mathbf{R}, AP_0(\mathbf{R}_+, X))$. Each function $g_+ \in AP_0(\mathbf{R}_+, AP_0(\mathbf{R}_+, X))$ can be extended to a function $g \in AP(\mathbf{R}, AP_0(\mathbf{R}_+, X))$ by setting

$$g(\theta) = \begin{cases} g_+(\theta), & \text{if } \theta \geq 0 \\ 0, & \text{if } \theta < 0 \end{cases}$$

It is clear that $G_*g \in AP(\mathbf{R}, AP_0(\mathbf{R}_+, X))$. Consider the function $f_+ : \mathbf{R}_+ \rightarrow AP_0(\mathbf{R}_+, X)$, defined by

$$f_+(r) = \int_0^r E^\tau g_+(r - \tau) d\tau \quad (r \geq 0).$$

It is easy to see that for all $t \geq 0$, we have

$$[f_+(\theta)](t) = \int_0^{\min(\theta, t)} U(t, t - \tau)(g_+(\theta - \tau))(t - \tau) d\tau, \quad \theta \geq 0,$$

and

$$[(G_*g)(\theta)](t) = \begin{cases} (f_+(\theta))(t), & \text{if } \theta \geq 0 \\ 0, & \text{if } \theta < 0 \end{cases}$$

Then $G_*g|_{\mathbf{R}_+} = f_+$ belongs to $AP_0(\mathbf{R}_+, AP_0(\mathbf{R}_+, X))$. From Theorem 1 ((3) \Rightarrow (1)) with \mathbf{T} replaced by E and X replaced by $AP_0(\mathbf{R}_+, X)$ it results that E is uniformly exponentially stable. Now is easy to see that \mathcal{U} is uniformly exponentially stable, cf. [CLMR, Theorem 2.2].

References

- [B] C. Buşe, *Exponential Stability for Periodic Evolution Families of Bounded Linear Operators*, submitted.
- [BP] C. Buşe and A. Pogan, *Individual Exponential Stability for Evolution Families of Linear and Bounded Operators*, submitted.
- [BDL] C. Buşe, S. S. Dragomir and V. Lupulescu, *Characterizations of stability for strongly continuous semigroups by boundedness of its convolution with almost periodic function*, submitted.
- [CLMR] S. Clark, Y. Latushkin, S. Montgomery-Smith and T. Randolph, *Stability radius and internal versus external stability in Banach spaces: an evolution semigroup approach*, *Tübinger Berichte zur Funktionalanalysis*, **7**(1998), 72-102.
- [DK] D. Daners, P. Koch Medina, *Abstract Evolution Equations, Periodic Problems and Applications*, Pitman Research Notes, 1992.

- [MRS] N. V. Minh, F. Rábiger and R. Schnaubelt, *Exponential stability, exponential expansiveness and exponential dichotomy of evolution equations on the half line*, Integral Equations Operator Theory, **32**(1998), 332-353.
- [NM] T. Naito and Nguyen Van Minh, *Evolution semigroups and spectral criteria for almost periodic solutions of periodic evolution equations*, J. Differential Equations **152**(1999), 358-376.
- [VS] Vũ Quốc Phóng and E. Schüler, *The operator equation $AX - XB = C$, admisibility and asymptotic behaviour of differential equations*, J. Diff. Equations, **145** (1998), 394-419.

D. Barbu and C. Buşe
West University of Timișoara,
Bd. V. Parvan 4,
1900 Timișoara, România
E-mail buse@hilbert.math.uvt.ro

S. S. Dragomir
Victoria University of Technology
P.O. Box 14428, MCMC
Melbourne, Victoria 8001, Australia
E-mail sever@matilda.vu.edu.au