



**VICTORIA UNIVERSITY**  
MELBOURNE AUSTRALIA

*Some Further Inequalities for Univariate Moments  
and Some New Ones for the Covariance*

This is the Published version of the following publication

Barnett, Neil S and Dragomir, Sever S (2000) Some Further Inequalities for Univariate Moments and Some New Ones for the Covariance. RGMIA research report collection, 3 (4).

The publisher's official version can be found at

Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/17342/>

# SOME FURTHER INEQUALITIES FOR UNIVARIATE MOMENTS AND SOME NEW ONES FOR THE COVARIANCE

N.S. BARNETT AND S.S. DRAGOMIR

ABSTRACT. In this paper some further inequalities for univariate moments are given with particular reference to the expectations of the extreme order statistics. In addition, some inequalities are obtained for the covariance of two continuous random variables.

## 1. INTRODUCTION

The expectation of a continuous random variable is given by

$$E(X) = \int_{-\infty}^{\infty} x f(x) dx,$$

where  $f(x)$  is the probability density function of the random variable,  $X$ . If the interval of definition is finite,  $(a, b)$ , then this can be further expressed, using integration by parts, as

$$(1.1) \quad E(X) = b - \int_a^b F(x) dx,$$

where  $F(x)$  is the associated cumulative distribution function. This result has been exploited variously to obtain inequalities involving the expectation and variance, see for example [1], [2], [3].

The aim of this paper is to provide some additional inequalities utilising a generalisation of (1.1) to higher moments, as well as providing some specific results for the extreme order statistics.

In addition, some results are obtained involving the covariance of two random variables using a bivariate generalisation of (1.1) and generalisations of the inequalities of Grüss and Ostrowski.

## 2. UNIVARIATE RESULTS

Denote by  $M_n$  the  $n^{\text{th}}$  moment  $\int_a^b x^n f(x) dx$  which, using integration by parts, can be expressed as

$$(2.1) \quad b^n - n \int_a^b x^{n-1} F(x) dx.$$

---

*Date:* August 25, 2000.

*1991 Mathematics Subject Classification.* Primary 26D15, 26D99; Secondary 65A99.

*Key words and phrases.* Grüss Inequality, Chebychev's Inequality, Cumulative Distribution Functions, Random Variables, Covariance.

Consider now

$$\left| \frac{1}{b-a} \int_a^b x^{n-1} F(x) dx - \frac{1}{(b-a)^2} \int_a^b x^{n-1} dx \int_a^b F(x) dx \right|$$

for which various inequalities can be found. Before exploiting some of these, we express the difference in terms of the distributional moments.

$$\frac{1}{(b-a)^2} \int_a^b x^{n-1} dx \int_a^b F(x) dx = \frac{(b^n - a^n)}{n(b-a)^2} \{b - E(X)\}.$$

We therefore have:-

$$\begin{aligned} (2.2) \quad & \left| \frac{1}{b-a} \int_a^b x^{n-1} F(x) dx - \frac{(b^n - a^n)}{n(b-a)^2} (b - E(X)) \right| \\ &= \left| \frac{b_n}{n(b-a)} - \frac{M_n}{n(b-a)} - \frac{b(b^n - a^n)}{n(b-a)^2} + \frac{M_1(b^n - a^n)}{n(b-a)^2} \right| \\ &= \left| \frac{ab(a^{n-1} - b^{n-1})}{n(b-a)^2} - \frac{M_n}{n(b-a)} + \frac{M_1(b^n - a^n)}{n(b-a)^2} \right| \\ &= \frac{1}{n(b-a)^2} |a^n(b - M_1) - b^n(a - M_1) - M_n(b-a)|. \end{aligned}$$

Now utilising various inequalities, we can obtain a number of results.

**Pre-Grüss.**

Using an inequality of [4] applied to (2.2), we have

$$\begin{aligned} (2.3) \quad & \frac{1}{n(b-a)^2} |a^n(b - M_1) - b^n(a - M_1) - M_n(b-a)| \\ &\leq \frac{1}{2} \left[ \frac{1}{(b-a)} \int_a^b x^{2(n-1)} dx - \left( \frac{1}{(b-a)} \int_a^b x^{n-1} dx \right)^2 \right]^{\frac{1}{2}} \\ &= \frac{1}{2} \left[ \frac{b^{2n-1} - a^{2n-1}}{(b-a)(2n-1)} - \left( \frac{b^n - a^n}{n(b-a)} \right)^2 \right]^{\frac{1}{2}}. \end{aligned}$$

The special case when  $n = 2$  gives:-

$$\begin{aligned} & \left| a^2(b - E(X)) - b^2(a - E(X)) - (b-a)(\sigma^2 + (E(X))^2) \right| \\ &\leq (b-a)^2 \left[ \frac{b^3 - a^3}{3(b-a)} - \left( \frac{b^2 - a^2}{2(b-a)} \right)^2 \right]^{\frac{1}{2}}, \end{aligned}$$

that is,

$$\begin{aligned} & |(b - E(X))(E(X) - a) - \sigma^2| \\ &\leq (b-a)^2 \left[ \frac{b^2 + ab + a^2}{3} - \frac{(b^2 + 2ab + a^2)}{4} \right]^{\frac{1}{2}} \\ &= \frac{(b-a)^3}{2\sqrt{3}}, \end{aligned}$$

which is Theorem 3 of [3].

**Pre-Chebyshev.**

Using a further result of [4] and (2.2) we can obtain

$$\begin{aligned} & \frac{1}{n(b-a)^2} |a^n(b-M_1) - b^n(a-M_1) - M_n(b-a)| \\ & \leq \frac{(b-a)}{2\sqrt{3}} \|f\|_\infty \left[ \frac{1}{(b-a)} \int_a^b x^{2(n-1)} dx - \left( \frac{1}{(b-a)} \int_a^b x^{n-1} dx \right)^2 \right]^{\frac{1}{2}}, \end{aligned}$$

where  $\|f\|_\infty$  is the supremum of the probability density function over  $(a, b)$ , giving

$$(2.4) \quad \begin{aligned} & |a^n(b-M_1) - b^n(a-M_1) - M_n(b-a)| \\ & \leq \frac{n(b-a)^3}{2\sqrt{3}} \|f\|_\infty \left[ \frac{b^{2n-1} - a^{2n-1}}{(b-a)(2n-1)} - \left( \frac{b^n - a^n}{n(b-a)} \right)^2 \right]^{\frac{1}{2}}. \end{aligned}$$

The special case where  $n = 2$  gives:-

$$|(b - E(X))(E(X) - a) - \sigma^2| \leq \frac{(b-a)^3}{6} \|f\|_\infty$$

which was obtained by Barnett and Dragomir in [5].

### 3. LIPSCHITZIAN MAPPINGS

If  $x^{n-1}F(x)$  is of the Lipschitzian type, then

$$|x^{n-1}F(x) - y^{n-1}F(y)| \leq L|x - y|,$$

where  $L \geq 0$  in which case

$$\left| x^{n-1}F(x) - \frac{1}{b-a} \int_a^b x^{n-1}F(x) dx \right| \leq \frac{L}{2} \left[ \left( \frac{x-a}{b-a} \right)^2 + \left( \frac{b-x}{b-a} \right)^2 \right] (b-a).$$

(Ostrowski's inequality, [7].)

Now,

$$\left| x^{n-1}F(x) - \frac{1}{b-a} \int_a^b x^{n-1}F(x) dx \right| = \left| x^{n-1}F(x) - \left\{ \frac{b^n - M_n}{n(b-a)} \right\} \right|$$

and thus we have;-

$$\left| x^{n-1}F(x) - \left\{ \frac{b^n - M_n}{n(b-a)} \right\} \right| \leq \frac{L}{2} \left[ \left( \frac{x-a}{b-a} \right)^2 + \left( \frac{b-x}{b-a} \right)^2 \right] (b-a).$$

If  $n = 2$ , then

$$\left| xF(x) - \left\{ \frac{b^2 - M_2}{2(b-a)} \right\} \right| \leq \frac{L}{2} \left[ \left( \frac{x-a}{b-a} \right)^2 + \left( \frac{b-x}{b-a} \right)^2 \right] (b-a).$$

Consider now the mapping  $F(x)$ ,  $x \in [a, b]$ , then the mapping is Lipschitzian if there exists  $L > 0$  such that

$$|F(x) - F(y)| \leq L|x - y|.$$

Now, if  $F(\cdot)$  is a cumulative distribution function, it is monotonic increasing between 0 and 1 over  $[a, b]$ . It is apparent that there exists  $z \in [x, y]$  such that

$$\frac{F(x) - F(y)}{x - y} \leq \left[ \frac{dF(x)}{dx} \right]_{x=z}.$$

Thus, if we choose  $L = \max \left[ \frac{dF(x)}{dx} \right]_{x=z}$  for  $z \in [a, b]$ , then this implies that  $F(x)$  is Lipschitzian, since

$$|F(x) - F(y)| \leq \|f\|_{\infty} |x - y|.$$

Consider similarly the mapping  $xF(x)$ , it is also monotonic increasing and by the same token, there exists  $z \in [x, y]$  such that

$$\left| \frac{xF(x) - yF(y)}{x - y} \right| \leq \left| \left[ \frac{d(xF(x))}{dx} \right]_{x=z} \right|.$$

In addition, we have that

$$\left| \left[ \frac{d(xF(x))}{dx} \right]_{x=z} \right| = \left| \left[ F(x) + x \frac{dF(x)}{dx} \right]_{x=z} \right|$$

and hence

$$\begin{aligned} \left| F(z) + z \left[ \frac{dF(x)}{dx} \right]_{x=z} \right| &\leq |F(z)| + \left| z \left[ \frac{dF(x)}{dx} \right]_{x=z} \right| \\ &\leq 1 + \|f\|_{\infty} \max\{|a|, |b|\}. \end{aligned}$$

Thus,  $L$  can be taken to be

$$1 + \|f\|_{\infty} \max\{|a|, |b|\}$$

and then

$$|xF(x) - yF(y)| < [1 + \|f\|_{\infty} \max\{|a|, |b|\}] |x - y|,$$

and so  $xF(x)$  is Lipschitzian.

Similarly it can be shown that  $x^{n-1}F(x)$  is Lipschitzian for  $n = 3, 4, \dots$

Thus,

$$\begin{aligned} &\left| xF(x) - \left\{ \frac{b^2 - M_2}{2(b-a)} \right\} \right| \\ &\leq \frac{1}{2} [1 + \|f\|_{\infty} \max\{|a|, |b|\}] \left[ \left( \frac{x-a}{b-a} \right)^2 + \left( \frac{b-x}{b-a} \right)^2 \right] (b-a). \end{aligned}$$

For  $x = a$  we get

$$\begin{aligned} |M_2 - b^2| &= \left| \sigma^2 + ((E(X))^2) - b^2 \right| \\ &\leq (b-a)^2 [1 + \|f\|_{\infty} \max\{|a|, |b|\}] \end{aligned}$$

and for  $x = b$  we have

$$\begin{aligned} |2b(b-a) - b^2 + M_2| &= |b^2 - 2ab + M_2| \\ &= \left| b(b-2a) + \sigma^2 + ((E(X))^2) \right| \\ &\leq (b-a)^2 [1 + \|f\|_{\infty} \max\{|a|, |b|\}]. \end{aligned}$$

## 4. DISTRIBUTIONS OF THE MAXIMUM, MINIMUM AND RANGE OF A SAMPLE

Consider a continuous random variable  $X$  with a non-zero probability density function over a finite interval  $[a, b]$ . Consider a random sample  $X_1, X_2, \dots, X_n$ . We consider the distribution function of the maximum, minimum and the range of the random sample.

**Maximum**

Let the cumulative distribution function of the maximum be  $G(x)$ , the probability density function be  $g(x)$ , and the corresponding functions for  $X$  be  $F(x)$  and  $f(x)$ . Then

$$G(x) = \Pr[\max \leq x] = \Pr[\text{all } X_1, \dots, X_n \leq x].$$

Therefore  $G(x) = [F(x)]^n$  and  $g(x) = n[F(x)]^{n-1} f(x)$ .

**Minimum**

Let the cumulative distribution function of the minimum and the probability density function be  $H(x)$  and  $h(x)$  respectively. Therefore

$$\begin{aligned} H(x) &= \Pr[\min \leq x] = 1 - \Pr[\min \geq x] \\ &= 1 - \Pr[\text{all } X_1, \dots, X_n \geq x] = 1 - [1 - F(x)]^n \end{aligned}$$

and

$$h(x) = n[1 - F(x)]^{n-1} f(x).$$

**Range**

Let the distribution function of the range be  $R(x)$  and the probability density function be  $r(x)$ . Consider

$$\begin{aligned} K(s, t) &= \Pr[\max \leq s, \min \leq t] \\ &= \Pr[\max \leq s] - \Pr[\max \leq s, \min \geq t] \\ &= [F(s)]^n - \Pr\{t \leq \text{all } X_1, \dots, X_n \leq s\} \\ &= [F(s)]^n - [F(s) - F(t)]^n. \end{aligned}$$

Therefore, the joint probability density function of the extreme order statistics can be found by differentiating this with respect to  $s$  and  $t$ , giving

$$k(s, t) = n(n-1)f(s)f(t)[F(s) - F(t)]^{n-2}, \quad s > t$$

which for a random variable defined on  $[a, b]$  gives

$$r(x) = n(n-1) \int_{s=a}^{b-x} f(s+x)f(s)[F(s+x) - F(s)]^{n-2} ds,$$

where  $0 < x < b - a$ .

## 5. APPLICATION OF GRÜSS' INEQUALITY TO POSITIVE INTEGER POWERS OF A FUNCTION

As a preliminary to the proof of Grüss' inequality we can establish the identity:

$$(5.1) \quad \frac{1}{b-a} \int_a^b g(x)f(x) dx = p + \left(\frac{1}{b-a}\right)^2 \int_a^b f(x) dx \cdot \int_a^b g(x) dx,$$

where it can be subsequently shown that

$$|p| \leq \frac{1}{4} [\Gamma - \gamma] [\Phi - \phi],$$

and  $\gamma, \Gamma, \phi, \Phi$  are respectively lower and upper bounds of  $f(x)$  and  $g(x)$ .

Applying this same identity to the square of a function, we have

$$(5.2) \quad \frac{1}{b-a} \int_a^b f^2(x) dx = p_1 + \left(\frac{1}{b-a}\right)^2 \left(\int_a^b f(x) dx\right)^2$$

Similarly,

$$\frac{1}{b-a} \int_a^b f^3(x) dx = p_2 + \frac{1}{b-a} \int_a^b f(x) dx \cdot \frac{1}{b-a} \int_a^b f^2(x) dx$$

and using (5.1), we have

$$(5.3) \quad \frac{1}{b-a} \int_a^b f^3(x) dx = p_2 + \frac{p_1}{b-a} \int_a^b f(x) dx + \left(\frac{1}{b-a}\right)^3 \left(\int_a^b f(x) dx\right)^3.$$

Continuing, we can show that for positive integers  $n$ ,  $n \geq 2$

$$(5.4) \quad \begin{aligned} & \frac{1}{b-a} \int_a^b f^n(x) dx - \left(\frac{1}{b-a}\right)^n \left(\int_a^b f(x) dx\right)^n \\ &= p_{n-1} + p_{n-2} \left(\frac{1}{b-a} \int_a^b f(x) dx\right) + p_{n-3} \left(\frac{1}{b-a} \int_a^b f(x) dx\right)^2 \\ & \quad + \cdots + p_1 \left(\frac{1}{b-a} \int_a^b f(x) dx\right)^{n-2}, \end{aligned}$$

giving:-

$$\begin{aligned} & \left| \frac{1}{b-a} \int_a^b f^n(x) dx - \left(\frac{1}{b-a}\right)^n \left(\int_a^b f(x) dx\right)^n \right| \\ & \leq |p_{n-1}| + |p_{n-2}| \left| \frac{1}{b-a} \int_a^b f(x) dx \right| + \cdots + |p_1| \left| \left(\frac{1}{b-a} \int_a^b f(x) dx\right)^{n-2} \right|, \end{aligned}$$

where

$$|p_1| \leq \frac{1}{4} (\Gamma - \gamma)^2, \quad \gamma < f(x) < \Gamma,$$

$$|p_2| \leq \frac{1}{4} (\Gamma - \gamma) (\Phi_1 - \phi_1), \quad \phi_1 < f^2(x) < \Phi_1,$$

$$|p_3| \leq \frac{1}{4} (\Gamma - \gamma) (\Phi_2 - \phi_2), \quad \phi_2 < f^3(x) < \Phi_2,$$

.....

Assuming that  $f(x) \geq 0$  and denoting  $\left(\frac{1}{b-a} \int_a^b f(x) dx\right)^n$  by  $\lambda^n$ , we have

$$\begin{aligned}
(5.5) \quad & \left| \frac{1}{b-a} \int_a^b f^n(x) dx - \left(\frac{1}{b-a}\right)^n \left(\int_a^b f(x) dx\right)^n \right| \\
& \leq \frac{1}{4} (\Gamma - \gamma) [\Gamma^{n-1} - \gamma^{n-1}] + \frac{1}{4} (\Gamma - \gamma) [\Gamma^{n-2} - \gamma^{n-2}] \lambda \\
& \quad + \frac{1}{4} (\Gamma - \gamma) [\Gamma^{n-3} - \gamma^{n-3}] \lambda^2 + \cdots + \frac{1}{4} (\Gamma - \gamma) (\Gamma - \gamma) \lambda^{n-2} \\
& = \frac{1}{4} (\Gamma - \gamma) \sum_{i=1}^{n-1} (\Gamma^{n-i} - \gamma^{n-i}) \lambda^{i-1}.
\end{aligned}$$

Now, if  $f$  is a p.d.f., the right hand side reduces to

$$\begin{aligned}
& \frac{1}{4} \Gamma \sum_{i=1}^{n-1} \Gamma^{n-i} \left(\frac{1}{b-a}\right)^{i-1} \\
& = \frac{1}{4} \Gamma^n \sum_{i=1}^{n-1} \left(\frac{1}{\Gamma(b-a)}\right)^{i-1} = \frac{1}{4} \Gamma^n \left(\frac{1 - \left(\frac{1}{\Gamma(b-a)}\right)^{n-1}}{1 - \frac{1}{\Gamma(b-a)}}\right) \\
& = \frac{\Gamma^n}{4\Gamma^{n-2}(b-a)^{n-2}} \left(\frac{\Gamma^{n-1}(b-a)^{n-1} - 1}{\Gamma(b-a) - 1}\right) \\
& = \frac{\Gamma^2}{4(b-a)^{n-2}} \left(\frac{\Gamma^{n-1}(b-a)^{n-1} - 1}{\Gamma(b-a) - 1}\right).
\end{aligned}$$

If we now consider this inequality for an associated cumulative distribution function  $F(\cdot)$ , we have that

$$\lambda = \frac{b - E(X)}{b - a}$$

and the right hand side of (5.5) becomes

$$\frac{1}{4} \sum_{i=1}^{n-1} \left(\frac{b - E(X)}{b - a}\right)^{i-1} = \frac{[(b-a)^{n-1} - (b - E(X))^{n-1}]}{4(E(X) - a)(b-a)^{n-2}}.$$

Thus, we have the two inequalities:-

$$(5.6) \quad \left| \frac{1}{b-a} \int_a^b f^n(x) dx - \left(\frac{1}{b-a}\right)^n \right| \leq \frac{\Gamma^2}{4(b-a)^{n-2}} \left(\frac{\Gamma^{n-1}(b-a)^{n-1} - 1}{\Gamma(b-a) - 1}\right)$$

and

$$(5.7) \quad \left| \frac{1}{b-a} \int_a^b F^n(x) dx - \left(\frac{b - E(X)}{b - a}\right)^n \right| \leq \frac{[(b-a)^{n-1} - (b - E(X))^{n-1}]}{4(E(X) - a)(b-a)^{n-2}}.$$

Similarly, we can develop an inequality for  $1 - F(x)$  by suitable substitution in (5.5), that is,

$$\lambda = \frac{E(X) - a}{b - a}$$



which gives

$$\begin{aligned}
 (5.8) \quad & \left| \frac{1}{b-a} \int_a^b (1-F(x))^n dx - \left( \frac{1}{b-a} \int_a^b (1-F(x)) dx \right)^n \right| \\
 &= \left| \frac{1}{b-a} \int_a^b (1-F(x))^n dx - \left( \frac{E(X)-a}{b-a} \right)^n \right| \\
 &\leq \frac{[(b-a)^{n-1} - (E(X)-a)^{n-1}]}{4(b-a)^{n-2}(b-E(X))}.
 \end{aligned}$$

## 6. INEQUALITIES FOR THE EXPECTATION OF THE EXTREME ORDER STATISTICS

As the p.d.f. of the maximum is

$$g(x) = n [F(x)]^{n-1} f(x),$$

then

$$E[X_{\max}] = n \int_a^b x [F(x)]^{n-1} f(x) dx.$$

Integrating by parts gives

$$\begin{aligned}
 E[X_{\max}] &= n \left[ \left[ \frac{x}{n} [F(x)]^n \right]_a^b - \frac{1}{n} \int_a^b (F(x))^n dx \right] \\
 &= b - \int_a^b F^n(x) dx
 \end{aligned}$$

giving, from (5.7)

$$(6.1) \quad \left| \frac{b - E[X_{\max}]}{b-a} - \left( \frac{b - E(X)}{b-a} \right)^n \right| \leq \frac{[(b-a)^{n-1} - (b - E(X))^{n-1}]}{4(E(X) - a)(b-a)^{n-2}}$$

and when  $E(X) = \frac{a+b}{2}$ , we have:-

$$(6.2) \quad \left| \frac{b - E[X_{\max}]}{b-a} - \frac{1}{2^n} \right| \leq \left( \frac{2^{n-1} - 1}{2^n} \right).$$

Consider now  $E(X_{\min}) = n \int_{x=a}^b x f(x) [1-F(x)]^{n-1} dx$ . Integration by parts gives:-

$$E[X_{\min}] = n \left[ \left[ -\frac{x}{n} [1-F(x)]^n \right]_a^b + \frac{1}{n} \int_a^b (1-F(x))^n dx \right]$$

and so

$$E(X_{\min}) = a + \int_a^b (1-F(x))^n dx.$$

Utilising (5.8) we have

$$(6.3) \quad \left| \frac{E(X_{\min}) - a}{b-a} - \left( \frac{E(X) - a}{b-a} \right)^n \right| \leq \frac{[(b-a)^{n-1} - (E(X) - a)^{n-1}]}{4(b-a)^{n-2}(b-E(X))}$$

and when  $E(X) = \frac{a+b}{2}$ , we have:-

$$(6.4) \quad \left| \frac{E(X_{\min}) - a}{b-a} - \frac{1}{2^n} \right| \leq \left( \frac{2^{n-1} - 1}{2^n} \right).$$

## 7. APPLICATIONS TO THE BETA DISTRIBUTION

The Beta probability density function is given by

$$\frac{1}{B(\alpha, \beta)} x^{\alpha-1} (1-x)^{\beta-1}, \quad 0 < x < 1, \quad \alpha, \beta > 0,$$

where

$$B(\alpha, \beta) = \int_0^1 x^{\alpha-1} (1-x)^{\beta-1} dx.$$

Clearly,

$$\begin{aligned} E(X) &= \frac{1}{B(\alpha, \beta)} \int_0^1 x^{\alpha+1-1} (1-x)^{\beta-1} dx = \frac{1}{B(\alpha, \beta)} B(\alpha+1, \beta) \\ &= \frac{\Gamma(\alpha+1) \Gamma(\alpha+\beta)}{\Gamma(\alpha+\beta+1) \Gamma(\alpha)} = \frac{\alpha \Gamma(\alpha) \Gamma(\alpha+\beta)}{(\alpha+\beta) \Gamma(\alpha+\beta) \Gamma(\alpha)} = \frac{\alpha}{(\alpha+\beta)}. \end{aligned}$$

Substituting  $a = 0$ ,  $b = 1$  and letting ' $\Gamma$ '  $\equiv m$ , from (5.6) we obtain

$$\left| \int_0^1 f^n(x) dx - 1 \right| \leq \frac{m^2 (1 - m^{n-1})}{4(1-m)}$$

and further,

$$\begin{aligned} & \left| \frac{1}{B^n(\alpha, \beta)} \int_0^1 x^{n(\alpha-1)} (1-x)^{n(\beta-1)} dx - 1 \right| \\ &= \left| \frac{B(n(\alpha-1)+1, n(\beta-1)+1)}{B^n(\alpha, \beta)} - 1 \right| \leq \frac{m^2 (1 - m^{n-1})}{4(1-m)} \end{aligned}$$

and  $m$  is the value of  $x$  for which  $f'(x) = 0$ , that is

$$\begin{aligned} & (1-x)^{\beta-1} (\alpha-1) x^{\alpha-2} + x^{\alpha-1} (\beta-1) (1-x)^{\beta-2} (-1) = 0. \\ & x^{\alpha-2} (1-x)^{\beta-2} \{(\alpha-1)(1-x) - x(\beta-1)\}, \\ \text{i.e. } m &= \frac{\alpha-1}{\alpha+\beta-2}, \quad \alpha, \beta > 1. \end{aligned}$$

We then have the inequality:-

$$\begin{aligned} & \left| \frac{B(n(\alpha-1)+1, n(\beta-1)+1)}{B^n(\alpha, \beta)} - 1 \right| \\ & \leq \frac{(\alpha-1)^2}{4(\alpha+\beta-2)(\beta-1)} \left\{ 1 - \left( \frac{\alpha-1}{\alpha+\beta-2} \right)^{n-1} \right\}. \end{aligned}$$

When  $\alpha = \beta$ , the right hand side becomes  $\frac{1}{8} \left( 1 - \frac{1}{2^{n-1}} \right)$ .

Consider now (6.1) when  $f(x)$  is the p.d.f. of a Beta distribution. This gives:-

$$\begin{aligned} & \left| 1 - E(X_{\max}) - \left(1 - \frac{\alpha}{\alpha + \beta}\right)^n \right| \\ &= \left| 1 - \left(\frac{\beta}{\alpha + \beta}\right)^n - E(X_{\max}) \right| \leq \frac{\left[1 - \left(1 - \frac{\alpha}{\alpha + \beta}\right)^{n-1}\right]}{4 \left(\frac{\alpha}{\alpha + \beta}\right)} \\ &= \frac{(\alpha + \beta)^{n-1} - \beta^{n-1}}{4\alpha(\alpha + \beta)^{n-2}} \end{aligned}$$

and when  $\alpha = \beta$ , this becomes:-

$$\left| 1 - \left(\frac{1}{2}\right)^n - E(X_{\max}) \right| \leq \frac{1 - \left(\frac{1}{2}\right)^{n-1}}{2}$$

and from (6.3)

$$\left| E(X_{\min}) - \left(\frac{\alpha}{\alpha + \beta}\right)^n \right| \leq \frac{\left[1 - \left(\frac{\alpha}{\alpha + \beta}\right)^{n-1}\right]}{4 \left(1 - \frac{\alpha}{\alpha + \beta}\right)} = \frac{(\alpha + \beta)^{n-1} - \alpha^{n-1}}{4\beta(\alpha + \beta)^{n-2}}$$

and when  $\alpha = \beta$

$$\left| E(X_{\min}) - \frac{1}{2^n} \right| \leq \frac{1 - \left(\frac{1}{2}\right)^{n-1}}{2}.$$

From these we can obtain further  $E(X_{\min}) \leq \frac{1}{2}$  and  $\frac{1}{2} \leq E(X_{\max}) \leq \frac{3}{2} - \left(\frac{1}{2}\right)^{n-1}$ .

## 8. SOME BOUNDS FOR JOINT MOMENTS AND PROBABILITIES USING OSTROWSKI TYPE INEQUALITIES FOR DOUBLE INTEGRALS

**Theorem 1.** *Let  $X, Y$  be two continuous random variables  $x \in [a, b]$ ,  $y \in [c, d]$  with probability density functions  $f_1(\cdot)$  and  $f_2(\cdot)$  respectively and with joint probability density function  $f(\cdot, \cdot)$  with associated cumulative distribution functions  $F_1(\cdot)$ ,  $F_2(\cdot)$  and  $F(\cdot, \cdot)$ . Then*

$$(8.1) \quad E(XY) = bE(Y) + dE(X) - bd + \int_{s=a}^b \int_{t=c}^d F(s, t) dsdt.$$

This is a generalisation of the result

$$E(X) = b - \int_a^b F(s) ds$$

and is equivalent to:-

$$(8.2) \quad E(XY) = bd - d \int_a^b F_1(s) ds - b \int_c^d F_2(t) dt + \int_{s=a}^b \int_{t=c}^d F(s, t) dsdt.$$

*Proof.*

$$E(XY) = \int_{t=c}^d t \left\{ \int_a^b s f(s, t) ds \right\} dt$$

and

$$\begin{aligned} \int_a^b s f(s, t) ds &= \left[ s \int_{u=a}^s f(u, t) du \right]_{s=a}^b - \int_{s=a}^b \left( \int_{u=a}^s f(u, t) du \right) ds \\ &= b f_2(t) - \int_{s=a}^b \left( \int_{u=a}^s f(u, t) du \right) ds, \end{aligned}$$

so

$$\begin{aligned} E(XY) &= b \int_c^d t f_2(t) dt - \int_{t=c}^d t \left[ \int_{s=a}^b \left( \int_{u=a}^s f(u, t) du \right) ds \right] dt \\ &= bE(Y) - \int_{s=a}^b \left( \int_{u=a}^s \left( \int_{t=c}^d t f(u, t) dt \right) du \right) ds \end{aligned}$$

Now

$$\begin{aligned} \int_{t=c}^d t f(u, t) dt &= \left[ t \int_{v=c}^t f(u, v) dv \right]_{t=c}^d - \int_{t=c}^d \left( \int_{v=c}^t f(u, v) dv \right) dt \\ &= d f_1(u) - \int_{t=c}^d \left( \int_{v=c}^t f(u, v) dv \right) dt, \end{aligned}$$

$$\begin{aligned} E(XY) &= bE(Y) - \int_{s=a}^b \left( \int_{u=a}^s \left\{ d f_1(u) - \int_{t=c}^d \left( \int_{v=c}^t f(u, v) dv \right) dt \right\} du \right) ds, \\ &= bE(Y) - d \int_{s=a}^b F_1(s) ds \\ &\quad + \int_{t=c}^d \left( \int_{s=a}^b \left( \int_{v=c}^t \left( \int_{u=a}^s f(u, v) du \right) dv \right) ds \right) dt \\ &= bE(Y) + dE(X) - bd + \int_{s=a}^b \int_{t=c}^d F(s, t) ds dt \end{aligned}$$

and, equivalently,

$$E(XY) = bd - d \int_a^b F_1(s) ds - b \int_c^d F_2(t) dt + \int_a^b \int_c^d F(s, t) ds dt.$$

□

In [6] Barnett and Dragomir proved the following theorem.

**Theorem 2.** *Let  $f : [a, b] \times [c, d] \rightarrow \mathbb{R}$  be continuous on  $[a, b] \times [c, d]$ ,  $f''_{x,y} = \frac{\partial^2 f}{\partial x \partial y}$  exists on  $(a, b) \times (c, d)$  and is bounded, i.e.,*

$$\|f''_{s,t}\|_{\infty} := \sup_{(x,y) \in (a,b) \times (c,d)} \left| \frac{\partial^2 f(x,y)}{\partial x \partial y} \right| < \infty$$

then we have the inequality:

$$\begin{aligned}
(8.3) \quad & \left| \int_a^b \int_c^d f(s, t) ds dt \right. \\
& \left. - \left[ (b-a) \int_c^d f(x, t) dt + (d-c) \int_a^b f(s, y) ds - (d-c)(b-a)f(x, y) \right] \right| \\
& \leq \left[ \frac{1}{4}(b-a)^2 + \left(x - \frac{a+b}{2}\right)^2 \right] \\
& \quad \times \left[ \frac{1}{4}(d-c)^2 + \left(y - \frac{c+d}{2}\right)^2 \right] \|f''_{s,t}\|_\infty
\end{aligned}$$

for all  $(x, y) \in [a, b] \times [c, d]$ .

If we apply this taking  $f(\cdot, \cdot)$  to be a joint cumulative distribution function  $F(\cdot, \cdot)$  with  $x = b$ ,  $y = d$  we obtain

$$\begin{aligned}
& \left| \int_a^b \int_c^d F(s, t) ds dt \right. \\
& \left. - (b-a) \int_c^d F(b, t) dt - (d-c) \int_a^b F(s, d) ds + (d-c)(b-a) \right| \\
& \leq \frac{1}{4}(b-a)^2(d-c)^2 \|F''_{s,t}\|_\infty
\end{aligned}$$

That is

$$\begin{aligned}
& \left| \int_a^b \int_c^d F(s, t) ds dt \right. \\
& \left. - (b-a) \int_c^d F_2(t) dt - (d-c) \int_a^b F_1(s) ds + (d-c)(b-a) \right| \\
& \leq \frac{1}{4}(b-a)^2(d-c)^2 \|F''_{s,t}\|_\infty.
\end{aligned}$$

Using (8.2), this gives

$$\begin{aligned}
& \left| E(XY) + a \int_c^d F_2(t) dt + c \int_a^b F_1(s) ds - ad - bc + ac \right| \\
& = |E(XY) + aE(Y) - cE(X) + ac| \leq \frac{1}{4}(b-a)^2(d-c)^2 \|F''_{s,t}\|_\infty \\
& = \frac{1}{4}(b-a)^2(d-c)^2 \|f\|_\infty,
\end{aligned}$$

providing bounds for  $E(XY)$  in terms of  $E(X)$  and  $E(Y)$ .

Since  $Cov(X, Y) = E(XY) - E(X) \cdot E(Y)$ , we can write the left hand side alternatively as:-

$$Cov(X, Y) + [c - E(Y)][a - E(X)].$$

We can similarly extract other bounds from (8.3) in the situations where

- (i)  $x = b$ ,  $y = c$ ,
- (ii)  $x = a$ ,  $y = d$ , and

(iii)  $x = a, y = c$

giving respectively

$$\begin{aligned} |E(XY) - dE(X) - aE(Y) + ad| &\leq \frac{1}{4} (b-a)^2 (d-c)^2 \|f\|_\infty, \\ |E(XY) - cE(X) - bE(Y) + bc| &\leq \frac{1}{4} (b-a)^2 (d-c)^2 \|f\|_\infty \end{aligned}$$

and

$$|E(XY) - dE(X) - bE(Y) + bd| \leq \frac{1}{4} (b-a)^2 (d-c)^2 \|f\|_\infty.$$

We can use the results of [8] by Dragomir, Cerone, Barnett and Roumeliotis to obtain further inequalities relating the first single and joint moments as well as some involving joint probabilities.

In [8], bounds were obtained for:-

$$\left| \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(s,t) dsdt - f(x,y) \right|$$

namely  $M_1(x) + M_2(y) + M_3(x,y)$  where these are as defined in [8]. For one particular case we have

$$\begin{aligned} M_1(x) &= \frac{\left[\frac{1}{2}(b-a) + \left|x - \frac{a+b}{2}\right|\right]}{(b-a)(d-c)} \left\| \frac{\partial f(s,t)}{\partial t} \right\|_1, \\ M_2(y) &= \frac{\left[\frac{1}{2}(d-c) + \left|y - \frac{c+d}{2}\right|\right]}{(b-a)(d-c)} \left\| \frac{\partial f(s,t)}{\partial s} \right\|_1 \end{aligned}$$

and

$$M_3(x,y) = \frac{\left[\frac{1}{2}(b-a) + \left|x - \frac{a+b}{2}\right|\right] \left[\frac{1}{2}(d-c) + \left|y - \frac{c+d}{2}\right|\right]}{(b-a)(d-c)} \left\| \frac{\partial^2 f(s,t)}{\partial s \partial t} \right\|_1.$$

It follows then that if we choose  $f$  to be the joint cumulative distribution function,  $F(x,y)$ , we can generate the following inequalities

$$\left| \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d F(s,t) dsdt \right| \leq M_1(a) + M_2(c) + M_3(a,c),$$

and

$$\left| \int_a^b \int_c^d F(s,t) dsdt - \Pr\{X \leq x, Y \leq y\} \right| \leq M_1(x) + M_2(y) + M_3(x,y).$$

The first of these simplifies to give:-

$$|E(XY) - bE(Y) - dE(X) + bd| \leq (b-a) + (d-c) + (d-c)(b-a).$$

## 9. FURTHER INEQUALITIES FOR THE COVARIANCE USING GRÜSS INEQUALITY

Consider functions  $f_1(\cdot)$ ,  $f_2(\cdot)$  and  $f(\cdot, \cdot)$  where  $f_1$  is integrable over  $[a, b]$ ,  $f_2$  is integrable over  $[c, d]$  and  $f(x, y)$  is integrable for  $x \in [a, b]$  and  $y \in [c, d]$ . Consider the integral

$$\int_a^b \int_c^d f_1(s) f_2(t) f(s,t) dsdt = \int_{s=a}^b f_1(s) \left( \int_{t=c}^d f_2(t) f(s,t) dt \right) ds.$$

We have:-

$$\begin{aligned} & \frac{1}{d-c} \int_c^d f_2(t) f(s,t) dt - \left( \frac{1}{d-c} \right)^2 \int_c^d f_2(t) dt \int_{t=c}^d f(s,t) dt \\ &= p_1(s) \quad (\text{say}). \end{aligned}$$

Hence,

$$\begin{aligned} & \int_a^b \int_c^d f_1(s) f_2(t) f(s,t) ds dt \\ &= \int_a^b f_1(s) \left\{ p_1(s)(d-c) + \frac{1}{d-c} \int_c^d f_2(t) dt \int_c^d f(s,t) dt \right\} ds \\ &= (d-c) \int_a^b p_1(s) f_1(s) ds + \frac{1}{d-c} \int_c^d f_2(t) dt \int_c^d \left( \int_a^b f_1(s) f(s,t) ds \right) dt \end{aligned}$$

and where

$$\frac{1}{b-a} \int_a^b f_1(s) \cdot f(s,t) ds - \frac{1}{(b-a)^2} \int_a^b f_1(s) ds \int_a^b f(s,t) ds = p_2(t).$$

Therefore

$$\begin{aligned} & \int_a^b \int_c^d f_1(s) f_2(t) f(s,t) ds dt \\ &= (d-c) \int_a^b p_1(s) f_1(s) ds + \frac{1}{(d-c)} \int_c^d f_2(t) dt \\ & \quad \times \int_c^d \left[ (b-a) p_2(t) + \frac{1}{b-a} \int_a^b f_1(s) ds \int_a^b f(s,t) ds \right] dt \\ &= (d-c) \int_a^b p_1(s) f_1(s) ds + \frac{b-a}{(d-c)} \int_c^d f_2(t) dt \int_c^d p_2(t) dt \\ & \quad + \frac{1}{(b-a)(d-c)} \int_c^d f_2(t) dt \int_a^b f_1(s) ds \int_a^b \int_c^d f(s,t) ds dt \end{aligned}$$

and thus

$$\begin{aligned} (9.1) \quad & \left| \int_a^b \int_c^d f_1(s) f_2(t) f(s,t) ds dt \right. \\ & \left. - \frac{1}{(b-a)(d-c)} \int_c^d f_2(t) dt \int_a^b f_1(s) ds \int_a^b \int_c^d f(s,t) ds dt \right| \\ & \leq (d-c) \|p_1\|_\infty \int_a^b |f_1(s)| ds + (b-a) \|p_2\|_\infty \int_c^d |f_2(t)| dt. \end{aligned}$$

**Case 1.** Now, if  $f_1(s) = s$  and  $f_2(t) = t$ , and  $f(\cdot, \cdot)$  is a joint probability density function, the left hand side becomes

$$\begin{aligned} & \left| E(XY) - \frac{1}{4(d-c)(b-a)} (d^2 - c^2)(b^2 - a^2) \right| \\ &= \left| E(XY) - \frac{1}{4}(b+a)(d+c) \right|. \end{aligned}$$

$$\begin{aligned}
|p_1(s)| &\leq \frac{1}{2} \|f\|_\infty \left[ \frac{(d^3 - c^3)}{3(d-c)} - \left( \frac{1}{d-c} \int_c^d t dt \right)^2 \right]^{\frac{1}{2}} \text{ see [4]} \\
&= \frac{1}{2} \|f\|_\infty \left[ \frac{(d-c)(d^2 + dc + c^2)}{3(d-c)} - \left( \frac{(d^2 - c^2)}{2(d-c)} \right)^2 \right]^{\frac{1}{2}} \\
&= \frac{1}{2} \|f\|_\infty \left[ \frac{d^2 + dc + c^2}{3} - \frac{(d+c)^2}{4} \right]^{\frac{1}{2}} \\
&= \frac{1}{2} \|f\|_\infty \times \frac{1}{2\sqrt{3}} [d^2 - 2dc + c^2]^{\frac{1}{2}} = \frac{1}{4\sqrt{3}} \|f\|_\infty (d-c).
\end{aligned}$$

Similarly,

$$p_2 \leq \frac{a^2 - b^2}{2}, \quad a < 0, \quad b < 0$$

Now

$$\int_a^b |f_1(s)| ds = \frac{a^2 + b^2}{2}, \quad a < 0, \quad b > 0$$

and similarly,

$$\int_c^d |f_2(t)| dt \leq \frac{b^2 - a^2}{2}, \quad a > 0, \quad b > 0.$$

Thus, we then have for  $a < 0, b > 0, c < 0, d > 0$ :-

$$\begin{aligned}
&\left| E(XY) - \frac{1}{4}(b+a)(d+c) \right| \\
&\leq \frac{1}{4\sqrt{3}} \|f\|_\infty \frac{(d-c)^2 (a^2 + b^2)}{2} + \frac{1}{4\sqrt{3}} \|f\|_\infty \frac{(b-a)^2 (c^2 + d^2)}{2} \\
&= \frac{1}{8\sqrt{3}} \left[ (d-c)^2 (a^2 + b^2) + (b-a)^2 (c^2 + d^2) \right] \|f\|_\infty \\
&= \frac{1}{4\sqrt{3}} \left[ (a^2 + b^2) (c^2 + d^2) + (ac + db) (ad + bc) \right] \|f\|_\infty.
\end{aligned}$$

**Case 2.** If  $f_1(s) = s$  and  $f_2(t) = 1$ , and  $f(\cdot, \cdot) = t\phi(s, t)$  where  $\phi(\cdot, \cdot)$  is a joint probability density function, then the left hand side is:-

$$\begin{aligned}
&\left| \int_a^b \int_c^d st\phi(s, t) ds dt - \frac{(d-c)(b^2 - a^2)}{2(d-c)(b-a)} E(Y) \right| \\
&= \left| E(XY) - \frac{1}{2}(a+b)E(Y) \right|.
\end{aligned}$$

$p_2$  is as above and

$$p_1 \leq \frac{1}{2} \|f\|_\infty \left[ \frac{1}{d-c} \int_c^d dt - \left( \frac{1}{d-c} \int_c^d dt \right)^2 \right]^{\frac{1}{2}} = 0$$



and hence

$$\left| E(XY) - \frac{1}{2}(a+b)E(Y) \right| \leq \frac{(b-a)^2(c^2+d^2)}{8\sqrt{3}} \|f\|_\infty$$

when  $a < 0, b > 0, c < 0, d > 0$ .

#### REFERENCES

- [1] N.S. BARNETT, S.S. DRAGOMIR and A. SOFO, Better bounds for an inequality of the Ostrowski type with applications, submitted.
- [2] N.S. BARNETT and S.S. DRAGOMIR, Some inequalities for random variables whose probability density functions are absolutely continuous using a pre-Chebyshev inequality, *Tamkang J. Math.*, accepted.
- [3] N.S. BARNETT, P. CERONE and S.S. DRAGOMIR, Further inequalities for the expectation and variance of a random variable defined on a finite interval, submitted.
- [4] M. MATIĆ, J.E. PEČARIĆ and N. UJEVIĆ, On new estimation of the remainder in Taylor's formula, *Math. Ineq. and Applic.*, **3**(2) (1999), 343-361.
- [5] N.S. BARNETT and S.S. DRAGOMIR, Some elementary inequalities for the expectation and variance of a random variable whose p.d.f. is defined on a finite interval, submitted.
- [6] N.S. BARNETT and S.S. DRAGOMIR, An Ostrowski type inequality for double integrals and applications for cubature formulae, *Soochow J. of Math.*, accepted.
- [7] S.S. DRAGOMIR, N.S. BARNETT and P. CERONE, An  $n$  dimensional version of Ostrowski's inequality for mappings of the Hölder type, *Kyungpook Math. J.*, **40**(1) (2000), 65-75.
- [8] S.S. DRAGOMIR, P. CERONE, N.S. BARNETT and J. ROUMELIOTIS, An inequality of the Ostrowski type for double integrals and applications for cubature formulae, *Tamsui Oxford J. Math. Sci.*, **16**(1) (2000), 1-16.

SCHOOL OF COMMUNICATIONS AND INFORMATICS, VICTORIA UNIVERSITY OF TECHNOLOGY, PO BOX 14428, MELBOURNE CITY MC, VICTORIA 8001, AUSTRALIA.

*E-mail address:* [neil@matilda.vu.edu.au](mailto:neil@matilda.vu.edu.au)

*URL:* <http://sci.vu.edu.au/staff/neilb.html>

*E-mail address:* [sever@matilda.vu.edu.au](mailto:sever@matilda.vu.edu.au)

*URL:* <http://rgmia.vu.edu.au/SSDragomirWeb.html>