



VICTORIA UNIVERSITY
MELBOURNE AUSTRALIA

An Algebraic Inequality, II

This is the Published version of the following publication

Guo, Bai-Ni and Qi, Feng (2001) An Algebraic Inequality, II. RGMIA research report collection, 4 (1).

The publisher's official version can be found at

Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/17358/>

AN ALGEBRAIC INEQUALITY, II

BAI-NI GUO AND FENG QI

ABSTRACT. In this article, using inequality between logarithmic mean and one-parameter mean, which can be deduced from monotonicity of the extended mean values, an integral analogue of J. S. Martins' inequality is proved. An open problem is proposed.

1. INTRODUCTION

In [1, 12, 35, 37], double inequalities were proved using mathematical induction and other techniques, which can be expressed as

$$\frac{n}{n+1} \leq \left(\frac{1}{n} \sum_{i=1}^n i^r \middle/ \frac{1}{n+1} \sum_{i=1}^{n+1} i^r \right)^{1/r} \leq \frac{\sqrt[n]{n!}}{\sqrt[n+1]{(n+1)!}}, \quad (1)$$

where $r > 0$ and $n \in \mathbb{N}$.

We call the left-hand side of inequality (1) H. Alzer's inequality [1], and the right-hand side of inequality (1) J. S. Martins' inequality [10]. The inequality between the two ends of (1) is called Minc-Sathre's inequality [11].

In [16], the second author generalised Alzer's inequality and obtained

$$\frac{n+k}{n+m+k} < \left(\frac{1}{n} \sum_{i=k+1}^{n+k} i^r \middle/ \frac{1}{n+m} \sum_{i=k+1}^{n+m+k} i^r \right)^{1/r}, \quad (2)$$

where r is a given positive real number, n and m are natural numbers, and k is a nonnegative integer. The lower bound in (2) is the best possible.

In [3], Martins' inequality was generalised: Let $\{a_i\}_{i=1}^{\infty}$ be an increasing sequence of positive real numbers and

Date: January 16, 2001.

2000 Mathematics Subject Classification. Primary 26D15.

Key words and phrases. Algebraic inequality, extended mean values, logarithmic mean, one-parameter mean.

The authors were supported in part by NSF of Henan Province (#004051800), SF for Pure Research of Natural Sciences of the Education Department of Henan Province (#1999110004), Doctor Fund of Jiaozuo Institute of Technology, and NNSF (#10001016) of China.

This paper was typeset using $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{L}\mathcal{A}\mathcal{T}\mathcal{E}\mathcal{X}$.

(1) for any positive integer $\ell > 1$,

$$\frac{a_{\ell+1}}{a_\ell} \leq \frac{a_\ell}{a_{\ell-1}}; \quad (3)$$

(2) for any positive integer $\ell > 1$,

$$\left(\frac{a_{\ell+1}}{a_\ell} \right)^\ell \geq \left(\frac{a_\ell}{a_{\ell-1}} \right)^{\ell-1}. \quad (4)$$

Then, for any natural numbers n and m , we have

$$\left(\frac{1}{n} \sum_{i=1}^n a_i^r \middle/ \frac{1}{n+m} \sum_{i=1}^{n+m} a_i^r \right)^{1/r} < \frac{\sqrt[n]{a_n!}}{\sqrt[n+m]{a_{n+m}!}}, \quad (5)$$

where $n, m \in \mathbb{N}$ and r is a positive number, $a_i!$ denotes $\prod_{i=1}^n a_i$. The upper bound is best possible.

As a corollary of inequality (5), we have: Let a and b be positive real numbers, k a nonnegative integer, and $m, n \in \mathbb{N}$. Then, for any real number $r > 0$, we have

$$\left(\frac{1}{n} \sum_{i=k+1}^{n+k} (ai+b)^r \middle/ \frac{1}{n+m} \sum_{i=k+1}^{n+m+k} (ai+b)^r \right)^{1/r} < \frac{\sqrt[n]{\prod_{i=k+1}^{n+k} (ai+b)}}{\sqrt[n+m]{\prod_{i=k+1}^{n+m+k} (ai+b)}}. \quad (6)$$

Inequalities (5) and (6) answer an open problem proposed in [15, 16].

The Alzer's inequality and inequality (2) have been generalised and extended by many mathematicians. For more information, please refer to [2, 4, 5, 8, 15, 20, 23, 26, 34]. The Minc-Sathre's inequality was generalised in [8, 15, 18, 20, 24, 26]. In [18, 24, 26], the following inequalities were obtained:

$$\frac{n+k+1}{n+m+k+1} < \left(\prod_{i=k+1}^{n+k} i \right)^{1/n} \middle/ \left(\prod_{i=k+1}^{n+m+k} i \right)^{1/(n+m)} \leq \sqrt{\frac{n+k}{n+m+k}}, \quad (7)$$

where $n, m \in \mathbb{N}$ and k is a nonnegative integer.

In [13], the second author proved the integral analogue of inequality (2): Let $b > a > 0$ and $\delta > 0$ be real numbers. Then, for any given positive $r \in \mathbb{R}$, we have

$$\begin{aligned} & \left(\frac{1}{b-a} \int_a^b x^r dx \middle/ \frac{1}{b+\delta-a} \int_a^{b+\delta} x^r dx \right)^{1/r} \\ &= \left(\frac{b+\delta-a}{b-a} \cdot \frac{b^{r+1}-a^{r+1}}{(b+\delta)^{r+1}-a^{r+1}} \right)^{1/r} \\ &> \frac{b}{b+\delta}. \end{aligned} \quad (8)$$

The lower bound in (8) is the best possible.

Inequality (8) has been generalised to inequalities for a positive functional in [6].

In this paper, by monotonicity of the extended mean values, from which an inequality between the logarithmic mean and one-parameter mean is deduced, we will prove an open problem proposed in [13] as follows:

Theorem 1. *Let $b > a > 0$ and $\delta > 0$ be real numbers. Then, for any positive $r \in \mathbb{R}$, we have*

$$\left(\frac{b + \delta - a}{b - a} \cdot \frac{b^{r+1} - a^{r+1}}{(b + \delta)^{r+1} - a^{r+1}} \right)^{1/r} < \frac{[b^b/a^a]^{1/(b-a)}}{[(b + \delta)^{b+\delta}/a^a]^{1/(b+\delta-a)}}. \quad (9)$$

The upper bound in (9) is the best possible.

At last, we propose an open problem.

2. LEMMAS

In [36], Stolarsky defined the extended mean values $E(r, s; x, y)$ and proved that it is continuous on the domain $\{(r, s; x, y) : r, s \in \mathbb{R}, x, y > 0\}$ as follows

$$E(r, s; x, y) = \left(\frac{r}{s} \cdot \frac{y^s - x^s}{y^r - x^r} \right)^{1/(s-r)}, \quad rs(r-s)(x-y) \neq 0; \quad (10)$$

$$E(r, 0; x, y) = \left(\frac{1}{r} \cdot \frac{y^r - x^r}{\ln y - \ln x} \right)^{1/r}, \quad r(x-y) \neq 0; \quad (11)$$

$$E(r, r; x, y) = e^{-1/r} \left(\frac{x^{x^r}}{y^{y^r}} \right)^{1/(x^r - y^r)}, \quad r(x-y) \neq 0; \quad (12)$$

$$E(0, 0; x, y) = \sqrt{xy}, \quad x \neq y; \quad (13)$$

$$E(r, s; x, x) = x, \quad x = y.$$

Leach and Sholander [9] showed that $E(r, s; x, y)$ are increasing with both r and s , or with both x and y . The monotonicities of E have also been researched by the authors and others in [7, 25] and [27]–[32] using different ideas and simpler methods. In [19], the logarithmic convexity of E was proved.

Most two variable means are special cases of E , for example

$$E(0, 1; x, y) = L(x, y), \quad (14)$$

$$E(r, r+1; x, y) = J_r(x, y), \quad r > 0. \quad (15)$$

They are called the logarithmic mean and the one-parameter mean, respectively, and, by the monotonicity of E , we have

$$J_r(x, y) > L(x, y), \quad r > 0. \quad (16)$$

Recently, the concepts of mean values has been generalised in [14, 17, 21, 22], [27]–[31] and [33].

3. PROOF OF THEOREM 1

Inequality (9) can be rewritten as follows

$$\frac{b^{r+1} - a^{r+1}}{(b-a)(b^b/a^a)^{r/(b-a)}} < \frac{(b+\delta)^{r+1} - a^{r+1}}{(b+\delta-a)((b+\delta)^{b+\delta}/a^a)^{r/(b+\delta-a)}}. \quad (17)$$

Therefore, it suffices to prove the function

$$\phi(x) = \frac{x^{r+1} - a^{r+1}}{(x-a)(x^x/a^a)^{r/(x-a)}} \quad (18)$$

is decreasing in $x > a > 0$.

Taking the logarithm and differentiating yields

$$\begin{aligned} \frac{d \ln \phi(x)}{dx} &= \frac{a^{2+r} + a^{2+r}r - a^{1+r}x - a^{1+r}rx - a^2x^r - a^2rx^r + ax^{1+r} + arx^{1+r}}{(a-x)^2(a^{1+r} - x^{1+r})} \\ &\quad + \frac{rx^{2+r} \ln x - a^{1+r}rx \ln x - arx^{1+r} \ln x + a^{1+r}r \ln(a^{-a}x^x) - rx^{1+r} \ln(a^{-a}x^x)}{(a-x)^2(a^{1+r} - x^{1+r})} \\ &= \frac{a[(1+r)(x-a)(x^r - a^r) + r(x^{1+r} - a^{1+r})(\ln a - \ln x)]}{(a-x)^2(a^{1+r} - x^{1+r})} \\ &= \frac{a(1+r)(x^r - a^r)(\ln a - \ln x) \left(\frac{r(x^{1+r} - a^{1+r})}{(r+1)(x^r - a^r)} - \frac{x-a}{\ln x - \ln a} \right)}{(a-x)^2(a^{1+r} - x^{1+r})} \\ &= \frac{a(1+r)(x^r - a^r)(\ln a - \ln x) (J_r(a, x) - L(a, x))}{(a-x)^2(a^{1+r} - x^{1+r})} \\ &< 0. \end{aligned} \quad (19)$$

Thus, the function $\ln \phi(x)$, and $\phi(x)$, is decreasing.

Here L'Hospital's rule yields

$$\lim_{r \rightarrow 0^+} \left(\frac{b+\delta-a}{b-a} \cdot \frac{b^{r+1} - a^{r+1}}{(b+\delta)^{r+1} - a^{r+1}} \right)^{1/r} = \frac{[b^b/a^a]^{1/(b-a)}}{[(b+\delta)^{b+\delta}/a^a]^{1/(b+\delta-a)}}. \quad (20)$$

Hence, the upper bound in inequality (9) is the best possible. The proof is complete.

4. OPEN PROBLEM

Lastly, we propose the following

Open Problem. Let $b > a > 0$ and $\delta > 0$ be real numbers, $f(x)$ a positive integrable function. Then, for any given positive $r \in \mathbb{R}$, we have

$$\begin{aligned} \frac{\sup_{x \in [a, b]} f(x)}{\sup_{x \in [a, b+\delta]} f(x)} &< \left(\frac{1}{b-a} \int_a^b f^r(x) dx \Big/ \frac{1}{b+\delta-a} \int_a^{b+\delta} f^r(x) dx \right)^{1/r} \\ &< \exp \left(\frac{\int_a^b \ln f(x) dx}{b-a} - \frac{\int_a^{b+\delta} \ln f(x) dx}{b+\delta-a} \right). \end{aligned} \quad (21)$$

The lower and upper bounds in (21) are the best possible.

REFERENCES

- [1] H. Alzer, *On an inequality of H. Minc and L. Sathre*, J. Math. Anal. Appl. **179** (1993), 396–402.
- [2] P. S. Bullen, *A Dictionary of Inequalities*, Pitman Monographs and Surveys in Pure and Applied Mathematics **97**, Addison Wesley Longman Limited, 1998.
- [3] Tsz Ho Chan, Peng Gao and Feng Qi, *On a generalization of Martins' inequality*, RGMIA Research Report Collection **4** (2001), no.1. <http://rgmia.vu.edu.au/v4n1.html>.
- [4] S. S. Dragomir and J. van der Hoek, *Some new analytic inequalities and their applications in guessing theory*, J. Math. Anal. Appl. **225** (1998), 542–556.
- [5] N. Elezović and J. Pečarić, *On Alzer's inequality*, J. Math. Anal. Appl. **223** (1998), 366–369.
- [6] B. Gavrea and I. Gavrea, *An inequality for linear positive functionals*, J. Ineq. Pure Appl. Math. **1** (2000), no. 1, Article 5. http://jipam.vu.edu.au/v1n1/004_99.html.
- [7] B.-N. Guo, S.-Q. Zhang, and F. Qi, *Elementary proofs of monotonicity for extended mean values of some functions with two parameters*, Shuxue de Shijian yu Renshi (Mathematics in Practice and Theory) **29** (1999), no.2, 169–174. (Chinese)
- [8] J.-C. Kuang, *Some extensions and refinements of Minc-Sathre inequality*, Math. Gaz. **83** (1999), 123–127.
- [9] E. Leach and M. Sholander, *Extended mean values*, Amer. Math. Monthly **85** (1978), 84–90.
- [10] J. S. Martins, *Arithmetic and geometric means, an applications to Lorentz sequence spaces*, Math Nachr. **139** (1988), 281–288.
- [11] H. Minc and L. Sathre, *Some inequalities involving $(n!)^{1/r}$* , Proc. Edinburgh Math. Soc. **14** (1964/65), 41–46.
- [12] N. Ozeki, *On some inequalities*, J. College Arts Sci. Chiba Univ. **4** (1965), no. 3, 211–214. (Japanese)
- [13] F. Qi, *An algebraic inequality*, J. Ineq. Pure Appl. Math. **1** (2001), in the press. <http://jipam.vu.edu.au>. RGMIA Res. Rep. Coll. **2** (1999), no. 1, Article 8, 81–83. <http://rgmia.vu.edu.au/v2n1.html>.
- [14] F. Qi, *Generalized abstracted mean values*, J. Ineq. Pure Appl. Math. **1** (2000), no. 1, Article 4. <http://jipam.vu.edu.au/v1n1/013.99.html>. RGMIA Res. Rep. Coll. **2** (1999), no. 5, Article 4, 633–642. <http://rgmia.vu.edu.au/v2n5.html>.
- [15] F. Qi, *Generalizations of Alzer's and Kuang's inequality*, Tamkang J. Math. **31** (2000), no. 3, 223–227. RGMIA Res. Rep. Coll. **2** (1999), no. 6, Article 12. <http://rgmia.vu.edu.au/v2n6.html>.
- [16] F. Qi, *Generalization of H. Alzer's inequality*, J. Math. Anal. Appl. **240** (1999), 294–297.
- [17] F. Qi, *Generalized weighted mean values with two parameters*, Proc. Roy. Soc. London Ser. A **454** (1998), no. 1978, 2723–2732.
- [18] F. Qi, *Inequalities and monotonicity of sequences involving $\sqrt[n]{(n+k)!/k!}$* , RGMIA Res. Rep. Coll. **2** (1999), no. 5, Article 8, 685–692. <http://rgmia.vu.edu.au/v2n5.html>.
- [19] F. Qi, *Logarithmic convexity of extended mean values*, Proc. Amer. Math. Soc. (2001), accepted. RGMIA Res. Rep. Coll. **2** (1999), no. 5, Article 5, 643–652. <http://rgmia.vu.edu.au/v2n5.html>.

- [20] F. Qi and B.-N. Guo, *Monotonicity of sequences involving convex function and sequence*, RGMIA Res. Rep. Coll. **3** (2000), no. 2, Article 14. <http://rgmia.vu.edu.au/v3n2.html>.
- [21] F. Qi, *On a two-parameter family of nonhomogeneous mean values*, Tamkang J. Math. **29** (1998), no. 2, 155–163.
- [22] F. Qi, *Studies on Problems in Topology and Geometry and on Generalized Weighted Abstracted Mean Values*, Thesis submitted for the degree of Doctor of Philosophy at University of Science and Technology of China, Hefei City, Anhui Province, China, Winter 1998. (Chinese)
- [23] F. Qi and L. Debnath, *On a new generalization of Alzer's inequality*, Internat. J. Math. & Math. Sci. **23** (2000), no. 12, 815–818.
- [24] F. Qi and B.-N. Guo, *Some inequalities involving the geometric mean of natural numbers and the ratio of gamma functions*, RGMIA Res. Rep. Coll. **4** (2001), no. 1, Article 3. <http://rgmia.vu.edu.au/v4n1.html>.
- [25] F. Qi and Q.-M. Luo, *A simple proof of monotonicity for extended mean values*, J. Math. Anal. Appl. **224** (1998), no. 2, 356–359.
- [26] F. Qi and Q.-M. Luo, *Generalization of H. Minc and J. Sathre's inequality*, Tamkang J. Math. **31** (2000), no. 2, 145–148. RGMIA Res. Rep. Coll. **2** (1999), no. 6, Article 14. <http://rgmia.vu.edu.au/v2n6.html>.
- [27] F. Qi and Q.-M. Luo, *Refinements and extensions of an inequality*, Mathematics and Informatics Quarterly **9** (1999), no. 1, 23–25.
- [28] F. Qi, J.-Q. Mei, Da-F. Xia, and S.-L. Xu, *New proofs of weighted power mean inequalities and monotonicity for generalized weighted mean values*, Math. Ineq. Appl. **3** (2000), no. 3, 377–383.
- [29] F. Qi, J.-Q. Mei, and S.-L. Xu, *Other proofs of monotonicity for generalized weighted mean values*, RGMIA Res. Rep. Coll. **2** (1999), no. 4, Article 6, 469–472. <http://rgmia.vu.edu.au/v2n4.html>.
- [30] F. Qi and S.-L. Xu, *Refinements and extensions of an inequality*, II, J. Math. Anal. Appl. **211** (1997), 616–620.
- [31] F. Qi and S.-L. Xu, *The function $(b^x - a^x)/x$: Inequalities and properties*, Proc. Amer. Math. Soc. **126** (1998), no. 11, 3355–3359.
- [32] F. Qi, S.-L. Xu, and L. Debnath, *A new proof of monotonicity for extended mean values*, Internat. J. Math. & Math. Sci. **22** (1999), no. 2, 415–420.
- [33] F. Qi and S.-Q. Zhang, *Note on monotonicity of generalized weighted mean values*, Proc. Roy. Soc. London Ser. A **455** (1999), no. 1989, 3259–3260.
- [34] J. Sándor, *Comments on an inequality for the sum of powers of positive numbers*, RGMIA Res. Rep. Coll. **2** (1999), no. 2, 259–261. <http://rgmia.vu.edu.au/v2n2.html>.
- [35] J. Sándor, *On an inequality of Alzer*, J. Math. Anal. Appl. **192** (1995), 1034–1035.
- [36] K. B. Stolarsky, *Generalizations of the logarithmic mean*, Math. Mag. **48** (1975), 87–92.
- [37] J. S. Ume, *An elementary proof of H. Alzer's inequality*, Math. Japon. **44** (1996), no. 3, 521–522.

DEPARTMENT OF MATHEMATICS, JIAOZUO INSTITUTE OF TECHNOLOGY, JIAOZUO CITY, HENAN
454000, THE PEOPLE'S REPUBLIC OF CHINA

DEPARTMENT OF MATHEMATICS, JIAOZUO INSTITUTE OF TECHNOLOGY, JIAOZUO CITY, HENAN
454000, THE PEOPLE'S REPUBLIC OF CHINA

E-mail address: qifeng@jzit.edu.cn

URL: <http://rgmia.vu.edu.au/qi.html>