



**VICTORIA UNIVERSITY**  
MELBOURNE AUSTRALIA

## *A Two Points Taylor's Formula for the Generalised Riemann Integral*

This is the Published version of the following publication

Dragomir, Sever S and Thompson, H. B (2008) A Two Points Taylor's Formula for the Generalised Riemann Integral. Research report collection, 11 (4).

The publisher's official version can be found at

Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/17655/>

# A TWO POINTS TAYLOR'S FORMULA FOR THE GENERALISED RIEMANN INTEGRAL

S.S. DRAGOMIR AND H.B. THOMPSON

ABSTRACT. A two points Taylor's formula for the generalised Riemann integral and various bounds for the remainder are established. Moreover, particular instances of interest are given.

## 1. INTRODUCTION

The generalised Riemann integral is variously known as the Kurzweil, the Riemann complete, and the gauge integral. It is also equivalent to the Perron integral, the descriptive  $D^*$ -integral of Luzin, and the restricted total integral (also called the  $T^*$ -integral) of Denjoy.

Newton introduced integration as antidifferentiation. Between 1912 and 1915, Denjoy [3], Luzin [10], and Perron [15], realising that the Lebesgue and Newton integrals did not properly contain one another, gave new definitions of the integral to encompass both the Newton and Lebesgue integrals. The equivalence of the Denjoy and Luzin integrals is not difficult to prove while the equivalence of these to the Perron integral is due to Hake [4], Looman [9], and Aleksandrov [1].

Kurzweil [7] introduced his integral for application to ordinary differential equations, and showed that it is equivalent to the Perron integral. Henstock [5] independently introduced this integral and developed its properties (see, e.g., [6]).

By a *tagged partition*  $\mathbf{T}$  of  $[a, b]$  we mean a set  $\{x_0, x_1, \dots, x_n; t_1, t_2, \dots, t_n\}$  satisfying

$$a = x_0 \leq t_1 \leq x_1 \leq t_2 \leq x_2 \leq \dots \leq x_n = b$$

for some  $n > 0$ . A positive function  $\delta : [a, b] \rightarrow \mathbb{R}^+ = (0, \infty)$  is called a *gauge* on  $[a, b]$ . Let  $\delta$  be a gauge on  $[a, b]$ . Then the partition  $\mathbf{T}$  is said to be  $\delta$ -fine if

$$[x_{i-1}, x_i] \subseteq (t_i - \delta(t_i), t_i + \delta(t_i))$$

for  $i = 1, 2, \dots, n$ .

Using bisection and the nested interval theorem it is easy to prove that for every gauge  $\delta$  on  $[a, b]$  there exists a  $\delta$ -fine partition of  $[a, b]$ .

**Definition 1.** Let  $f : [a, b] \rightarrow \mathbb{R}$ . Then  $I$  is said to be the *generalised Riemann integral* of  $f$  on  $[a, b]$  (denoted by  $\int_a^b f(t) dt$ ) if, given  $\epsilon > 0$ , there exists a gauge  $\delta$  on  $[a, b]$  such that

$$\left| \sum_{i=1}^n f(t_i)(x_i - x_{i-1}) - I \right| < \epsilon,$$

---

2000 *Mathematics Subject Classification.* Primary 41A55, Secondary 26D15, 26D10.

*Key words and phrases.* Generalised Riemann integral, Kurzweil integral, Perron integral, Taylor's formula, Integral inequalities,

whenever the partition  $\mathbf{T}$  is  $\delta$ -fine. We call  $f$  integrable on  $[a, b]$  if its generalised Riemann integral exists.

We are ready to state the fundamental theorem and its associated theorem on integration by parts.

**Definition 2.** (See [14]) Let  $f : [a, b] \rightarrow \mathbb{R}$  be given. A function  $F : [a, b] \rightarrow \mathbb{R}$  is a *primitive* of  $f$  on  $[a, b]$  provided  $F$  is continuous on  $[a, b]$  and  $F'(x) = f(x)$  for all  $x$  in  $[a, b]$ , except possibly at a finite or countably infinite set of values of  $x$ .

**Theorem 1.** (The fundamental theorem, [18, Theorem 5]) If  $f$  has a primitive  $F$  on  $[a, b]$ , then  $f$  is integrable and

$$(1.1) \quad \int_a^b f(t) dt = F(b) - F(a).$$

**Remark 1.** To guarantee the validity of (1.1) and of the integral form of Taylor's theorem in the case of the Riemann or Lebesgue integrals, additional assumptions such as the integrability of  $f$  are required. In particular, there is a function  $F$  having a bounded derivative everywhere on  $[a, b]$  but such that  $f = F'$  is not Riemann integrable on  $[a, b]$ . Also, the function  $F$  defined by  $F(x) = x^2 \sin 1/x^2$ , for  $x \neq 0$ ,  $F(0) = 0$  is differentiable everywhere but  $f = F'$  is not Lebesgue integrable on  $[a, b]$  if  $a \neq b$  and  $0 \in [a, b]$ .

As an immediate consequence of the fundamental theorem, one obtains the following.

**Theorem 2.** (Integration by parts; See [14]) If  $g$  and  $h$  have primitives  $G$  and  $H$ , respectively, on  $[a, b]$ , then  $gH$  is integrable if and only if  $Gh$  is integrable. Moreover

$$(1.2) \quad \int_a^b g(t)H(t) dt = G(b)H(b) - G(a)H(a) - \int_a^b G(t)h(t) dt.$$

**Remark 2.** The integrability of  $gH$  and hence of  $Gh$  is necessary for (1.2) to hold as can be seen by setting

$$F(x) = x^2 \sin x^{-4}, G(x) = x^2 \cos x^{-4} \text{ for } x \neq 0 \text{ and } F(0) = 0 = G(0).$$

See [11, Ex. 13]. In our case  $g$  will be continuous on  $[a, b]$  so  $gH$  will be integrable on  $[a, b]$ .

We also recall Taylor's theorem for the generalised Riemann integral obtained in [21]:

**Lemma 1.** Let  $f, f^{(1)}, \dots, f^{(n)}$  be continuous on  $[\alpha, \beta]$  and suppose that  $f^{(n+1)}$  exists on  $[\alpha, \beta]$ , except possibly at a countable number of points. Then

$$(1.3) \quad f(\beta) = \sum_{k=0}^n \frac{1}{k!} f^{(k)}(\alpha) (\beta - \alpha)^k + R_{n,\alpha}(\beta),$$

where

$$(1.4) \quad R_{n,\alpha}(\beta) = \frac{1}{n!} \int_{\alpha}^{\beta} f^{(n+1)}(t) (\beta - t)^n dt$$

and the integral in (1.4) is taken in the generalised Riemann sense.

The main aim of this paper is to provide a two points Taylor's formula for the generalised Riemann integral and establish various bounds for the remainder. Particular instances of interest also will be given.

## 2. IDENTITIES

The following identity can be stated:

**Theorem 3.** *Let  $f, f^{(1)}, \dots, f^{(n)}$  be continuous on  $[a, b]$  and suppose that  $f^{(n+1)}$  exists on  $[a, b]$ , except possibly at a countable number of points. Then for any  $x \in [a, b]$  and for any  $\lambda \in [0, 1]$  we have the representation*

(2.1)

$$f(x) = \lambda f(a) + (1 - \lambda) f(b) + \sum_{k=1}^n \frac{1}{k!} \left[ \lambda f^{(k)}(a) (x - a)^k + (-1)^k (1 - \lambda) f^{(k)}(b) (b - x)^k \right] + S_{n,\lambda}(x),$$

where the remainder  $S_{n,\lambda}(x)$  is given by

$$(2.2) \quad S_{n,\lambda}(x) := \frac{1}{n!} \int_a^b f^{(n+1)}(t) K_{n,\lambda}(x, t) dt$$

and the kernel  $K_{n,\lambda}(\cdot, \cdot) : [a, b]^2 \rightarrow \mathbb{R}$  is defined by

$$(2.3) \quad K_{n,\lambda}(x, t) := \begin{cases} \lambda (x - t)^n & \text{if } a \leq t \leq x, \\ (-1)^{n+1} (1 - \lambda) (t - x)^n & \text{if } x < t \leq b. \end{cases}$$

*Proof.* Using Lemma 1 we can write the following two identities

$$(2.4) \quad f(x) = \sum_{k=0}^n \frac{1}{k!} f^{(k)}(a) (x - a)^k + \frac{1}{n!} \int_a^x f^{(n+1)}(t) (x - t)^n dt$$

and

$$(2.5) \quad f(x) = \sum_{k=0}^n \frac{(-1)^k}{k!} f^{(k)}(b) (b - x)^k + \frac{(-1)^{n+1}}{n!} \int_x^b f^{(n+1)}(t) (t - x)^n dt$$

for each  $x \in [a, b]$ .

Now, if we multiply (2.4) with  $\lambda$  and (2.5) with  $(1 - \lambda)$  and add the resulting equalities, a simple calculation yields the desired identity (2.1). ■

**Corollary 1.** *With the assumptions in Theorem 3 we have for each  $x \in [a, b]$*

$$\begin{aligned} f(x) &= \frac{1}{b - a} [(b - x) f(a) + (x - a) f(b)] \\ &+ \frac{(b - x)(x - a)}{b - a} \cdot \sum_{k=1}^n \frac{1}{k!} \left\{ (x - a)^{k-1} f^{(k)}(a) + (-1)^k (b - x)^{k-1} f^{(k)}(b) \right\} \\ &+ \frac{1}{n! (b - a)} \int_a^b L_n(x, t) f^{(n+1)}(t) dt, \end{aligned}$$

where

$$L_n(x, t) = \begin{cases} (x - t)^n (b - x) & \text{if } a \leq t \leq x, \\ (-1)^{n+1} (t - x)^n (x - a) & \text{if } x < t \leq b, \end{cases}$$

and

$$\begin{aligned} f(x) &= \frac{1}{b-a} [(x-a)f(a) + (b-x)f(b)] \\ &+ \frac{1}{b-a} \sum_{k=1}^n \frac{1}{k!} \left\{ (x-a)^{k+1} f^{(k)}(a) + (-1)^k (b-x)^{k+1} f^{(k)}(b) \right\} \\ &+ \frac{1}{n!(b-a)} \int_a^b P_n(x, t) f^{(n+1)}(t) dt, \end{aligned}$$

where

$$P_n(x, t) = \begin{cases} (x-t)^n (x-a) & \text{if } a \leq t \leq x, \\ (-1)^{n+1} (t-x)^n (b-x) & \text{if } x < t \leq b, \end{cases}$$

respectively.

The proof is obvious. Choose  $\lambda = (b-x)/(b-a)$  and  $\lambda = (x-a)/(b-a)$ , respectively, in Theorem 3. The details are omitted.

**Remark 3.** We observe that each of the identities from Corollary 1 provide the possibility to approximate the value of a function at the midpoint in terms of its values at the end points as well as in terms of the values of its derivatives at the same points. To be more precise, we can state the identity

$$\begin{aligned} f\left(\frac{a+b}{2}\right) &= \frac{f(a) + f(b)}{2} + \sum_{k=1}^n \frac{1}{2^{k+1}k!} \left[ f^{(k)}(a) + (-1)^k f^{(k)}(b) \right] (b-a)^k \\ &+ \frac{1}{2 \cdot n!} \int_a^b M_n(t) f^{(n+1)}(t) dt, \end{aligned}$$

where

$$M_n(t) = \begin{cases} \left(\frac{a+b}{2} - t\right)^n & \text{if } a \leq t \leq \frac{a+b}{2}, \\ (-1)^{n+1} \left(t - \frac{a+b}{2}\right)^n & \text{if } \frac{a+b}{2} < t \leq b. \end{cases}$$

**Corollary 2.** With the assumption in Theorem 3 we have for each  $\lambda \in [0, 1]$

$$\begin{aligned} f[\lambda a + (1-\lambda)b] &= \lambda f(a) + (1-\lambda)f(b) \\ &+ \lambda(1-\lambda) \sum_{k=1}^n \frac{1}{k!} \left[ (1-\lambda)^{k-1} f^{(k)}(a) + (-1)^k \lambda^{k-1} f^{(k)}(b) \right] (b-a)^k \\ &+ \frac{1}{n!} \int_a^b f^{(n+1)}(t) K_{n,\lambda}(t) dt, \end{aligned}$$

where

$$K_{n,\lambda}(t) := \begin{cases} \lambda [\lambda a + (1-\lambda)b - t]^n & \text{if } a \leq t \leq \lambda a + (1-\lambda)b, \\ (-1)^{n+1} (1-\lambda) \{t - [\lambda a + (1-\lambda)b]\}^n & \text{if } \lambda a + (1-\lambda)b < t \leq b. \end{cases}$$

**Remark 4.** To the best of our knowledge the representation results from this section are new even in the non Kurzweil setting.

## 3. UPPER AND LOWER BOUNDS FOR THE REMAINDER

Consider the polynomials

$$(3.1) \quad T_{n,\lambda}(x) := \sum_{k=1}^n \frac{1}{k!} \left[ \lambda f^{(k)}(a) (x-a)^k + (-1)^k (1-\lambda) f^{(k)}(b) (b-x)^k \right],$$

where  $x \in [a, b]$  and  $\lambda \in [0, 1]$ .

When upper and lower bounds for the  $(n+1)^{\text{th}}$  derivative of the function  $f$  are available, we may state the following result.

**Theorem 4.** *Assume that  $f : [a, b] \rightarrow \mathbb{R}$  is such that  $f^{(1)}, \dots, f^{(n)}$  are continuous on  $[a, b]$  and  $f^{(n+1)}$  exists, except possibly at a countable number of points of  $[a, b]$ . Assume that for  $x \in (a, b)$  there exists the constants  $\gamma_{n+1}^{(i)}(x)$ ,  $\Gamma_{n+1}^{(i)}(x)$ ,  $i \in \{1, 2\}$  so that*

$$(3.2) \quad \gamma_{n+1}^{(1)}(x) \leq f^{(n+1)}(t) \leq \Gamma_{n+1}^{(1)}(x) \quad \text{for } t \in [a, x]$$

and

$$(3.3) \quad \gamma_{n+1}^{(2)}(x) \leq f^{(n+1)}(t) \leq \Gamma_{n+1}^{(2)}(x) \quad \text{for } t \in [x, b].$$

If  $n = 2m - 1$  ( $m \geq 1$ ), then

$$(3.4) \quad \begin{aligned} & \frac{1}{(2m)!} \left[ \lambda \gamma_{2m}^{(1)}(x) (x-a)^{2m} + (1-\lambda) \gamma_{2m}^{(2)}(x) (b-x)^{2m} \right] \\ & \leq f(x) - T_{2m-1,\lambda}(x) \\ & \leq \frac{1}{(2m)!} \left[ \lambda \Gamma_{2m}^{(1)}(x) (x-a)^{2m} + (1-\lambda) \Gamma_{2m}^{(2)}(x) (b-x)^{2m} \right], \end{aligned}$$

for any  $\lambda \in [0, 1]$ .

If  $n = 2m$  ( $m \geq 1$ ), then

$$(3.5) \quad \begin{aligned} & \frac{1}{(2m+1)!} \left[ \lambda \gamma_{2m+1}^{(1)}(x) (x-a)^{2m+1} - (1-\lambda) \Gamma_{2m+1}^{(2)}(x) (b-x)^{2m+1} \right] \\ & \leq f(x) - T_{2m,\lambda}(x) \\ & \leq \frac{1}{(2m+1)!} \left[ \lambda \Gamma_{2m+1}^{(1)}(x) (x-a)^{2m+1} - (1-\lambda) \gamma_{2m+1}^{(2)}(x) (b-x)^{2m+1} \right] \end{aligned}$$

for any  $\lambda \in [0, 1]$ .

*Proof.* For  $n = 2m - 1$ , we have the representation

$$(3.6)$$

$$\begin{aligned} & f(x) - T_{2m-1,\lambda}(x) \\ & = \frac{1}{(2m-1)!} \left[ \lambda \int_a^x (x-t)^{2m-1} f^{(2m)}(t) dt + (1-\lambda) \int_x^b (t-x)^{2m-1} f^{(2m)}(t) dt \right] \end{aligned}$$

for any  $x \in [a, b]$  and  $\lambda \in [0, 1]$ . Using assumptions (3.2) and (3.3) for  $n = 2m - 1$ , we get

$$(3.7) \quad \begin{aligned} \gamma_{2m}^{(1)}(x) \cdot \frac{(x-a)^{2m}}{2m} & \leq \int_a^x (x-t)^{2m-1} f^{(2m)}(t) dt \\ & \leq \Gamma_{2m}^{(1)}(x) \cdot \frac{(x-a)^{2m}}{2m} \end{aligned}$$

and

$$(3.8) \quad \gamma_{2m}^{(2)}(x) \cdot \frac{(b-x)^{2m}}{2m} \leq \int_x^b (t-x)^{2m-1} f^{(2m)}(t) dt \\ \leq \Gamma_{2m}^{(2)}(x) \cdot \frac{(b-x)^{2m}}{2m}.$$

Using (3.6) – (3.8) we easily deduce (3.4).

For  $n = 2m$ , we have the representation

$$(3.9) \quad f(x) - T_{2m,\lambda}(x) \\ = \frac{1}{(2m)!} \left[ \lambda \int_a^x (x-t)^{2m} f^{(2m+1)}(t) dt - (1-\lambda) \int_x^b (t-x)^{2m} f^{(2m+1)}(t) dt \right].$$

On making use of assumptions (3.2) and (3.3) for  $n = 2m$ , we get

$$(3.10) \quad \gamma_{2m+1}^{(1)}(x) \cdot \frac{(x-a)^{2m+1}}{2m+1} \leq \int_a^x (x-t)^{2m} f^{(2m+1)}(t) dt \\ \leq \Gamma_{2m+1}^{(1)}(x) \cdot \frac{(x-a)^{2m+1}}{2m+1}$$

and

$$(3.11) \quad -\Gamma_{2m+1}^{(2)}(x) \cdot \frac{(b-x)^{2m+1}}{2m+1} \leq -\int_x^b (t-x)^{2m} f^{(2m+1)}(t) dt \\ \leq -\gamma_{2m+1}^{(2)}(x) \cdot \frac{(b-x)^{2m+1}}{2m+1}.$$

Finally, the identity (3.9) and the inequalities (3.10) and (3.11) yield the desired result (3.5). The details are omitted. ■

When global bounds for the  $(n+1)^{\text{th}}$  derivative are available, the following more convenient result may be stated.

**Corollary 3.** *Under the assumptions of Theorem 4, if there exist constants  $\gamma_{n+1}$ ,  $\Gamma_{n+1}$  so that*

$$(3.12) \quad -\infty < \gamma_{n+1} \leq f^{(n+1)}(t) \leq \Gamma_{n+1} < \infty \quad \text{for } t \in [a, b],$$

then for  $n = 2m - 1$  ( $m \geq 1$ ) we have

$$(3.13) \quad \frac{\gamma_{2m}}{(2m)!} \left[ \lambda (x-a)^{2m} + (1-\lambda) (b-x)^{2m} \right] \\ \leq f(x) - T_{2m-1,\lambda}(x) \\ \leq \frac{\Gamma_{2m}}{(2m)!} \left[ \lambda (x-a)^{2m} + (1-\lambda) (b-x)^{2m} \right]$$

for any  $x \in [a, b]$  and for any  $\lambda \in [0, 1]$  while for  $n = 2m$  ( $m \geq 1$ ) we have

$$(3.14) \quad \begin{aligned} & \frac{1}{(2m+1)!} \left[ \lambda \gamma_{2m+1} (x-a)^{2m+1} - (1-\gamma) \Gamma_{2m+1} (b-x)^{2m+1} \right] \\ & \leq f(x) - T_{2m,\lambda}(x) \\ & \leq \frac{1}{(2m+1)!} \left[ \lambda \Gamma_{2m+1} (x-a)^{2m+1} - (1-\gamma) \gamma_{2m+1} (b-x)^{2m+1} \right] \end{aligned}$$

for each  $x \in [a, b]$  and  $\lambda \in [0, 1]$ .

Now, let us consider the polynomials

$$(3.15) \quad \begin{aligned} P_n(x) &= \frac{1}{b-a} [(b-x)f(a) + (x-a)f(b)] \\ &+ \frac{(b-x)(x-a)}{b-a} \cdot \sum_{k=1}^n \frac{1}{k!} \left\{ (x-a)^{k-1} f^{(k)}(a) + (-1)^k (b-x)^{k-1} f^{(k)}(b) \right\} \end{aligned}$$

and

$$(3.16) \quad \begin{aligned} Q_n(x) &= \frac{1}{b-a} [(x-a)f(a) + (b-x)f(b)] \\ &+ \frac{1}{b-a} \sum_{k=1}^n \frac{1}{k!} \left\{ (x-a)^{k+1} f^{(k)}(a) + (-1)^k (b-x)^{k+1} f^{(k)}(b) \right\} \end{aligned}$$

which are obtained from  $T_{n,\lambda}(x)$  by choosing  $\lambda = (b-x)/(b-a)$  and  $\lambda = (x-a)/(b-a)$ , respectively. Then we may state the following additional result.

**Corollary 4.** *Under the assumptions of Theorem 4, if there exist constants  $\gamma_{n+1}$ ,  $\Gamma_{n+1}$  so that (3.12) is valid, then for  $n = 2m-1$  ( $m \geq 1$ ) we have*

$$(3.17) \quad \begin{aligned} & \frac{\gamma_{2m}}{(2m)!} \cdot \frac{(x-a)(b-x)}{b-a} \left[ (x-a)^{2m-1} + (b-x)^{2m-1} \right] \\ & \leq f(x) - P_{2m-1}(x) \\ & \leq \frac{\Gamma_{2m}}{(2m)!} \cdot \frac{(x-a)(b-x)}{b-a} \left[ (x-a)^{2m-1} + (b-x)^{2m-1} \right] \end{aligned}$$

for any  $x \in [a, b]$ , while for  $n = 2m$  ( $m \geq 1$ ) we have

$$(3.18) \quad \begin{aligned} & \frac{1}{(2m+1)!} \cdot \frac{(x-a)(b-x)}{b-a} \left[ \gamma_{2m+1} (x-a)^{2m} - \Gamma_{2m+1} (b-x)^{2m} \right] \\ & \leq f(x) - P_{2m}(x) \\ & \leq \frac{1}{(2m+1)!} \cdot \frac{(x-a)(b-x)}{b-a} \left[ \Gamma_{2m+1} (x-a)^{2m} - \gamma_{2m+1} (b-x)^{2m} \right] \end{aligned}$$

for each  $x \in [a, b]$ .

Also, for  $n = 2m-1$  ( $m \geq 1$ ) we have

$$(3.19) \quad \begin{aligned} & \frac{\gamma_{2m}}{(2m)!(b-a)} \left[ (x-a)^{2m+1} + (b-x)^{2m+1} \right] \\ & \leq f(x) - Q_{2m-1}(x) \\ & \leq \frac{\Gamma_{2m}}{(2m)!(b-a)} \left[ (x-a)^{2m+1} + (b-x)^{2m+1} \right] \end{aligned}$$



for any  $x \in [a, b]$ , while for  $n = 2m$  ( $m \geq 1$ ) we have

$$(3.20) \quad \begin{aligned} & \frac{1}{(2m+1)!(b-a)} \left[ \gamma_{2m+1}(x-a)^{2m+2} - \Gamma_{2m+1}(b-x)^{2m+2} \right] \\ & \leq f(x) - Q_{2m}(x) \\ & \leq \frac{1}{(2m+1)!(b-a)} \left[ \Gamma_{2m+1}(x-a)^{2m+2} - \gamma_{2m+1}(b-x)^{2m+2} \right] \end{aligned}$$

for any  $x \in [a, b]$ .

**Remark 5.** Assume  $f : [a, b] \rightarrow \mathbb{R}$  is  $2m$ -differentiable. If  $f$  is  $2m$ -convex, that is,  $f^{(2m)}(t) \geq 0$  for any  $t \in (a, b)$ , it follows from (2.1) - (2.3) that

$$(3.21) \quad \begin{aligned} f(x) & \geq \lambda f(a) + (1-\lambda)f(b) \\ & + \sum_{k=1}^{2m-1} \frac{1}{k!} \left[ \lambda(x-a)^k f^{(k)}(a) + (-1)^k (1-\lambda) f^{(k)}(b) (b-x)^k \right] \end{aligned}$$

for any  $x \in [a, b]$  and  $\lambda \in [0, 1]$ . Moreover, if we choose  $x = \lambda a + (1-\lambda)b$  in (2.1), then we get the inequality

$$\begin{aligned} f(\lambda a + (1-\lambda)b) & \geq \lambda f(a) + (1-\lambda)f(b) \\ & + \lambda(1-\lambda) \sum_{k=1}^{2m-1} \frac{1}{k!} \left[ (1-\lambda)^{k-1} f^{(k)}(a) + (-1)^k \lambda^{k-1} f^{(k)}(b) \right] (b-a)^k \end{aligned}$$

that holds for any  $\lambda \in [0, 1]$ .

#### 4. ERROR BOUNDS IN TERMS OF $p$ -NORMS

Moreover, the following result providing error bounds for the approximation of  $f$  in terms of the polynomials

$$(4.1) \quad T_{n,\lambda}(x) := \sum_{k=1}^n \frac{1}{k!} \left[ \lambda f^{(k)}(a) (x-a)^k + (-1)^k (1-\lambda) f^{(k)}(b) (b-x)^k \right]$$

may be stated.

**Theorem 5.** Assume that  $f : [a, b] \rightarrow \mathbb{R}$  is such that  $f^{(1)}, \dots, f^{(n)}$  are continuous on  $[a, b]$  and  $f^{(n+1)}$  exists, except possibly at a countable number of points of  $[a, b]$ .

Then, for any  $x \in [a, b]$  and  $\lambda \in [0, 1]$ , we have

$$(4.2) \quad |f(x) - T_{n,\lambda}(x)| \leq \frac{\lambda}{n!} \times \begin{cases} \frac{(x-a)^{n+1}}{n+1} \|f^{(n+1)}\|_{[a,x],\infty} & \text{if } f^{(n+1)} \in L_\infty[a, x], \\ \frac{(x-a)^{n+\frac{1}{q}}}{(nq+1)^{\frac{1}{q}}} \|f^{(n+1)}\|_{[a,x],p} & \text{if } f^{(n+1)} \in L_p[a, x] \\ & \text{for } p > 1, \frac{1}{p} + \frac{1}{q} = 1, \\ (x-a)^n \|f^{(n+1)}\|_{[a,x],1} & \text{if } f^{(n+1)} \in L_1[a, x], \end{cases} \\ + \frac{(1-\lambda)}{n!} \times \begin{cases} \frac{(b-x)^{n+1}}{n+1} \|f^{(n+1)}\|_{[x,b],\infty} & \text{if } f^{(n+1)} \in L_\infty[x, b], \\ \frac{(b-x)^{n+\frac{1}{\beta}}}{(n\beta+1)^{\frac{1}{\beta}}} \|f^{(n+1)}\|_{[x,b],\alpha} & \text{if } f^{(n+1)} \in L_\alpha[x, b] \\ & \text{for } \alpha > 1, \frac{1}{\alpha} + \frac{1}{\beta} = 1, \\ (b-x)^n \|f^{(n+1)}\|_{[x,b],1} & \text{if } f^{(n+1)} \in L_1[x, b], \end{cases}$$

where (4.2) should be seen as all nine possible combinations.

*Proof.* Using the representation (2.1), we have

$$(4.3) \quad |f(x) - T_{n,\lambda}(x)| = \frac{1}{n!} \left| \lambda \int_a^x (x-t)^n f^{(n+1)}(t) dt + (1-\lambda) (-1)^{n+1} \int_x^b (t-x)^n f^{(n+1)}(t) dt \right| \\ \leq \left[ \lambda \int_a^x (x-t)^n |f^{(n+1)}(t)| dt + (1-\lambda) \int_x^b (t-x)^n |f^{(n+1)}(t)| dt \right].$$

It follows from Hölder's integral inequality, that

$$\int_a^x (x-t)^n |f^{(n+1)}(t)| dt \leq \begin{cases} \frac{(x-a)^{n+1}}{n+1} \|f^{(n+1)}\|_{[a,x],\infty} & \text{if } f^{(n+1)} \in L_\infty[a, x], \\ \frac{(x-a)^{n+\frac{1}{q}}}{(nq+1)^{\frac{1}{q}}} \|f^{(n+1)}\|_{[a,x],p} & \text{if } f^{(n+1)} \in L_p[a, x] \\ & \text{for } p > 1, \frac{1}{p} + \frac{1}{q} = 1, \\ (x-a)^n \|f^{(n+1)}\|_{[a,x],1} & \text{if } f^{(n+1)} \in L_1[a, x], \end{cases}$$

and

$$\int_x^b (t-x)^n |f^{(n+1)}(t)| dt \leq \begin{cases} \frac{(b-x)^{n+1}}{n+1} \|f^{(n+1)}\|_{[x,b],\infty} & \text{if } f^{(n+1)} \in L_\infty[x, b], \\ \frac{(b-x)^{n+\frac{1}{\beta}}}{(n\beta+1)^{\frac{1}{\beta}}} \|f^{(n+1)}\|_{[x,b],\alpha} & \text{if } f^{(n+1)} \in L_\alpha[x, b] \\ & \text{for } \alpha > 1, \frac{1}{\alpha} + \frac{1}{\beta} = 1, \\ (b-x)^n \|f^{(n+1)}\|_{[x,b],1} & \text{if } f^{(n+1)} \in L_1[x, b], \end{cases}$$

which together with (4.3) provide the desired result (4.2). ■

**Remark 6.** The result in (4.2) has some instances of interest which are perhaps more useful for applications. Namely, if  $f^{(n+1)} \in L_\infty[a, b]$ , then for each  $x \in [a, b]$

we have

$$\begin{aligned}
 (4.4) \quad |f(x) - T_{n,\lambda}(x)| & \leq \frac{1}{(n+1)!} \left[ \lambda (x-a)^n \|f^{(n+1)}\|_{[a,x],\infty} + (1-\lambda) (b-x)^n \|f^{(n+1)}\|_{[x,b],\infty} \right] \\
 & \leq \frac{1}{(n+1)!} [\lambda (x-a)^n + (1-\lambda) (b-x)^n] \|f^{(n+1)}\|_{[a,b],\infty} \\
 & =: N_1(\lambda, x).
 \end{aligned}$$

If we denote by

$$M_1(\lambda, x) := \lambda (x-a)^n + (1-\lambda) (b-x)^n,$$

then we also have the following upper bounds for  $M_1(\lambda, x)$ :

$$(4.5) \quad M_1(\lambda, x) \leq \begin{cases} \left[ \frac{1}{2} + \left| \lambda - \frac{1}{2} \right| \right] [(x-a)^n + (b-x)^n], \\ \left[ \lambda^s + (1-\lambda)^s \right]^{\frac{1}{s}} [(x-a)^{wn} + (b-x)^{wn}]^{\frac{1}{w}}, \\ \left[ \frac{1}{2} (b-a) + \left| x - \frac{a+b}{2} \right| \right]^{n-1}, \end{cases}$$

which yield three different bounds for  $N_1(\lambda, x)$ .

Now, in the case where  $f^{(n+1)} \in L_p[a, b]$  ( $p > 1$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ ), then we have

$$\begin{aligned}
 (4.6) \quad |f(x) - T_{n,\lambda}(x)| & \leq \frac{1}{n! (nq+1)^{\frac{1}{q}}} \left[ \lambda (x-a)^{n+\frac{1}{q}} \|f^{(n+1)}\|_{[a,x],p} \right. \\
 & \quad \left. + (1-\lambda) (b-x)^{n+\frac{1}{q}} \|f^{(n+1)}\|_{[x,b],p} \right] \\
 & \leq \frac{1}{n! (nq+1)^{\frac{1}{q}}} \left[ \lambda^q (x-a)^{nq+1} + (1-\lambda)^q (b-x)^{nq+1} \right]^{\frac{1}{q}} \|f^{(n+1)}\|_{[a,b],p} \\
 & =: N_p(\lambda, x), \quad x \in [a, b], \quad \lambda \in [0, 1].
 \end{aligned}$$

If we denote

$$M_q(\lambda, x) := \lambda^q (x-a)^{nq+1} + (1-\lambda)^q (b-x)^{nq+1}, \quad q > 1$$

then we also have the following upper bounds for  $M_q(\lambda, x)$

$$(4.7) \quad M_q(\lambda, x) \leq \begin{cases} \left[ \frac{1}{2} + \left| \lambda - \frac{1}{2} \right| \right]^q [(x-a)^{nq+1} + (b-x)^{nq+1}], \\ \left[ \lambda^{sq} + (1-\lambda)^{sq} \right]^{\frac{1}{s}} [(x-a)^{(nq+1)w} + (b-x)^{(nq+1)w}]^{\frac{1}{w}} \\ \text{for } s > 1, 1/s + 1/w = 1, \\ \left[ \lambda^q + (1-\lambda)^q \right] \left[ \frac{1}{2} (b-a) + \left| x - \frac{a+b}{2} \right| \right]^{nq+1}, \end{cases}$$

which provides three different bounds for  $N_p(\lambda, x)$ .

Finally, we have

$$\begin{aligned}
 (4.8) \quad & |f(x) - T_{n,\lambda}(x)| \\
 & \leq \frac{1}{n!} \left[ \lambda (x-a)^n \|f^{(n+1)}\|_{[a,x],1} + (1-\lambda) (b-x)^n \|f^{(n+1)}\|_{[x,b],1} \right] \\
 & \leq \frac{1}{n!} \max \{ \lambda (x-a)^n, (1-\lambda) (b-x)^n \} \|f^{(n+1)}\|_{[a,b],1} \\
 & \leq \frac{1}{n!} \left( \frac{1}{2} + \left| \lambda - \frac{1}{2} \right| \right) \left( \frac{1}{2} (b-a) + \left| x - \frac{a+b}{2} \right| \right)^n \|f^{(n+1)}\|_{[a,b],1}
 \end{aligned}$$

for any  $x \in [a, b]$  and  $\lambda \in [0, 1]$ .

**Remark 7.** If in the above inequalities we chose  $\lambda = (b-x)/(b-a)$  or  $\lambda = (x-a)/(b-a)$ , then we obtain various error bounds resulting from approximating the function  $f$  by the polynomials  $P_n$  and  $Q_n$  which are defined in the equations (3.15) and (3.16), respectively. The details are left to the interested reader.

## REFERENCES

- [1] A. Aleksandrov, Über die Äquivalenz des Perronschen und des Denjoyschen Integralbegriffes, *Math. Z.*, **20**(1924), 213–222.
- [2] J. C. Burkill, *A First Course in Mathematical Analysis*, Cambridge University Press, Cambridge, 1967.
- [3] A. Denjoy, Une extension de l'intégrale de M. Lebesgue, *C.R. Acad. Sci. Paris*, **154**(1912), 895–862.
- [4] H. Hake, Über de la Vallée Poussins Ober-und Unterfunktionen, *Math. Ann.* **83**(1921), 119–142.
- [5] R. Henstock, Definitions of Riemann type of variational integrals, *Proc. London Math. Soc.* **1**(1961), 402–418.
- [6] R. Henstock, A Riemann-type integral of Lebesgue power, *Canadian J. Math.* **20**(1968), 79–87.
- [7] J. Kurzweil, Generalised ordinary differential equations and continuous dependence on a parameter, *Czechoslovak Math. J.*, **82**(1957), 418–449.
- [8] J. Kurzweil, On Fubini theorem for general Perron integral, *Czechoslovak Math. J.*, **98**(1973), 286–297.
- [9] H. Looman, Über die Perronsche Integral Definition, *Math. Ann.* **93**(1935), 153–156.
- [10] N.N. Luzin, Sur les propriétés de l'intégrale de M. Denjoy, *C. R. Acad. Sci. Paris*, **155** (1912) 1475–1478.
- [11] J. Mäřík, Foundation of the theory of an integral in Euclidean space, (translated into English by L. I. Trudzik, Dept. of Math., Univ. of Melbourne, Parkville, Victoria 3052, Australia) *Časopis Pěst. Mat.* **77**(1952), 125–144.
- [12] J. Mawhin, *L'introduction à l'Analyse*, CABAY, Louvain-La-Neuve, 1984.
- [13] E.J. McShane, *A Riemann-type Integral that Includes Lebesgue-Stieltjes*, *Bochner and Stochastic Integrals*, Vol. **88**, Memoirs of Amer. Math. Soc., 1969.
- [14] R.M. McLeod, *The Generalized Riemann Integral*, Carus Mathematical Monographs, Mathematical Association of America, 1980.
- [15] O. Perron, Über den Integralbegriff, Sitzber, *Heidelberg Akad. Wiss. Abt. A* **16** (1914), 1–16.
- [16] M. Spivak, *Calculus*, W.A. Benjamin, New York, 1967.
- [17] K.R. Stromberg, *An Introduction to Classical Real Analysis*, Wadsworth, Belmont, California, 1981.
- [18] C. Swartz and B.S. Thomson, The Teaching of Mathematics; More on the Fundamental Theorem of Calculus, *Amer. Math. Monthly*, **95**(1988), 644–641.
- [19] H.B. Thompson, Taylor's theorem with the integral remainder under very weak differentiability assumptions, *Austral. Math. Soc. Gazette*, **12**(1985), 1–6.
- [20] R. Výborný, Kurzweil's integral and arclength, *Austral. Math. Soc. Gazette*, **8**(1981), 19–22.

- [21] H.B. Thompson, Taylor's Theorem Using the Generalised Riemann Integral, *Amer. Math. Monthly*, **96**(4) (1989), 346-350.

SCHOOL OF COMPUTER SCIENCE & MATHEMATICS, VICTORIA UNIVERSITY, PO Box 14428,  
MELBOURNE, VICTORIA, AUSTRALIA, 8001

*E-mail address:* Sever.Dragomir@vu.edu.au

*URL:* <http://rgmia.vu.edu.au/dragomir>

MATHEMATICS, SPS, THE UNIVERSITY OF QUEENSLAND, BRISBANE, QUEENSLAND, AUSTRALIA,  
4072

*E-mail address:* hbt@maths.uq.edu.au

*URL:* <http://www.maths.uq.edu.au/~hbt>