



VICTORIA UNIVERSITY
MELBOURNE AUSTRALIA

Inequalities involving the sequence $3\sqrt{a + 3\sqrt{a + \dots + 3\sqrt{a}}}$

This is the Published version of the following publication

Luo, Qiu-Ming, Qi, Feng, Barnett, Neil S and Dragomir, Sever S (2001)
Inequalities involving the sequence $3\sqrt{a + 3\sqrt{a + \dots + 3\sqrt{a}}}$. RGMIA
research report collection, 4 (4).

The publisher's official version can be found at

Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/17662/>

INEQUALITIES INVOLVING THE SEQUENCE $\sqrt[3]{a + \sqrt[3]{a + \cdots + \sqrt[3]{a}}}$

QIU-MING LUO, FENG QI, NEIL S. BARNETT, AND SEVER S. DRAGOMIR

ABSTRACT. In this article, the convergence of the sequence

$$\underbrace{\sqrt[3]{a + \sqrt[3]{a + \cdots + \sqrt[3]{a}}}}_n$$

is proved, and some inequalities involving this sequence are established for $a > 0$. As by-product, two identities involving irrational numbers are obtained. Two open problems are proposed.

1. INTRODUCTION

Let $a > 0$ and \mathbb{N} be the set of natural numbers. Denote

$$S_n(a) = \sqrt{\underbrace{a + \sqrt{a + \cdots + \sqrt{a}}}_n}, \quad (1)$$

$$f_n(a) = \frac{a - S_{n+1}(a)}{a - S_n(a)}. \quad (2)$$

In 1993, J.-Ch. Kuang sought the lower and upper bounds of $f_n(a)$, and conjectured that

$$f_n(a) > \frac{1}{a^2} \quad (3)$$

for all $n \in \mathbb{N}$. See [2, pp. 505–506 and p. 778].

2000 *Mathematics Subject Classification.* Primary 26D15; Secondary 40A05.

Key words and phrases. Inequality, sequence, ratio, convergence, identity, irrational number.

The first and second authors were supported in part by NSF (#10001016) of China, SF for the Prominent Youth of Henan Province, SF of Henan Innovation Talents at Universities, NSF of Henan Province (#004051800), SF for Pure Research of Natural Science of the Education Department of Henan Province (#1999110004), Doctor Fund of Jiaozuo Institute of Technology, China.

This paper was typeset using $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{L}\mathcal{A}\mathcal{T}\mathcal{E}\mathcal{X}$.

In 1999, as a reading note in [2], the second author raised the issue of the convergence of $S_{n,t}(a)$ and the bounds of $f_{n,t}(a)$, where, for $a > 0$ and $t \neq 0$,

$$S_{n,t}(a) = \sqrt[t]{\underbrace{a + \sqrt[t]{a} + \cdots + \sqrt[t]{a}}_n}, \quad (4)$$

$$f_{n,t}(a) = \frac{a - S_{n+1,t}(a)}{a - S_{n,t}(a)}. \quad (5)$$

Recently, the conjecture made by J.-Ch. Kuang was considered in [3], and the following result obtained.

Theorem A. *Let $a > 0$ and $n \in \mathbb{N}$.*

(1) *For $a \geq 2$, we have*

$$\frac{1}{a^2} < \frac{2(a + \sqrt{a} - a^2)}{(\sqrt{a} - a)(\sqrt{1 + 4a} + 2a + 1)} < f_n(a) < 1; \quad (6)$$

(2) *For $1 \leq a < 2$, there is a number $n_0 \in \mathbb{N}$ such that*

$$f_n(a) > 1 \geq \frac{1}{a^2} \quad (7)$$

holds for $n > n_0$;

(3) *For $0 < a < 1$, we have*

$$1 < f_n(a) \leq \frac{\sqrt{a + \sqrt{a}} - a}{\sqrt{a} - a}. \quad (8)$$

In this article, motivated by the reading note in [2] and the paper [3], we give an explicit solution to the problem involving the convergence of $S_{n,t}(a)$ and the bounds of $f_{n,t}(a)$ defined by (4) and (5) in the case of $t = 3$.

2. CONVERGENCE AND INEQUALITIES FOR $S_{n,t}(a)$

In this section, we first discuss the convergence of the sequence $S_{n,t}(a)$, and then obtain several inequalities for it.

Theorem 1. *Let $a > 0$ and $n \in \mathbb{N}$. The sequence $\{S_{n,3}(a)\}_{n=1}^{\infty}$ is strictly increasing.*

(1) *If $0 < a \leq \frac{2}{3\sqrt{3}}$, we have*

$$\lim_{n \rightarrow \infty} S_{n,3}(a) = \frac{2}{\sqrt{3}} \cos \left(\frac{1}{3} \arccos \frac{3a\sqrt{3}}{2} \right); \quad (9)$$

(2) If $a > \frac{2}{3\sqrt{3}}$, we have

$$\lim_{n \rightarrow \infty} S_{n,3}(a) = \sqrt[3]{\frac{a}{2} + \sqrt{\frac{a^2}{4} - \frac{1}{27}}} + \sqrt[3]{\frac{a}{2} - \sqrt{\frac{a^2}{4} - \frac{1}{27}}}. \quad (10)$$

Proof. By induction, it is easy to prove that the sequence $\{S_{n,3}(a)\}_{n=1}^{\infty}$ is strictly increasing for $a > 0$ and $\sqrt[3]{a} \leq S_{n,3}(a) < \sqrt[3]{a} + 1$, therefore, the sequence $\{S_{n,3}(a)\}_{n=1}^{\infty}$ converges.

Suppose $\lim_{n \rightarrow \infty} S_{n,3}(a) = x$, then, from $S_{n,3}^3(a) = a + S_{n-1,3}(a)$, it can be deduced that $x^3 - x - a = 0$.

From Cardano's formula [1] for the solution of a cubic equation of a single variable, the proof of Theorem 1 follows. \square

Using monotonicity of the sequence $\{S_{n,3}(a)\}_{n=1}^{\infty}$ and Theorem 1, the following inequalities are obtained.

Theorem 2. Let $a > 0$ and $n \in \mathbb{N}$.

(1) If $0 < a \leq \frac{2}{3\sqrt{3}}$, then

$$a < \sqrt[3]{a} \leq S_{n,3}(a) \leq \frac{2}{\sqrt{3}} \cos \left(\frac{1}{3} \arccos \frac{3a\sqrt{3}}{2} \right); \quad (11)$$

(2) If $\frac{2}{3\sqrt{3}} < a < 1$, we have

$$a < \sqrt[3]{a} \leq S_{n,3}(a) < \sqrt[3]{\frac{a}{2} + \sqrt{\frac{a^2}{4} - \frac{1}{27}}} + \sqrt[3]{\frac{a}{2} - \sqrt{\frac{a^2}{4} - \frac{1}{27}}}; \quad (12)$$

(3) If $1 \leq a < \sqrt{2}$, there exists a number $n_0 \in \mathbb{N}$ such that

$$\sqrt[3]{a} \leq S_{n_0,3}(a) \leq a < S_{n,3}(a) < \sqrt[3]{\frac{a}{2} + \sqrt{\frac{a^2}{4} - \frac{1}{27}}} + \sqrt[3]{\frac{a}{2} - \sqrt{\frac{a^2}{4} - \frac{1}{27}}} \quad (13)$$

holds for $n > n_0$;

(4) If $a \geq \sqrt{2}$, then

$$\sqrt[3]{a} \leq S_{n,3}(a) < \sqrt[3]{\frac{a}{2} + \sqrt{\frac{a^2}{4} - \frac{1}{27}}} + \sqrt[3]{\frac{a}{2} - \sqrt{\frac{a^2}{4} - \frac{1}{27}}} \leq a. \quad (14)$$

Proof. We verify the inequalities (13) and (14), the rest follow similarly.

For $x \geq \frac{2}{3\sqrt{3}}$, we introduce a function $\psi(x)$ defined by

$$\psi(x) \triangleq g(x) - x \triangleq \sqrt[3]{\frac{x}{2} + \sqrt{\frac{x^2}{4} - \frac{1}{27}}} + \sqrt[3]{\frac{x}{2} - \sqrt{\frac{x^2}{4} - \frac{1}{27}}} - x. \quad (15)$$

We also claim that $\psi(x) \leq 0$ if and only if $x \geq \sqrt{2}$.

Direct calculation reveals that

$$g^3(x) = g(x) + x. \quad (16)$$

We have, then

$$g'(x) = \frac{1}{3g^2(x) - 1}, \quad g''(x) = -\frac{6g(x)}{[3g^2(x) - 1]^3}. \quad (17)$$

It is clear that both the terms of $g(x)$ are positive for $x \geq \frac{2}{3\sqrt{3}}$. Using the arithmetic-geometric mean inequality yields that $g(x) > \frac{2\sqrt{3}}{3}$ for $x \geq \frac{2}{3\sqrt{3}}$. This leads to $3g^2(x) - 1 > 3$ for $x \geq \frac{2}{3\sqrt{3}}$. Therefore, the first derivative of $g(x)$ satisfies $g'(x) > 0$ and the second derivative $g''(x) < 0$ for $x \geq \frac{2}{3\sqrt{3}}$. This means that the function $g(x)$ is increasing and concave on $[\frac{2}{3\sqrt{3}}, \infty)$.

Straightforward computation yields

$$\psi\left(\frac{2}{3\sqrt{3}}\right) = \frac{4}{3\sqrt{3}}, \quad \lim_{x \rightarrow \infty} \psi(x) = -\infty. \quad (18)$$

This implies that the curve $y = g(x)$ and the straight line $y = x$ intersect at a unique point on $[\frac{2}{3\sqrt{3}}, \infty)$. Thus, there exists a unique point $x_0 \in (\frac{2}{3\sqrt{3}}, \infty)$ such that $\psi(x) > 0$ for $x \in (\frac{2}{3\sqrt{3}}, x_0)$ and $\psi(x) < 0$ for (x_0, ∞) .

Since $\psi(\sqrt{2}) = 0$, consequently $x_0 = \sqrt{2}$. The proof is complete. \square

Remark 1. Now we provide another proof for the claim that $\psi(x) \leq 0$ if and only if $x \geq \sqrt{2}$.

Firstly, we prove that $g(x) = x$ holds if and only if $x = \sqrt{2}$. Letting $x = \sqrt{2}$ in (16), we have $g^3(\sqrt{2}) - g(\sqrt{2}) - \sqrt{2} = 0$, which is equivalent to $[g(\sqrt{2}) - \sqrt{2}][g^2(\sqrt{2}) + \sqrt{2}g(\sqrt{2}) + 1] = 0$, thus $g(\sqrt{2}) = \sqrt{2}$. Conversely, letting $g(x) = x \geq \frac{2}{3\sqrt{3}}$, then equation (16) reduces to $x^3 - 2x = 0$, and so $x = \sqrt{2}$.

Secondly, we verify that $g(x) < x$ is valid if and only if $x > \sqrt{2}$. If $g(x) < x$, then equation (16) can be rewritten as $x - g(x) = g^3(x) - 2g(x) = g(x)[g^2(x) - 2] > 0$, then $x > g(x) > \sqrt{2}$. Conversely, if $x > \sqrt{2}$, then $g^3(x) - g(x) - \sqrt{2} > g^3(x) - g(x) - x = 0$, which is equivalent to $[g(x) - \sqrt{2}][g^2(x) + \sqrt{2}g(x) + 1] > 0$, and so $g(x) > \sqrt{2}$. Therefore, $g(x) - x = 2g(x) - g^3(x) = g(x)[2 - g^2(x)] < 0$, which means that $g(x) < x$.

The proof is complete.

Corollary 1. *The irrational number $\sqrt{2}$ can be expressed as*

$$\sqrt{2} = \sqrt[3]{\frac{1}{\sqrt{2}} - \frac{5}{3\sqrt{6}}} + \sqrt[3]{\frac{1}{\sqrt{2}} + \frac{5}{3\sqrt{6}}}, \quad (19)$$

which is equivalent to

$$\sqrt[3]{3\sqrt{3} - 5} + \sqrt[3]{3\sqrt{3} + 5} = \sqrt{3} \cdot \sqrt[3]{4}. \quad (20)$$

Proof. Identity (20) follows from simplifying (19) directly.

Raising both sides of $A = \sqrt[3]{3\sqrt{3} - 5} + \sqrt[3]{3\sqrt{3} + 5}$ to the power of 3 shows that A satisfies the cubic equation $x^3 - 3\sqrt[3]{2}x - 6\sqrt{3} = 0$. By Cardano's formula in [1], it follows that $A = \sqrt{3} \cdot \sqrt[3]{4}$. The proof is complete. \square

3. INEQUALITIES FOR THE SEQUENCE $f_{n,3}(a)$

From the monotonicity and inequalities for the sequence $\{S_{n,3}(a)\}_{n=1}^{\infty}$, we will derive some inequalities for the sequence $\{f_{n,3}(a)\}_{n=1}^{\infty}$.

Theorem 3. *Let $a > 0$ and $n \in \mathbb{N}$.*

(1) *When $0 < a < 1$, we have*

$$1 < f_{n,3}(a) \leq \frac{\sqrt[3]{a + \sqrt[3]{a}} - a}{\sqrt[3]{a} - a}; \quad (21)$$

(2) *When $1 \leq a < \sqrt{2}$, there exists a number $n_0 \in \mathbb{N}$ such that*

$$f_{n,3}(a) > 1 > \frac{1}{a} > \frac{1}{a^2} \quad (22)$$

holds for all $n > n_0$;

(3) *When $a \geq \sqrt{2}$, we have*

$$1 > f_{n,3}(a) > \frac{1}{a^2 + a\alpha + \alpha^2} \left(1 + \frac{a^3 - 2a}{a - \sqrt[3]{a}} \right), \quad (23)$$

where

$$\alpha = \sqrt[3]{\frac{a}{2} + \sqrt{\frac{a^2}{4} - \frac{1}{27}}} + \sqrt[3]{\frac{a}{2} - \sqrt{\frac{a^2}{4} - \frac{1}{27}}}. \quad (24)$$

Proof. For $0 < a < 1$, since the sequence $\{S_{n,3}(a)\}_{n=1}^{\infty}$ is strictly increasing with $S_{n,3}(a) > a$, then $a - S_{n+1,3}(a) < a - S_{n,3}(a) < 0$, and $f_{n,3}(a) = \frac{a - S_{n+1,3}(a)}{a - S_{n,3}(a)} > 1$.

By standard argument, it follows that

$$\begin{aligned}
f_{n+1,3}(a) &= \frac{a - S_{n+2,3}(a)}{a - S_{n+1,3}(a)} \\
&= \frac{1}{a^2 + aS_{n+2,3}(a) + S_{n+2,3}^2(a)} \frac{a^3 - S_{n+2,3}^3(a)}{a - S_{n+1,3}(a)} \\
&= \frac{1}{a^2 + aS_{n+2,3}(a) + S_{n+2,3}^2(a)} \left[1 + \frac{2a - a^3}{S_{n+1,3}(a) - a} \right] \\
&< \frac{1}{a^2 + aS_{n+1,3}(a) + S_{n+1,3}^2(a)} \left[1 + \frac{2a - a^3}{S_{n,3}(a) - a} \right] \\
&= \frac{a - S_{n+1,3}(a)}{a - S_{n,3}(a)} \\
&= f_{n,3}(a).
\end{aligned} \tag{25}$$

This implies that the sequence $\{f_{n,3}(a)\}_{n=1}^{\infty}$ is strictly decreasing, therefore

$$f_{n,3}(a) \leq f_{1,3}(a) = \frac{\sqrt[3]{a + \sqrt[3]{a}} - a}{\sqrt[3]{a} - a}. \tag{26}$$

For $1 \leq a < \sqrt{2}$, and (13), there exists a number $n_0 \in \mathbb{N}$ such that $a - S_{n+1,3}(a) < a - S_{n,3}(a) < 0$ holds for $n > n_0$. Hence

$$f_{n,3}(a) = \frac{a - S_{n+1,3}(a)}{a - S_{n,3}(a)} > 1 > \frac{1}{a} > \frac{1}{a^2}, \quad n > n_0.$$

For $n > n_0$, the formula (25) is also valid. Thus, the sequence $\{f_{n,3}(a)\}_{n=n_0+1}^{\infty}$ is strictly decreasing, and

$$\frac{1}{a^2} < \frac{1}{a} < 1 < f_{n,3}(a) < \frac{a - S_{n_0+2,3}(a)}{a - S_{n_0+1,3}(a)}, \quad n > n_0. \tag{27}$$

For $a \geq \sqrt{2}$, and (14), we have $0 < a - S_{n+1,3}(a) < a - S_{n,3}(a)$ for $n \in \mathbb{N}$. Then $f_{n,3}(a) = \frac{a - S_{n+1,3}(a)}{a - S_{n,3}(a)} < 1$. From a combination of the following formula (28),

$$f_{n,3}(a) = \frac{a - S_{n+1,3}(a)}{a - S_{n,3}(a)} = \frac{1}{a^2 + aS_{n+1,3}(a) + S_{n+1,3}^2(a)} \left[1 + \frac{2a - a^3}{S_{n,3}(a) - a} \right], \tag{28}$$

and the inequalities in (14), then (23) follows.

The proof is complete. \square

4. OPEN PROBLEMS

It is natural to pose the following questions.

- (1) Can we prove or disprove the convergence of the sequence $\{S_{n,t}(a)\}_{n=1}^{\infty}$ for a positive real number a and nonzero real number $t \neq 0$?

- (2) Can we establish sharp lower and upper sharp bounds for the sequence $\{f_{n,t}(a)\}_{n=1}^{\infty}$ for a positive real number a and nonzero real number $t \neq 0$?

Acknowledgements. This paper was finalized during the second author visited RGMIA between November 1, 2001 and January 31, 2002, as a Visiting Professor with grants from the Victoria University of Technology and Jiaozuo Institute of Technology.

REFERENCES

- [1] C. M. Bender and S. A. Orszag, *Advanced Mathematical Methods for Scientists and Engineers*, McGraw-Hill, 1978. Chinese edition, translated by J.-Ch. Li, F.-Q. Zhuang, and B.-Y. Wang, Science Press, 1992, p. 280.
- [2] J.-Ch. Kuang, *Chángyòng Bùděngshì (Applied Inequalities)*, 2nd edition, Hunan Education Press, Changsha, China, 1993. (Chinese)
- [3] B.-Y. Xi, *Discussion on a conjecture about an inequality*, Shùxué Tōngbào (Bulletin of Mathematics) (2000), no. 11, 21–22. (Chinese)

(Luo) DEPARTMENT OF BROADCAST-TELEVISION-TEACHING, JIAOZUO UNIVERSITY, JIAOZUO CITY, HENAN 454002, CHINA

E-mail address: luoqm236@sohu.com

(Qi) DEPARTMENT OF MATHEMATICS, JIAOZUO INSTITUTE OF TECHNOLOGY, JIAOZUO CITY, HENAN 454000, CHINA

E-mail address: qifeng@jz.it.edu.cn or qifeng618@hotmail.com

URL: <http://rgmia.vu.edu.au/qi.html>

(Barnett) SCHOOL OF COMMUNICATIONS AND INFORMATICS, VICTORIA UNIVERSITY OF TECHNOLOGY, P. O. BOX 14428, MELBOURNE CITY MC, VICTORIA 8001, AUSTRALIA

E-mail address: neil@matilda.vu.edu.au

URL: <http://sci.vu.edu.au/staff/neilb.html>

(Dragomir) SCHOOL OF COMMUNICATIONS AND INFORMATICS, VICTORIA UNIVERSITY OF TECHNOLOGY, P. O. BOX 14428, MELBOURNE CITY MC, VICTORIA 8001, AUSTRALIA

E-mail address: sever.dragomir@vu.edu.au

URL: <http://rgmia.vu.edu.au/SSDragomirWeb.html>