



VICTORIA UNIVERSITY
MELBOURNE AUSTRALIA

Lower and upper bounds of $\zeta(3)$

This is the Published version of the following publication

Luo, Qiu-Ming, Wei, Zong-Li and Qi, Feng (2001) Lower and upper bounds of $\zeta(3)$. RGMIA research report collection, 4 (4).

The publisher's official version can be found at

Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/17668/>

LOWER AND UPPER BOUNDS OF $\zeta(3)$

QIU-MING LUO, ZONG-LI WEI, AND FENG QI

ABSTRACT. In this short note, using refinements of Jordan's inequality and an integral expression of $\zeta(3)$, the lower and upper bounds of $\zeta(3)$ are obtained, and some related results are improved.

1. INTRODUCTION

The Riemann zeta function can be defined by the integral

$$\zeta(x) = \frac{1}{\Gamma(x)} \int_0^\infty \frac{u^{x-1}}{e^u - 1} du, \quad (1)$$

where $x > 1$. If x is an integer n , we obtain the most common form of the function $\zeta(n)$, which is given by

$$\zeta(n) = \sum_{k=1}^{\infty} \frac{1}{k^n}. \quad (2)$$

For $n = 1$, the zeta function reduces to the harmonic series, which is divergent.

The Riemann zeta function can also be defined in terms of multiple integrals by

$$\zeta(n) = \underbrace{\int_0^1 \cdots \int_0^1}_n \frac{\prod_{i=1}^n dx_i}{1 - \prod_{i=1}^n x_i}. \quad (3)$$

An additional identity is

$$\lim_{s \rightarrow 1} \zeta(s) - \frac{1}{s-1} = \gamma, \quad (4)$$

where γ is the Euler-Mascheroni constant.

2000 *Mathematics Subject Classification.* 11R40, 42A16.

Key words and phrases. Lower and upper bounds, Riemann zeta function, Jordan inequality.

The third author was supported in part by NSF (#10001016) of China, SF for the Prominent Youth of Henan Province, SF of Henan Innovation Talents at Universities, NSF of Henan Province (#004051800), SF for Pure Research of Natural Science of the Education Department of Henan Province (#1999110004), Doctor Fund of Jiaozuo Institute of Technology, China.

This paper was typeset using $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{L}\mathcal{A}\mathcal{T}\mathcal{E}\mathcal{X}$.

The Euler product formula can also be written as

$$\zeta(x) = \left[\prod_{i=2}^{\infty} (1 - i^{-x}) \right]^{-1}. \quad (5)$$

The function $\zeta(n)$ was proved to be transcendental for all even n . For $n = 2k$ we have:

$$\zeta(n) = \frac{2^{n-1} |B_n| \pi^n}{n!}, \quad (6)$$

where B_n is the Bernoulli number. Another intimate connection with the Bernoulli numbers is provided by

$$B_n = (-1)^{n+1} n \zeta(1-n). \quad (7)$$

The Riemann zeta function is related to the gamma function and it has important applications in mathematics, especially in the Analytic Number Theory.

The Riemann zeta function may be computed analytically for even n using either Contour integration or Parseval's theorem with the appropriate Fourier series.

No analytic form for $\zeta(n)$ is known for odd $n = 2k + 1$, but $\zeta(2k + 1)$ can be expressed as the sum limit

$$\zeta(2k + 1) = \left(\frac{\pi}{2}\right)^{2k+1} \lim_{t \rightarrow \infty} \frac{1}{t^{2k+1}} \sum_{i=1}^{\infty} \left[\cot\left(\frac{i}{2t+1}\right) \right]^{2k+1}. \quad (8)$$

The values for the first few integral arguments are

$$\begin{aligned} \zeta(0) &= -\frac{1}{2}, & \zeta(1) &= \infty, & \zeta(2) &= \frac{\pi^2}{6}, \\ \zeta(3) &= 1.2020569032 \dots, & \zeta(4) &= \frac{\pi^4}{90}, & \zeta(5) &= 1.0369277551 \dots \end{aligned}$$

In [9, p. 81], using the Jordan inequality $1 < \frac{x}{\sin x} \leq \frac{\pi}{2}$ for $x \in (0, \frac{\pi}{2}]$ and two integral expressions

$$\sum_{i=0}^{\infty} \frac{1}{(2i+1)^3} = \frac{1}{4} \int_0^{\pi/2} \frac{x(\pi-x)}{\sin x} dx, \quad (9)$$

$$\sum_{i=1}^{\infty} \frac{1}{i^3} = \frac{1}{6\pi} \int_0^{\pi} \left(\frac{x(\pi-x)}{\sin x} \right)^2 dx, \quad (10)$$

the following estimates were given

$$\frac{3\pi^2}{32} \leq \sum_{i=0}^{\infty} \frac{1}{(2i+1)^3} \leq \frac{3\pi^3}{64}, \quad \sum_{i=1}^{\infty} \frac{1}{i^3} < \frac{29}{24}. \quad (11)$$

There is much literature devoted to evaluations and proofs of $\zeta(2) = \frac{\pi^2}{6}$. Please refer to the related references in this paper.

In this short note, using an integral expression of $\zeta(3)$ and refinements of the Jordan inequality, we obtain the following

Theorem 1. *The value of $\zeta(3)$ can be evaluated by the following*

$$\begin{aligned}
 1.201 \dots &= \frac{1}{14\sqrt{5}} \left\{ \pi\sqrt{30(\sqrt{30}-5)} \ln \left[\pi^2 + 4\pi\sqrt{5+\sqrt{30}} + 8\sqrt{30} \right] \right. \\
 &\quad - \pi\sqrt{30(\sqrt{30}-5)} \ln \left[\pi^2 - 4\pi\sqrt{5+\sqrt{30}} + 8\sqrt{30} \right] \\
 &\quad + 2\pi\sqrt{30(5+\sqrt{30})} \arctan \left[\frac{\pi + 2\sqrt{\sqrt{30}+5}}{2\sqrt{\sqrt{30}-5}} \right] \\
 &\quad + 2\pi\sqrt{30(5+\sqrt{30})} \arctan \left[\frac{\pi - 2\sqrt{\sqrt{30}+5}}{2\sqrt{\sqrt{30}-5}} \right] \\
 &\quad + 120 \arctan \left[\frac{\pi + 2\sqrt{\sqrt{30}+5}}{2\sqrt{\sqrt{30}-5}} \right] \\
 &\quad \left. - 120 \arctan \left[\frac{\pi - 2\sqrt{\sqrt{30}+5}}{2\sqrt{\sqrt{30}-5}} \right] - 240 \arctan \sqrt{\frac{\sqrt{30}+5}{\sqrt{30}-5}} \right\} \\
 &< \zeta(3) = \sum_{i=0}^{\infty} \frac{1}{i^3} = \frac{8}{7} \sum_{i=0}^{\infty} \frac{1}{(2i+1)^3} \\
 &< \frac{2}{7} \left\{ 3 \ln(24 - \pi^2) - 9 \ln 2 - 3 \ln 3 + \pi\sqrt{6} \operatorname{arctanh} \frac{\pi\sqrt{6}}{12} \right\} \\
 &= 7 \left\{ 6 \ln \frac{24 - \pi^2}{4} - 6 \ln 6 - \pi\sqrt{6} \ln \frac{12 - \pi\sqrt{6}}{2} + \pi\sqrt{6} \ln \frac{12 + \pi\sqrt{6}}{2} \right\} \\
 &= 1.217 \dots .
 \end{aligned} \tag{12}$$

It is obvious that inequality (12) improves inequality (11).

2. LOWER AND UPPER BOUNDS OF $\zeta(3)$

It is well-known that, for $x \in [0, \frac{\pi}{2}]$, we have

$$x - \frac{1}{6}x^3 \leq \sin x \leq x - \frac{1}{6}x^3 + \frac{1}{120}x^5. \tag{13}$$

The inequalities in (13) can be found in [7, 10, 11] and [9, p. 309]. The inequalities in (13) are a refinement of the well-known Jordan inequality $\frac{2}{\pi}x \leq \sin x \leq x$ for $x \in [0, \frac{\pi}{2}]$.

It is easy to see that

$$\sum_{i=0}^{\infty} \frac{1}{(2i+1)^3} = \frac{7}{8}\zeta(3). \tag{14}$$

From formula (9) and inequality (13), it follows by direct calculation that

$$\begin{aligned}
1.051 \dots &= \frac{1}{16\sqrt{5}} \left\{ \pi \sqrt{30(\sqrt{30}-5)} \ln \left[\pi^2 + 4\pi \sqrt{5 + \sqrt{30}} + 8\sqrt{30} \right] \right. \\
&\quad - \pi \sqrt{30(\sqrt{30}-5)} \ln \left[\pi^2 - 4\pi \sqrt{5 + \sqrt{30}} + 8\sqrt{30} \right] \\
&\quad + 2\pi \sqrt{30(5 + \sqrt{30})} \arctan \left[\frac{\pi + 2\sqrt{\sqrt{30} + 5}}{2\sqrt{\sqrt{30} - 5}} \right] \\
&\quad + 2\pi \sqrt{30(5 + \sqrt{30})} \arctan \left[\frac{\pi - 2\sqrt{\sqrt{30} + 5}}{2\sqrt{\sqrt{30} - 5}} \right] \\
&\quad + 120 \arctan \left[\frac{\pi + 2\sqrt{\sqrt{30} + 5}}{2\sqrt{\sqrt{30} - 5}} \right] \\
&\quad \left. - 120 \arctan \left[\frac{\pi - 2\sqrt{\sqrt{30} + 5}}{2\sqrt{\sqrt{30} - 5}} \right] - 240 \arctan \sqrt{\frac{\sqrt{30} + 5}{\sqrt{30} - 5}} \right\} \quad (15) \\
&= \frac{1}{4} \int_0^{\pi/2} \frac{\pi - x}{1 - \frac{1}{6}x^2 + \frac{1}{120}x^4} dx \\
&< \sum_{i=0}^{\infty} \frac{1}{(2i+1)^3} \\
&< \frac{1}{4} \int_0^{\pi/2} \frac{\pi - x}{1 - \frac{1}{6}x^2} dx \\
&= \frac{3 \ln(24 - \pi^2) - 9 \ln 2 - 3 \ln 3 + \pi\sqrt{6} \operatorname{arctanh} \frac{\pi\sqrt{6}}{12}}{4} \\
&= \frac{6 \ln \frac{24 - \pi^2}{4} - 6 \ln 6 - \pi\sqrt{6} \ln \frac{12 - \pi\sqrt{6}}{2} + \pi\sqrt{6} \ln \frac{12 + \pi\sqrt{6}}{2}}{8} \\
&= 1.0654 \dots
\end{aligned}$$

The proof of Theorem 1 follows from a combination of (14) with (15).

REFERENCES

- [1] T. M. Apostol, *A proof that Euler missed: Evaluating $\zeta(2)$ the easy way*, Math. Intel. **5** (1983), 59–60.
- [2] R. Ayoub, *Euler and the zeta function*, Amer. Math. Monthly **81** (1974), 1067–1086.
- [3] D. Borwein and J. Borwein, *On an intriguing integral and some series related to $\zeta(4)$* , Proc. Amer. Math. Soc. **123** (1995), 1191–1198.
- [4] B. R. Choe, *An elementary proof of $\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$* , Amer. Math. Monthly **94** (1987), 662–663.

- [5] Group of compilation, *Shùxué Shǒucè (Handbook of Mathematics)*, The People's Education Press, Beijing, China, 1979. (Chinese)
- [6] G. Klambauer, *Mathematical Analysis*, Chinese edition, translated by Ben-Wang Sun, The People's Press of Hunan, Changsha City, Hunan, China, 1981. English edition, Marcel Dekker, Inc., New York, 1975.
- [7] G. Klambauer, *Problems and Propositions in Analysis*, Marcel Dekker, Inc., New York and Basel, 1979, pp. 171–173, pp. 266–267, pp. 280–284.
- [8] R. A. Kortram, *Simple proofs for $\sum_{k=1}^{\infty} \frac{1}{k^2} = \frac{\pi^2}{6}$ and $\sin x = x \prod_{k=1}^{\infty} \left(1 - \frac{x^2}{(k\pi)^2}\right)$* , *Math. Mag.* **69** (1996), 122–125.
- [9] J.-Ch. Kuang, *Chángyòng Bùděngshì (Applied Inequalities)*, 2nd edition, Hunan Education Press, Changsha City, Hunan Province, China, 1993. (Chinese)
- [10] F. Qi, *Extensions and sharpenings of Jordan's and Kober's inequality*, *Gōngkē Shùxué (Journal of Mathematics for Technology)* **12** (1996), no. 4, 98–102. (Chinese)
- [11] F. Qi and Q.-D. Hao, *Refinements and sharpenings of Jordan's and Kober's inequality*, *Mathematics and Informatics Quarterly* **8** (1998), no. 3, 116–120.
- [12] M. E. Taylor, *Partial Differential Equations I*, *Applied Mathematical Sciences* **15**, Springer-Verlag, 1996. Reprinted in China by Beijing World Publishing Corporation, 1999, pp. 237–239.
- [13] E. W. Weisstein, *CRC Concise Encyclopedia of Mathematics on CD-ROM*, 1999. Available online at <http://www.math.ustc.edu.cn/Encyclopedia/contents/RiemannZetaFunction.html>.

(Luo) DEPARTMENT OF BROADCAST-TELEVISION-TEACHING, JIAOZUO UNIVERSITY, JIAOZUO CITY, HENAN 454002, CHINA

(Wei) DEPARTMENT OF MATHEMATICS, LUOYANG NORMAL COLLEGE, LUOYANG CITY, HENAN 471022, CHINA

E-mail address: weizl@lync.edu.cn

(Qi) DEPARTMENT OF MATHEMATICS, JIAOZUO INSTITUTE OF TECHNOLOGY, JIAOZUO CITY, HENAN 454000, CHINA

E-mail address: qifeng@jzit.edu.cn

URL: <http://rgmia.vu.edu.au/qi.html>