



**VICTORIA UNIVERSITY**  
MELBOURNE AUSTRALIA

## *Generalizations of Bernoulli's Numbers and Polynomials*

This is the Published version of the following publication

Luo, Qiu-Ming, Guo, Bai-Ni and Qi, Feng (2002) Generalizations of Bernoulli's Numbers and Polynomials. RGMIA research report collection, 5 (2).

The publisher's official version can be found at

Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/17699/>

# GENERALIZATIONS OF BERNOULLI'S NUMBERS AND POLYNOMIALS

QIU-MING LUO, BAI-NI GUO, AND FENG QI

ABSTRACT. In this paper, the concepts of Bernoulli numbers  $B_n$ , Bernoulli polynomials  $B_n(x)$ , the generalized Bernoulli numbers  $B_n(a, b)$  are further generalized to one which is called as the generalized Bernoulli polynomials, and many basic properties and some relationships between  $B_n$ ,  $B_n(x)$  and  $B_n(a, b)$  are established.

## 1. INTRODUCTION

It is well known that Bernoulli's numbers and polynomials play important roles in mathematics, they are main objects in the Theory of Special Functions [4]. Their definitions can be given as follows.

**Definition 1.** The numbers  $B_n$ ,  $0 \leq n \leq \infty$ , are called Bernoulli numbers if

$$\phi(t) = \frac{t}{e^t - 1} = \sum_{n=0}^{\infty} \frac{B_n}{n!} t^n, \quad |t| < 2\pi. \quad (1)$$

**Definition 2.** The functions  $B_n(x)$ ,  $0 \leq n \leq \infty$ , are called Bernoulli polynomials if they satisfy

$$\phi(x; t) = \frac{te^{xt}}{e^t - 1} = \sum_{n=0}^{\infty} \frac{B_n(x)}{n!} t^n, \quad |t| < 2\pi, \quad x \in \mathbb{R}. \quad (2)$$

The usual definition of Bernoulli polynomials is

$$\frac{t^\sigma e^{ut}}{(e^t - 1)^\sigma} = \sum_{n=0}^{\infty} \frac{B_n^\sigma(u)}{n!} t^n, \quad |t| < 2\pi. \quad (3)$$

---

2000 *Mathematics Subject Classification.* Primary 11B68, 33E20; Secondary 26A48, 40A30.

*Key words and phrases.* Bernoulli numbers, Bernoulli polynomials, generalized Bernoulli numbers, generalized Bernoulli polynomials.

The authors were supported in part by NNSF (#10001016) of China, SF for the Prominent Youth of Henan Province, SF of Henan Innovation Talents at Universities, NSF of Henan Province (#004051800), SF for Pure Research of Natural Science of the Education Department of Henan Province (#1999110004), Doctor Fund of Jiaozuo Institute of Technology, China.

This paper was typeset using  $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{L}\mathcal{A}\mathcal{T}\mathcal{E}\mathcal{X}$ .

In [2], the last two authors generalized the concept of Bernoulli numbers as follows.

**Definition 3.** Let  $a, b > 0$  and  $a \neq b$ . The generalized Bernoulli numbers  $B_n(a, b)$  are defined by

$$\phi(t; a, b) = \frac{t}{b^t - a^t} = \sum_{n=0}^{\infty} \frac{B_n(a, b)}{n!} t^n, \quad |t| < \frac{2\pi}{|\ln b - \ln a|}. \quad (4)$$

Among other things, some basic properties and relationships between  $B_n$ ,  $B_n(x)$  and  $B_n(a, b)$  were also studied in [2] initially and originally.

In this article, we first give the following definition of the generalized Bernoulli polynomials, which generalizes the concepts stated above, and then research their basic properties and relationships to Bernoulli numbers  $B_n$ , Bernoulli polynomials  $B_n(x)$  and the generalized Bernoulli numbers  $B_n(a, b)$ .

## 2. DEFINITIONS AND PROPERTIES OF GENERALIZED BERNOULLI POLYNOMIALS

It is easy to see that the following definition is natural and essential generalizations of the concepts of Bernoulli numbers  $B_n$ , Bernoulli polynomials  $B_n(x)$  and the generalized Bernoulli numbers  $B_n(a, b)$ .

**Definition 4.** Let  $a, b, c > 0$  and  $a \neq b$ . The generalized Bernoulli polynomials  $B_n(x; a, b, c)$  for nonnegative integer  $n$  are defined by

$$\phi(x; t; a, b, c) = \frac{tc^{xt}}{b^t - a^t} = \sum_{n=0}^{\infty} \frac{B_n(x; a, b, c)}{n!} t^n, \quad |t| < \frac{2\pi}{|\ln b - \ln a|}, \quad x \in \mathbb{R}. \quad (5)$$

The generalized Bernoulli polynomials  $B_n(x; a, b, c)$  have the following properties, which are stated as theorems below.

**Theorem 1.** Let  $a, b, c > 0$  and  $a \neq b$ . For  $x \in \mathbb{R}$  and  $n \geq 0$ , we have

$$B_n(x; 1, e, e) = B_n(x), \quad B_n(0; a, b, c) = B_n(a, b), \quad B_n(0; 1, e, e) = B_n, \quad (6)$$

$$B_n(x; a, b, 1) = B_n(a, b), \quad B_n(x; 1, e, 1) = B_n,$$

$$B_n(x; a, b, c) = \sum_{k=0}^n \binom{n}{k} [\ln c]^{n-k} B_k(a, b) x^{x-k}, \quad (7)$$

$$B_n(x; a, b, c) = \sum_{k=0}^n \binom{n}{k} [\ln c]^{n-k} [\ln b - \ln a]^{k-1} B_k \left( \frac{\ln a}{\ln a - \ln b} \right) x^{x-k}, \quad (8)$$

$$B_n(x; a, b, c) = \sum_{k=0}^n \sum_{j=0}^k (-1)^{k-j} \binom{n}{k} \binom{k}{j} [\ln c]^{n-k} [\ln a]^{k-j} \left[ \ln \frac{b}{a} \right]^{j-1} B_j x^{x-k}. \quad (9)$$

*Proof.* Applying Definition 3 to the term  $\frac{t}{b^t - a^t}$  and expanding the exponential function  $c^{xt}$  at  $t = 0$  yields

$$\begin{aligned} \frac{t c^{xt}}{b^t - a^t} &= \left( \sum_{k=0}^{\infty} \frac{B_k(a, b)}{k!} t^k \right) \left( \sum_{i=0}^{\infty} \frac{x^i (\ln c)^i}{i!} t^i \right) \\ &= \sum_{k=0}^{\infty} \sum_{i=0}^k \frac{(\ln c)^{k-i}}{i!(k-i)!} B_i(a, b) x^{k-i} t^k \\ &= \sum_{n=0}^{\infty} \left( \sum_{k=0}^n \binom{n}{k} (\ln c)^{n-k} B_k(a, b) x^{n-k} \right) \frac{t^n}{n!}. \end{aligned} \quad (10)$$

Combining (10) and (5) and equating their coefficients of  $t^n$  produces formula (7).

The following two formulae were provided in [2]:

$$B_n(a, b) = (\ln b - \ln a)^{n-1} B_n \left( \frac{\ln a}{\ln a - \ln b} \right), \quad (11)$$

$$B_n(a, b) = \sum_{i=0}^n (-1)^{n-i} (\ln b - \ln a)^{i-1} (\ln a)^{n-i} \binom{n}{i} B_i. \quad (12)$$

Substituting (11) and (12) into (7) leads to (8) and (9).

Formulae in (6) are obvious.  $\square$

Now we give some results about derivatives and integrals of the generalized Bernoulli polynomials  $B_n(x; a, b, c)$  as follows.

**Theorem 2.** *Let  $a, b, c > 0$ ,  $a \neq b$ ,  $n \geq 0$ , and  $x \in \mathbb{R}$ . For any nonnegative integer  $\ell$  and real numbers  $\alpha$  and  $\beta$ , we have*

$$\frac{\partial^\ell B_n(x; a, b, c)}{\partial x^\ell} = \frac{n!}{(n-\ell)!} (\ln c)^\ell B_{n-\ell}(x; a, b, c), \quad (13)$$

$$\int_\alpha^\beta B_n(t; a, b, c) dt = \frac{1}{(n+1) \ln c} [B_{n+1}(\beta; a, b, c) - B_{n+1}(\alpha; a, b, c)], \quad (14)$$

where  $B_0(x; a, b, c) = \frac{1}{\ln b - \ln a}$ .

*Proof.* Formula (13) follows from standard arguments and induction.

Integrating on both sides of (13) with respect to variable  $x$  for  $\ell = 1$  turns out formula (14).  $\square$

**Theorem 3.** *Let  $a, b, c > 0$ ,  $a \neq b$ ,  $n \geq 0$  and  $x \in \mathbb{R}$ . Then*

$$B_n(x+1; a, b, c) = \sum_{k=0}^n \binom{n}{k} (\ln c)^{n-k} B_k(x; a, b, c), \quad (15)$$

$$B_n(x+1; a, b, c) = B_n \left( x; \frac{a}{c}, \frac{b}{c}, c \right), \quad (16)$$

and, for  $m \geq 2$ ,

$$B_m(x+1; a, b, c) = B_m(x; a, b, c) + m(\ln c)^{m-1}x^{m-1} + \sum_{k=0}^{m-1} \binom{m}{k} [(\ln a)^{m-k} - (\ln b)^{m-k} + (\ln c)^{m-k}] B_k(x; a, b, c). \quad (17)$$

*Proof.* By definition of the generalized Bernoulli polynomials, we have

$$\frac{tc^{(x+1)t}}{b^t - a^t} = \sum_{n=0}^{\infty} \frac{B_n(x+1; a, b, c)}{n!} t^n \quad (18)$$

and

$$\begin{aligned} \frac{tc^{(x+1)t}}{b^t - a^t} &= \frac{tc^{xt}}{b^t - a^t} \cdot c^t \\ &= \left( \sum_{n=0}^{\infty} \frac{B_n(x; a, b, c)}{n!} t^n \right) \left( \sum_{k=0}^{\infty} \frac{(\ln c)^k}{k!} t^k \right) \\ &= \sum_{n=0}^{\infty} \frac{\sum_{k=0}^n \binom{n}{k} (\ln c)^{n-k} B_k(x; a, b, c)}{n!} t^n. \end{aligned} \quad (19)$$

Combining (18) and (19) and equating their coefficients of  $t^n$  leads to formula (15).

Similarly, since

$$\frac{tc^{(x+1)t}}{b^t - a^t} = \frac{tc^{xt}}{\left(\frac{b}{c}\right)^t - \left(\frac{a}{c}\right)^t} = \sum_{n=0}^{\infty} \frac{B_n\left(x; \frac{a}{c}, \frac{b}{c}, c\right)}{n!} t^n, \quad (20)$$

equating the coefficients of  $t^n$  in (18) and (20) leads to formula (16).

Straightforward computation gives us that

$$\begin{aligned} \frac{tc^{(x+1)t}}{b^t - a^t} &= tc^{xt} + \frac{tc^{xt}(a^t - b^t + c^t)}{b^t - a^t} \\ &= \sum_{n=0}^{\infty} \frac{(\ln c)^n x^n}{n!} t^{n+1} \\ &\quad + \left( \sum_{n=0}^{\infty} \frac{B_n(x; a, b, c)}{n!} t^n \right) \left( \sum_{\ell=0}^{\infty} \frac{[(\ln a)^\ell - (\ln b)^\ell + (\ln c)^\ell]}{\ell!} t^\ell \right) \\ &= \sum_{n=0}^{\infty} \frac{(\ln c)^n x^n}{n!} t^{n+1} \\ &\quad + \sum_{n=0}^{\infty} \left[ \sum_{\ell=0}^n \binom{n}{\ell} [(\ln a)^{n-\ell} - (\ln b)^{n-\ell} + (\ln c)^{n-\ell}] B_\ell(x; a, b, c) \right] \frac{t^n}{n!} \end{aligned} \quad (21)$$

$$\begin{aligned}
&= B_0(x; a, b, c) + [1 + B_1(x; a, b, c) + B_0(x; a, b, c)(\ln a - \ln b + \ln c)]t \\
&\quad + \sum_{n=2}^{\infty} [n(\ln c)^{n-1}x^{n-1} + B_n(x; a, b, c)] \frac{t^n}{n!} \\
&\quad + \sum_{n=2}^{\infty} \left\{ \sum_{\ell=0}^{n-1} \binom{n}{\ell} [(\ln a)^{n-\ell} - (\ln b)^{n-\ell} + (\ln c)^{n-\ell}] B_\ell(x; a, b, c) \right\} \frac{t^n}{n!}.
\end{aligned}$$

Equating (5) and (21) yields (17).  $\square$

**Corollary 1.** For  $n \geq 1$ ,  $b > 0$  and  $x \in \mathbb{R}$ , we have

$$B_n(x+1; 1, b, b) = B_n(x; 1, b, b) + n(\ln b)^{n-1}x^{n-1}. \quad (22)$$

*Remark 1.* If taking  $b = e$  in (22), the following well known result is deduced

$$B_n(x+1) = B_n(x) + nx^{n-1}, \quad n \geq 1. \quad (23)$$

Similarly, from (13) it follows that

$$B'_i(t) = iB_{i-1}(t), \quad B_0(t) = 1. \quad (24)$$

Actually, the Bernoulli polynomials  $B_i(t)$ ,  $i \in \mathbb{N}$ , are uniquely determined by formulae (23) and (24), see [1, 23.1.5 and 23.1.6] or [4].

**Theorem 4.** Let  $a, b, c > 0$ ,  $a \neq b$ ,  $n \geq 0$ , and  $x \in \mathbb{R}$ . Then

$$B_n(1-x; a, b, c) = (-1)^n B_n\left(x; \frac{c}{b}, \frac{c}{a}, c\right) \quad (25)$$

$$= B_n\left(-x; \frac{a}{c}, \frac{b}{c}, \frac{1}{c}\right), \quad (26)$$

$$B_n(x+y; a, b, c) = \sum_{k=0}^n \binom{n}{k} (\ln c)^{n-k} B_k(x; a, b, c) y^{n-k} \quad (27)$$

$$= \sum_{k=0}^n \binom{n}{k} (\ln c)^{n-k} B_k(y; a, b, c) x^{n-k}. \quad (28)$$

*Proof.* From Definition 5, it follows that

$$\frac{tc^{(1-x)t}}{b^t - a^t} = \sum_{n=0}^{\infty} \frac{B_n(1-x; a, b, c)}{n!} t^n. \quad (29)$$

Meanwhile, we have

$$\frac{tc^{(1-x)t}}{b^t - a^t} = \frac{tc^{-xt}}{\left(\frac{b}{c}\right)^t - \left(\frac{a}{c}\right)^t} = \sum_{n=0}^{\infty} \frac{B_n\left(-x; \frac{a}{c}, \frac{b}{c}, c\right)}{n!} t^n, \quad (30)$$

$$\frac{tc^{(1-x)t}}{b^t - a^t} = \frac{-tc^{x(-t)}}{\left(\frac{c}{a}\right)^{-t} - \left(\frac{c}{b}\right)^{-t}} = \sum_{n=0}^{\infty} (-1)^n \frac{B_n\left(x; \frac{c}{b}, \frac{c}{a}, c\right)}{n!} t^n. \quad (31)$$

Therefore, formulae (25) and (26) follow from equating series-expansions in (29), (30) and (31).

Similarly, we have

$$\frac{tc^{(x+y)t}}{b^t - a^t} = \sum_{n=0}^{\infty} \frac{B_n(x+y; a, b, c)}{n!} t^n, \quad (32)$$

$$\begin{aligned} \frac{tc^{(x+y)t}}{b^t - a^t} &= \frac{tc^{xt}}{b^t - a^t} \cdot c^{yt} \\ &= \left( \sum_{n=0}^{\infty} \frac{B_n(x; a, b, c)}{n!} t^n \right) \left( \sum_{i=0}^{\infty} \frac{y^i (\ln c)^i}{i!} t^i \right) \end{aligned} \quad (33)$$

$$\begin{aligned} &= \sum_{n=0}^{\infty} \left( \sum_{k=0}^n \binom{n}{k} y^{n-k} (\ln c)^{n-k} B_k(x; a, b, c) \right) \frac{t^n}{n!}, \\ \frac{tc^{(x+y)t}}{b^t - a^t} &= \frac{tc^{yt}}{b^t - a^t} \cdot c^{xt} \\ &= \sum_{n=0}^{\infty} \left( \sum_{k=0}^n \binom{n}{k} x^{n-k} (\ln c)^{n-k} B_k(y; a, b, c) \right) \frac{t^n}{n!}. \end{aligned} \quad (34)$$

Hence, formulae (27) and (28) follow from equating series-expansions in (32), (33) and (34). The proof is complete.  $\square$

**Theorem 5.** *Let  $m, n$  be natural numbers. Then, for any positive number  $b$ , the following identity holds:*

$$\begin{aligned} \sum_{j=1}^m j^n &= \frac{1}{(n+1)(\ln b)^n} [B_{n+1}(m+1; 1, b, b) - B_{n+1}(0; 1, b, b)] \\ &= \frac{1}{(n+1)(\ln b)^n} [B_{n+1}(m+1; 1, b, b) - B_{n+1}(1; 1, b, b)]. \end{aligned} \quad (35)$$

*Proof.* Rewriting formula (22) yields

$$x^{n-1} = \frac{1}{n(\ln b)^{n-1}} [B_n(x+1; 1, b, b) - B_n(x; 1, b, b)], \quad (36)$$

which implies

$$j^n = \frac{1}{(n+1)(\ln b)^n} [B_{n+1}(j+1; 1, b, b) - B_{n+1}(j; 1, b, b)]. \quad (37)$$

Summing up on both sides of (37) from 0 to  $m$  or from 1 to  $m$  with respect to  $j$  leads to formula (35) easily.  $\square$

*Remark 2.* The calculation of values of  $\sum_{j=1}^m j^n$  is an interesting problem, and there has been a rich literature, for example [3].

*Remark 3.* At last, it is pointed out that the Bernoulli's and Euler's numbers and the Bernoulli's and Euler's polynomials can be further generalized to more general results in this manner. These conclusions will be published in some subsequent papers.

**Acknowledgements.** This paper was finalized while the third author's visit to the RGMIA between November 1, 2001 and January 31, 2002, as a Visiting Professor with grants from the Victoria University of Technology and Jiaozuo Institute of Technology.

#### REFERENCES

- [1] M. Abramowitz and I. A. Stegun (Eds), *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, National Bureau of Standards, Applied Mathematics Series 55, 4th printing, Washington, 1965.
- [2] B.-N. Guo and F. Qi, *Generalisation of Bernoulli polynomials*, Internat. J. Math. Ed. Sci. Tech. (2002), in the press. RGMIA Res. Rep. Coll. 4 (2001), no. 4, Art. 10, 591–595. Available online at <http://rgmia.vu.edu.au/v4n4.html>.
- [3] S.-L. Guo and F. Qi, *Recursion formulae for  $\sum_{m=0}^n m^k$* , Z. Anal. Anwendungen **18** (1999), no. 4, 1123–1130.
- [4] Zh.-X. Wang and D.-R. Guo, *Tèshū Hánshù Gàilùn (Introduction to Special Function)*, The Series of Advanced Physics of Peking University, Peking University Press, Beijing, China, 2000. (Chinese)

(Luo) DEPARTMENT OF BROADCAST-TELEVISION-TEACHING, JIAOZUO UNIVERSITY, JIAOZUO CITY, HENAN 454002, CHINA

*E-mail address:* [luoqm236@sohu.com](mailto:luoqm236@sohu.com)

(Guo) DEPARTMENT OF APPLIED MATHEMATICS AND INFORMATICS, JIAOZUO INSTITUTE OF TECHNOLOGY, JIAOZUO CITY, HENAN 454000, CHINA

*E-mail address:* [guobaini@jzit.edu.cn](mailto:guobaini@jzit.edu.cn)

(Qi) DEPARTMENT OF APPLIED MATHEMATICS AND INFORMATICS, JIAOZUO INSTITUTE OF TECHNOLOGY, JIAOZUO CITY, HENAN 454000, CHINA

*E-mail address:* [qifeng@jzit.edu.cn](mailto:qifeng@jzit.edu.cn) or [qifeng618@hotmail.com](mailto:qifeng618@hotmail.com)

*URL:* <http://rgmia.vu.edu.au/qi.html>