



**VICTORIA UNIVERSITY**  
MELBOURNE AUSTRALIA

*A Refinement of the Grüss Inequality and Applications*

This is the Published version of the following publication

Cerone, Pietro and Dragomir, Sever S (2002) A Refinement of the Grüss Inequality and Applications. RGMIA research report collection, 5 (2).

The publisher's official version can be found at

Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/17716/>

# A REFINEMENT OF THE GRÜSS INEQUALITY AND APPLICATIONS

P. CERONE AND S.S. DRAGOMIR

ABSTRACT. A sharp refinement of the Grüss inequality in the general setting of measurable spaces and abstract Lebesgue integrals is proven. Some consequential particular inequalities are mentioned.

## 1. INTRODUCTION

Let  $(\Omega, \mathcal{A}, \mu)$  be a measurable space consisting of a set  $\Omega$ , a  $\sigma$ -algebra  $\mathcal{A}$  of parts of  $\Omega$  and a countably additive and positive measure  $\mu$  on  $\mathcal{A}$  with values in  $\mathbb{R} \cup \{\infty\}$ .

For a  $\mu$ -measurable function  $w : \Omega \rightarrow \mathbb{R}$ , with  $w(x) \geq 0$  for  $\mu$ -a.e.  $x \in \Omega$ , consider the Lebesgue space  $L_w(\Omega, \mu) := \{f : \Omega \rightarrow \mathbb{R}, f \text{ is } \mu\text{-measurable and } \int_{\Omega} w(x) |f(x)| d\mu(x) < \infty\}$ . Assume  $\int_{\Omega} w(x) d\mu(x) > 0$ .

If  $f, g : \Omega \rightarrow \mathbb{R}$  are  $\mu$ -measurable functions and  $f, g, fg \in L_w(\Omega, \mu)$ , then we may consider the Čebyšev functional

$$(1.1) \quad T_w(f, g) := \frac{1}{\int_{\Omega} w(x) d\mu(x)} \int_{\Omega} w(x) f(x) g(x) d\mu(x) \\ - \frac{1}{\int_{\Omega} w(x) d\mu(x)} \int_{\Omega} w(x) f(x) d\mu(x) \\ \times \frac{1}{\int_{\Omega} w(x) d\mu(x)} \int_{\Omega} w(x) g(x) d\mu(x).$$

The following result is known in the literature as the Grüss inequality

$$(1.2) \quad |T_w(f, g)| \leq \frac{1}{4} (\Gamma - \gamma) (\Delta - \delta),$$

provided

$$(1.3) \quad -\infty < \gamma \leq f(x) \leq \Gamma < \infty, \quad -\infty < \delta \leq g(x) \leq \Delta < \infty$$

for  $\mu$ -a.e.  $x \in \Omega$ .

The constant  $\frac{1}{4}$  is sharp in the sense that it cannot be replaced by a smaller constant.

Note that if  $\Omega = \{1, \dots, n\}$  and  $\mu$  is the discrete measure on  $\Omega$ , then we obtain the discrete Grüss inequality

$$(1.4) \quad \left| \frac{1}{W_n} \sum_{i=1}^n w_i x_i y_i - \frac{1}{W_n} \sum_{i=1}^n w_i x_i \cdot \frac{1}{W_n} \sum_{i=1}^n w_i y_i \right| \leq \frac{1}{4} (\Gamma - \gamma) (\Delta - \delta),$$

*Date:* April 10, 2002.

*1991 Mathematics Subject Classification.* Primary 26D15, 26D20; Secondary 26D10.

*Key words and phrases.* Grüss inequality, Measurable functions, Lebesgue integral, Perturbed Rules.

provided  $\gamma \leq x_i \leq \Gamma$ ,  $\delta \leq y_i \leq \Delta$  for each  $i \in \{1, \dots, n\}$  and  $w_i \geq 0$  with  $W_n := \sum_{i=1}^n w_i > 0$ .

The following result was proved in Cheng and Sun [4].

**Theorem 1.** *Let  $f, g : [a, b] \rightarrow \mathbb{R}$  be two integrable functions such that  $\delta \leq g(x) \leq \Delta$  for some constants  $\delta, \Delta$  for all  $x \in [a, b]$ , then*

$$(1.5) \quad \left| \frac{1}{b-a} \int_a^b f(x)g(x) dx - \frac{1}{(b-a)^2} \int_a^b f(x) dx \int_a^b g(x) dx \right| \\ \leq \frac{\Delta - \delta}{2} \int_a^b \left| f(x) - \frac{1}{b-a} \int_a^b f(y) dy \right| dx.$$

They used the result (1.5) to obtain perturbed trapezoidal rules.

In the current paper we obtain bounds for  $|T_w(f, g)|$  under the general setting expressed in (1.1). A bound which is shown to be *sharp* is obtained in Section 2. The sharpness of (1.5) was not demonstrated in [4]. Sharp results were obtained for a perturbed interior point rule (Ostrowski-Grüss) inequalities in Cheng [3]. Some particular instances of the results in Section 2 are investigated in Sections 4 and 5, recapturing earlier work. Results are presented in Section 3, for Lebesgue measurable functions and for a discrete weighted Čebyšev functional involving  $n$ -tuples.

## 2. AN INTEGRAL INEQUALITY

With the assumptions as presented in the Introduction and if  $f \in L_w(\Omega, \mu)$  then we may define

$$(2.1) \quad D_w(f) := D_{w,1}(f) \\ := \frac{1}{\int_{\Omega} w(x) d\mu(x)} \int_{\Omega} w(x) \\ \times \left| f(x) - \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} w(y) f(y) d\mu(y) \right| d\mu(x).$$

The following fundamental result holds.

**Theorem 2.** *Let  $w, f, g : \Omega \rightarrow \mathbb{R}$  be  $\mu$ -measurable functions with  $w \geq 0$   $\mu$ -a.e. on  $\Omega$  and  $\int_{\Omega} w(y) d\mu(y) > 0$ . If  $f, g, fg \in L_w(\Omega, \mu)$  and there exists the constants  $\delta, \Delta$  such that*

$$(2.2) \quad -\infty < \delta \leq g(x) \leq \Delta < \infty \quad \text{for } \mu\text{-a.e. } x \in \Omega,$$

then we have the inequality

$$(2.3) \quad |T_w(f, g)| \leq \frac{1}{2} (\Delta - \delta) D_w(f).$$

The constant  $\frac{1}{2}$  is sharp in the sense that it cannot be replaced by a smaller quantity.

*Proof.* Obviously, we have

$$(2.4) \quad T_w(f, g) = \frac{1}{\int_{\Omega} w(x) d\mu(x)} \int_{\Omega} w(x) \\ \times \left( f(x) - \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} f(y) w(y) d\mu(y) \right) g(x) d\mu(x).$$

Consider the measurable subsets  $\Omega_+$  and  $\Omega_-$ , of  $\Omega$ , defined by

$$\Omega_+ := \left\{ x \in \Omega \left| f(x) - \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} w(y) f(y) d\mu(y) \geq 0 \right. \right\}$$

and

$$\Omega_- := \left\{ x \in \Omega \left| f(x) - \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} w(y) f(y) d\mu(y) < 0 \right. \right\}.$$

Obviously,  $\Omega = \Omega_+ \cup \Omega_-$ ,  $\Omega_+ \cap \Omega_- = \emptyset$  and if we define

$$\begin{aligned} I_+(f, g, w) &:= \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega_+} w(x) \\ &\quad \times \left( f(x) - \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} w(y) f(y) d\mu(y) \right) g(x) d\mu(x) \end{aligned}$$

and

$$\begin{aligned} I_-(f, g, w) &:= \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega_-} w(x) \\ &\quad \times \left( f(x) - \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} w(y) f(y) d\mu(y) \right) g(x) d\mu(x) \end{aligned}$$

then we have

$$(2.5) \quad T_w(f, g) = I_+(f, g, w) + I_-(f, g, w).$$

Since  $-\infty < \delta \leq g(x) \leq \Delta < \infty$  for  $\mu$ -a.e.  $x \in \Omega$  and  $w(x) \geq 0$  for  $\mu$ -a.e.  $x \in \Omega$ , we may write:

$$(2.6) \quad \begin{aligned} I_+(f, g, w) &\leq \frac{\Delta}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega_+} w(x) \\ &\quad \times \left( f(x) - \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} f(y) w(y) d\mu(y) \right) d\mu(x) \end{aligned}$$

and

$$(2.7) \quad \begin{aligned} I_-(f, g, w) &\leq \frac{\delta}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega_-} w(x) \\ &\quad \times \left( f(x) - \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} f(y) w(y) d\mu(y) \right) d\mu(x). \end{aligned}$$

Since

$$\begin{aligned} 0 &= \int_{\Omega} w(x) \left( f(x) - \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} f(y) w(y) d\mu(y) \right) d\mu(x) \\ &= \int_{\Omega_+} w(x) \left( f(x) - \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} f(y) w(y) d\mu(y) \right) d\mu(x) \\ &\quad + \int_{\Omega_-} w(x) \left( f(x) - \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} f(y) w(y) d\mu(y) \right) d\mu(x) \end{aligned}$$

we get

$$\begin{aligned} & \int_{\Omega_-} w(x) \left( f(x) - \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} f(y) w(y) d\mu(y) \right) d\mu(x) \\ &= - \int_{\Omega_+} w(x) \left( f(x) - \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} f(y) w(y) d\mu(y) \right) d\mu(x) \end{aligned}$$

and thus, from (2.7), we deduce

$$(2.8) \quad I_-(f, g, w) \leq \frac{-\delta}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega_+} w(x) \times \left( f(x) - \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} w(y) f(y) d\mu(y) \right) d\mu(x).$$

Consequently, by adding (2.6) with (2.8), we deduce

$$(2.9) \quad T_w(f, g) \leq \frac{\Delta - \delta}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega_+} w(x) \times \left( f(x) - \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} w(y) f(y) d\mu(y) \right) d\mu(x).$$

On the other hand,

$$\begin{aligned} & \int_{\Omega} w(x) \left| f(x) - \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} f(y) w(y) d\mu(y) \right| d\mu(x) \\ &= \int_{\Omega_+} w(x) \left| f(x) - \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} f(y) w(y) d\mu(y) \right| d\mu(x) \\ &+ \int_{\Omega_-} w(x) \left| f(x) - \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} f(y) w(y) d\mu(y) \right| d\mu(x) \\ &= \int_{\Omega_+} w(x) \left( f(x) - \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} f(y) w(y) d\mu(y) \right) d\mu(x) \\ &- \int_{\Omega_-} w(x) \left( f(x) - \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} f(y) w(y) d\mu(y) \right) d\mu(x) \\ &= 2 \int_{\Omega_+} w(x) \left( f(x) - \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} f(y) w(y) d\mu(y) \right) d\mu(x), \end{aligned}$$

and thus, by (2.9) we deduce

$$(2.10) \quad T_w(f, g) \leq \frac{1}{2} (\Delta - \delta) \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} w(x) \times \left| f(x) - \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} f(y) w(y) d\mu(y) \right| d\mu(x).$$

Now, if we write the inequality (2.10) for  $-f$  instead of  $f$  and taking into account that  $T_w(-f, g) = -T_w(f, g)$ , we deduce

$$(2.11) \quad \begin{aligned} -T(f, g) &\leq \frac{1}{2} (\Delta - \delta) \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} w(x) \\ &\times \left| f(x) - \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} f(y) w(y) d\mu(y) \right| d\mu(x), \end{aligned}$$

giving the desired inequality (2.3).

To prove the sharpness of the constant  $\frac{1}{2}$ , assume that (2.3) holds for  $\Omega = [a, b]$  and  $w \equiv 1$ , with a constant  $C > 0$ . That is,

$$(2.12) \quad |T(f, g)| \leq C(\Delta - \delta) \frac{1}{b-a} \int_a^b \left| f(x) - \frac{1}{b-a} \int_a^b f(y) dy \right| dx,$$

where

$$T(f, g) = \frac{1}{b-a} \int_a^b f(x)g(x) dx - \frac{1}{b-a} \int_a^b f(x) dx \cdot \frac{1}{b-a} \int_a^b g(x) dx$$

and the integral  $\int_a^b$  is the usual Lebesgue integral on  $[a, b]$ .

Choose in (2.12)  $g = f$  and  $f : [a, b] \rightarrow \mathbb{R}$  defined by

$$f(x) = \begin{cases} -1 & \text{if } x \in [a, \frac{a+b}{2}], \\ 1 & \text{if } x \in (\frac{a+b}{2}, b], \end{cases}$$

then, obviously,

$$\begin{aligned} T(f, f) &= \frac{1}{b-a} \int_a^b f^2(x) dx - \left( \frac{1}{b-a} \int_a^b f(x) dx \right)^2 = 1, \\ D(f) &= \frac{1}{b-a} \int_a^b \left| f(x) - \frac{1}{b-a} \int_a^b f(y) dy \right| dx = 1, \\ \delta &= -1, \quad \Delta = 1, \end{aligned}$$

and by (2.12) we get  $2C \geq 1$  giving  $C \geq \frac{1}{2}$ . ■

For  $f \in L_{p,w}(\Omega, \mathcal{A}, \mu) := \{f : \Omega \rightarrow \mathbb{R}, \int_{\Omega} w(x) |f(x)|^p d\mu(x) < \infty\}$ ,  $p \geq 1$  we may also define

$$(2.13) \quad \begin{aligned} D_{w,p}(f) &:= \left[ \frac{1}{\int_{\Omega} w(x) d\mu(x)} \int_{\Omega} w(x) \right. \\ &\quad \times \left. \left| f(x) - \frac{1}{\int_{\Omega} w(y) d\mu(y)} \int_{\Omega} w(y) f(y) d\mu(y) \right|^p d\mu(x) \right]^{\frac{1}{p}} \\ &= \frac{\left\| f - \frac{1}{\int_{\Omega} w d\mu} \int_{\Omega} w f d\mu \right\|_{\Omega,p}}{\left[ \int_{\Omega} w(x) d\mu(x) \right]^{\frac{1}{p}}} \end{aligned}$$

where  $\|\cdot\|_{\Omega,p}$  is the usual  $p$ -norm on  $L_{p,w}(\Omega, \mathcal{A}, \mu)$ , namely,

$$\|h\|_{\Omega,p} := \left( \int_{\Omega} w |h|^p d\mu \right)^{\frac{1}{p}}, \quad p \geq 1.$$

Using Hölder's inequality we get

$$(2.14) \quad D_{w,1}(f) \leq D_{w,p}(f) \quad \text{for } p \geq 1, f \in L_{p,w}(\Omega, \mathcal{A}, \mu);$$

and, in particular for  $p = 2$

$$(2.15) \quad D_{w,1}(f) \leq D_{w,2}(f) = \left[ \frac{\int_{\Omega} w f^2 d\mu}{\int_{\Omega} w d\mu} - \left( \frac{\int_{\Omega} w f d\mu}{\int_{\Omega} w d\mu} \right)^2 \right]^{\frac{1}{2}},$$

if  $f \in L_{2,w}(\Omega, \mathcal{A}, \mu)$ .

For  $f \in L_\infty(\Omega, \mathcal{A}, \mu) := \left\{ f : \Omega \rightarrow \mathbb{R}, \|f\|_{\Omega, \infty} := \operatorname{ess\,sup}_{x \in \Omega} |f(x)| < \infty \right\}$  we also have

$$(2.16) \quad D_{w,p}(f) \leq D_{w,\infty}(f) := \left\| f - \frac{1}{\int_\Omega w d\mu} \int_\Omega w f d\mu \right\|_{\Omega, \infty}.$$

The following corollary may be useful in practice.

**Corollary 1.** *With the assumptions of Theorem 2, we have*

$$(2.17) \quad \begin{aligned} & |T_w(f, g)| \\ & \leq \frac{1}{2}(\Delta - \delta) D_w(f) \\ & \leq \frac{1}{2}(\Delta - \delta) D_{w,p}(f) \quad \text{if } f \in L_p(\Omega, \mathcal{A}, \mu), 1 < p < \infty; \\ & \leq \frac{1}{2}(\Delta - \delta) \left\| f - \frac{1}{\int_\Omega w d\mu} \int_\Omega w f d\mu \right\|_{\Omega, \infty} \quad \text{if } f \in L_\infty(\Omega, \mathcal{A}, \mu). \end{aligned}$$

**Remark 1.** *The inequalities in (2.17) are in order of increasing coarseness. If we assume that  $-\infty < \gamma \leq f(x) \leq \Gamma < \infty$  for  $\mu$ -a.e.  $x \in \Omega$ , then by the Grüss inequality for  $g = f$  we have for  $p = 2$*

$$(2.18) \quad \left[ \frac{\int_\Omega w f^2 d\mu}{\int_\Omega w d\mu} - \left( \frac{\int_\Omega w f d\mu}{\int_\Omega w d\mu} \right)^2 \right]^{\frac{1}{2}} \leq \frac{1}{2}(\Gamma - \gamma).$$

By (2.17), we deduce the following sequence of inequalities

$$(2.19) \quad \begin{aligned} |T_w(f, g)| & \leq \frac{1}{2}(\Delta - \delta) \frac{1}{\int_\Omega w d\mu} \int_\Omega w \left| f - \frac{1}{\int_\Omega w d\mu} \int_\Omega w f d\mu \right| d\mu \\ & \leq \frac{1}{2}(\Delta - \delta) \left[ \frac{\int_\Omega w f^2 d\mu}{\int_\Omega w d\mu} - \left( \frac{\int_\Omega w f d\mu}{\int_\Omega w d\mu} \right)^2 \right]^{\frac{1}{2}} \\ & \leq \frac{1}{4}(\Delta - \delta)(\Gamma - \gamma) \end{aligned}$$

for  $f, g : \Omega \rightarrow \mathbb{R}$ ,  $\mu$ -measurable functions and so that  $-\infty < \gamma \leq f(x) < \Gamma < \infty$ ,  $-\infty < \delta \leq g(x) \leq \Delta < \infty$  for  $\mu$ -a.e.  $x \in \Omega$ . Thus, the inequality (2.19) is a refinement of Grüss' inequality (1.2).

It is well known that if  $f \in L_{2,w}(\Omega, \mathcal{A}, \mu)$ , then the following Schwartz's type inequality holds:

$$(2.20) \quad \frac{1}{\int_\Omega w d\mu} \int_\Omega w f^2 d\mu \geq \left( \frac{1}{\int_\Omega w d\mu} \int_\Omega w f d\mu \right)^2.$$

Using the above results, we may point out the following counterpart result.

**Proposition 1.** *Assume that the  $\mu$ -measurable function  $f : \Omega \rightarrow \mathbb{R}$  satisfies the assumption:*

$$(2.21) \quad -\infty < \gamma \leq f(x) \leq \Gamma < \infty \quad \text{for a.e. } x \in \Omega.$$

Then one has the inequality

$$(2.22) \quad \begin{aligned} 0 &\leq \frac{1}{\int_{\Omega} w d\mu} \int_{\Omega} w f^2 d\mu - \left( \frac{1}{\int_{\Omega} w d\mu} \int_{\Omega} w f d\mu \right)^2 \\ &\leq \frac{1}{2} (\Gamma - \gamma) \frac{1}{\int_{\Omega} w d\mu} \int_{\Omega} w \left| f - \frac{1}{\int_{\Omega} w d\mu} \int_{\Omega} w f d\mu \right| d\mu \\ &\quad \left( \leq \frac{1}{4} (\Gamma - \gamma)^2 \right). \end{aligned}$$

The constant  $\frac{1}{2}$  is sharp.

The proof follows by the inequality (2.3) for  $g = f$ .

The following proposition also holds.

**Proposition 2.** *Assume that the measurable functions  $f, g : \Omega \rightarrow \mathbb{R}$  satisfy (1.3) (the condition in Grüss' inequality). Then*

$$(2.23) \quad \begin{aligned} |T_w(f, g)| &\leq \frac{1}{2} [(\Gamma - \gamma) (\Delta - \delta)]^{\frac{1}{2}} [D_w(f) D_w(g)]^{\frac{1}{2}} \\ &\leq \frac{1}{4} (\Delta - \delta) (\Gamma - \gamma). \end{aligned}$$

The constant  $\frac{1}{2}$  in the first inequality and  $\frac{1}{4}$  in the second inequality are sharp.

*Proof.* By (2.19) we have

$$|T_w(f, g)| \leq \frac{1}{2} (\Delta - \delta) D_w(f)$$

and

$$|T_w(f, g)| \leq \frac{1}{2} (\Gamma - \gamma) D_w(g)$$

from which, by multiplication, gives the first part of (2.23).

The second part and the sharpness of the constants are obvious. ■

### 3. SOME PARTICULAR INEQUALITIES

The following particular inequalities are of interest.

1. Let  $w, f, g : [a, b] \rightarrow \mathbb{R}$  be Lebesgue measurable functions with  $w \geq 0$  a.e. on  $[a, b]$  and  $\int_a^b w(y) dy > 0$ . If  $f, g, fg \in L_w[a, b]$ , where

$$L_w[a, b] := \left\{ f : [a, b] \rightarrow \mathbb{R} \mid \int_a^b w(x) |f(x)| dx < \infty \right\}$$

and

$$(3.1) \quad -\infty < \delta \leq g(x) \leq \Delta < \infty \quad \text{for a.e. } x \in [a, b],$$

then we have the inequalities

$$(3.2) \quad \left| \frac{1}{\int_a^b w(x) dx} \int_a^b w(x) f(x) g(x) dx - \frac{1}{\int_a^b w(x) dx} \int_a^b w(x) f(x) dx \cdot \frac{1}{\int_a^b w(x) dx} \int_a^b w(x) g(x) dx \right|$$



$$\begin{aligned}
&\leq \frac{1}{2} (\Delta - \delta) \frac{1}{\int_a^b w(x) dx} \int_a^b w(x) \left| f(x) - \frac{1}{\int_a^b w(y) dy} \int_a^b w(y) f(y) dy \right| dx \\
&\leq \frac{1}{2} (\Delta - \delta) \left[ \frac{\int_a^b w(x) \left| f(x) - \frac{1}{\int_a^b w(y) dy} \int_a^b w(y) f(y) dy \right|^p dx}{\int_a^b w(x) dx} \right]^{\frac{1}{p}} \\
&\text{if } f \in L_{p,w}[a, b], \quad 1 < p < \infty, \\
&\leq \frac{1}{2} (\Delta - \delta) \operatorname{ess\,sup}_{x \in [a, b]} \left| f(x) - \frac{1}{\int_a^b w(y) dy} \int_a^b w(y) f(y) dy \right| \text{ if } f \in L_\infty[a, b].
\end{aligned}$$

The constant  $\frac{1}{2}$  is sharp in the first inequality in (3.2).

The following counterpart of Schwartz's inequality holds

$$\begin{aligned}
(3.3) \quad 0 &\leq \frac{1}{\int_a^b w(y) dy} \int_a^b w(x) f^2(x) dx - \left( \frac{1}{\int_a^b w(y) dy} \int_a^b w(x) f(x) dx \right)^2 \\
&\leq \frac{1}{2} (\Delta - \gamma) \frac{1}{\int_a^b w(y) dy} \int_a^b w(x) \left| f(x) - \frac{1}{\int_a^b w(y) dy} \int_a^b w(y) f(y) dy \right| dx \\
&\quad \left( \leq \frac{1}{4} (\Gamma - \gamma)^2 \right),
\end{aligned}$$

provided  $-\infty < \gamma \leq f(x) \leq \Gamma < \infty$  for a.e.  $x \in [a, b]$ . The constant  $\frac{1}{2}$  is sharp.

If  $w(x) = 1$ ,  $x \in [a, b]$ , then we recapture the result in [4] as depicted here by (1.5).

**2.** Let  $\bar{\mathbf{a}} = (a_1, \dots, a_n)$ ,  $\bar{\mathbf{b}} = (b_1, \dots, b_n)$ ,  $\bar{\mathbf{p}} = (p_1, \dots, p_n)$  be  $n$ -tuples of real numbers with  $p_i \geq 0$  ( $i \in \{1, \dots, n\}$ ) and  $\sum_{i=1}^n p_i = 1$ . If

$$(3.4) \quad b \leq b_i \leq B, \quad i \in \{1, \dots, n\},$$

then one has the inequality

$$\begin{aligned}
(3.5) \quad &\left| \sum_{i=1}^n p_i a_i b_i - \sum_{i=1}^n p_i a_i \cdot \sum_{i=1}^n p_i b_i \right| \\
&\leq \frac{1}{2} (B - b) \sum_{i=1}^n p_i \left| a_i - \sum_{j=1}^n p_j a_j \right| \\
&\leq \frac{1}{2} (B - b) \left[ \sum_{i=1}^n p_i \left| a_i - \sum_{j=1}^n p_j a_j \right|^p \right]^{\frac{1}{p}} \quad \text{if } 1 < p < \infty \\
&\leq \frac{1}{2} (B - b) \max_{i=1, n} \left| a_i - \sum_{j=1}^n p_j a_j \right|.
\end{aligned}$$

The constant  $\frac{1}{2}$  is sharp in the first inequality.

If  $p_i = 1$ ,  $i \in \{1, \dots, n\}$ , the following unweighted inequality may be stated

$$\begin{aligned}
 (3.6) \quad 0 &\leq \frac{1}{n} \sum_{i=1}^n a_i b_i - \frac{1}{n} \sum_{i=1}^n a_i \cdot \frac{1}{n} \sum_{i=1}^n b_i \\
 &\leq \frac{1}{2} (B - b) \frac{1}{n} \sum_{i=1}^n \left| a_i - \frac{1}{n} \sum_{j=1}^n a_j \right| \\
 &\leq \frac{1}{2} (B - b) \left( \frac{1}{n} \sum_{i=1}^n \left| a_i - \frac{1}{n} \sum_{j=1}^n a_j \right|^p \right)^{\frac{1}{p}} \\
 &\leq \frac{1}{2} (B - b) \max_{i=1, \dots, n} \left| a_i - \frac{1}{n} \sum_{j=1}^n a_j \right|.
 \end{aligned}$$

The following counterpart of Schwartz's inequality also holds

$$\begin{aligned}
 (3.7) \quad 0 &\leq \sum_{i=1}^n p_i a_i^2 - \left( \sum_{i=1}^n p_i a_i \right)^2 \leq \frac{1}{2} (A - a) \sum_{i=1}^n p_i \left| a_i - \sum_{j=1}^n p_j a_j \right| \\
 &\quad \left( \leq \frac{1}{4} (A - a)^2 \right),
 \end{aligned}$$

provided  $a \leq a_i \leq A$  for each  $i \in \{1, \dots, n\}$  and  $\sum_{i=1}^n p_i = 1$ . The constant  $\frac{1}{2}$  is sharp.

#### 4. APPLICATIONS FOR OSTROWSKI'S INEQUALITY

If  $\varphi : [a, b] \rightarrow \mathbb{R}$  is an absolutely continuous function on  $[a, b]$  such that  $\varphi' \in L_\infty[a, b]$ , then the following inequality is known in the literature as Ostrowski's inequality

$$\begin{aligned}
 (4.1) \quad \left| \varphi(x) - \frac{1}{b-a} \int_a^b \varphi(t) dt \right| \\
 \leq \left[ \frac{1}{4} + \left( \frac{x - \frac{a+b}{2}}{b-a} \right)^2 \right] \|\varphi'\|_\infty (b-a), \quad x \in [a, b],
 \end{aligned}$$

where  $\|\varphi'\|_\infty := \operatorname{ess\,sup}_{\alpha \in [a, b]} |\varphi'(x)|$ . The constant  $\frac{1}{4}$  is best possible.

A simple proof of this fact, as mentioned in [1], may be accomplished by the use of the Montgomery identity

$$(4.2) \quad \varphi(x) = \frac{1}{b-a} \int_a^b \varphi(t) dt + \frac{1}{b-a} \int_a^b K(x, t) \varphi'(t) dt,$$

where the kernel  $K : [a, b]^2 \rightarrow \mathbb{R}$  is defined by

$$(4.3) \quad K(x, t) := \begin{cases} t - a & \text{if } a \leq t \leq x \\ t - b & \text{if } a \leq x < t \leq b. \end{cases}$$

We will now use the unweighted version of the inequality (3.2), namely, (1.5) (obtained by Cheng and Sun [4]) to procure the next result concerning a perturbed version of Ostrowski's inequality (4.1).

The following result also obtained by Cheng [3] is recaptured in a simpler manner. A weighted version of this result was obtained by Roumeliotis [5].

**Theorem 3.** *Assume that  $\varphi : [a, b] \rightarrow \mathbb{R}$  is an absolutely continuous function on  $[a, b]$  such that  $\varphi' : [a, b] \rightarrow \mathbb{R}$  satisfies the condition*

$$(4.4) \quad -\infty < \gamma \leq \varphi'(x) \leq \Gamma < \infty \quad \text{for a.e. } x \in [a, b].$$

Then we have the inequality

$$(4.5) \quad \left| \varphi(x) - \frac{1}{b-a} \int_a^b \varphi(t) dt - \left(x - \frac{a+b}{2}\right) [\varphi; a, b] \right| \leq \frac{1}{8} (b-a) (\Gamma - \gamma)$$

for any  $x \in [a, b]$ , where  $[\varphi; a, b] = \frac{\varphi(b) - \varphi(a)}{b-a}$  is the divided difference. The constant  $\frac{1}{8}$  is best possible.

*Proof.* We apply inequality (3.1) for the choices  $w(t) = 1$ ,  $f(t) = K(x, t)$  defined by (4.3),  $g(t) = \varphi'(t)$ ,  $t \in [a, b]$  to get

$$(4.6) \quad \left| \frac{1}{b-a} \int_a^b K(x, t) \varphi'(t) dt - \frac{1}{b-a} \int_a^b K(x, t) dt \cdot \frac{1}{b-a} \int_a^b \varphi'(t) dt \right| \\ \leq \frac{1}{2} (\Gamma - \gamma) \cdot \frac{1}{b-a} \int_a^b \left| K(x, t) - \frac{1}{b-a} \int_a^b K(x, s) ds \right| dt.$$

We obviously have,

$$\frac{1}{b-a} \int_a^b K(x, t) dt = x - \frac{a+b}{2}$$

and

$$\frac{1}{b-a} \int_a^b \varphi'(t) dt = \frac{\varphi(b) - \varphi(a)}{b-a}.$$

Also

$$I(x) := \frac{1}{b-a} \int_a^b \left| K(x, t) - \left(x - \frac{a+b}{2}\right) \right| dt \\ = \frac{1}{b-a} \left[ \int_a^x \left| t - a - x + \frac{a+b}{2} \right| dt + \int_x^b \left| t - b - x + \frac{a+b}{2} \right| dt \right] \\ = \frac{1}{b-a} \left[ \int_a^x \left| t - x + \frac{b-a}{2} \right| dt + \int_x^b \left| t - x - \frac{b-a}{2} \right| dt \right].$$

Straight forward substitution of  $u = t - x + \frac{b-a}{2}$  and  $v = t - x - \frac{b-a}{2}$  gives

$$I(x) = \frac{1}{b-a} \left[ \int_{\frac{a+b}{2}-x}^{\frac{b-a}{2}} |u| du + \int_{-\frac{b-a}{2}}^{\frac{a+b}{2}-x} |v| dv \right] \\ = \frac{1}{b-a} \int_{-\frac{b-a}{2}}^{\frac{b-a}{2}} |u| du = \frac{2}{b-a} \int_0^{\frac{b-a}{2}} u du = \frac{b-a}{4}.$$

Substitution of the above into (4.6) produces (4.5). The sharpness of the constant was proved in [3]. ■

## 5. APPLICATION FOR THE GENERALISED TRAPEZOID INEQUALITY

If  $\varphi : [a, b] \rightarrow \mathbb{R}$  is an absolutely continuous function on  $[a, b]$  so that  $\varphi' \in L_\infty [a, b]$ , then the following inequality is known as the generalised trapezoid inequality

$$(5.1) \quad \left| (x-a)\varphi(a) + (b-x)\varphi(b) - \int_a^b \varphi(t) dt \right| \leq \left[ \frac{1}{4}(b-a)^2 + \left(x - \frac{a+b}{2}\right)^2 \right] \|\varphi'\|_\infty$$

for any  $x \in [a, b]$ . The constant  $\frac{1}{4}$  is best possible.

A simple proof of this fact is accomplished by using the identity [2]

$$(5.2) \quad \int_a^b \varphi(t) dt = (x-a)\varphi(a) + (b-x)\varphi(b) + \int_a^b (x-t)\varphi'(t) dt.$$

Utilising the inequality (3.1) we may point out the following perturbed version of (5.1).

**Theorem 4.** *Assume that  $\varphi : [a, b] \rightarrow \mathbb{R}$  is an absolutely continuous function on  $[a, b]$  so that  $\varphi' : [a, b] \rightarrow \mathbb{R}$  satisfies the condition (4.4). Then we have the inequality*

$$(5.3) \quad \left| \frac{1}{b-a} \int_a^b \varphi(t) dt - \left[ \left(\frac{x-a}{b-a}\right)\varphi(a) + \left(\frac{b-x}{b-a}\right)\varphi(b) \right] - \left(x - \frac{a+b}{2}\right) [\varphi; a, b] \right| \leq \frac{1}{8} (b-a) (\Gamma - \gamma)$$

for any  $x \in [a, b]$ , where  $[\varphi; a, b]$  is the divided difference. The constant  $\frac{1}{8}$  is sharp.

*Proof.* We apply inequality (3.2) for the choices  $f(t) = (x-t)$ ,  $g(t) = \varphi'(t)$ ,  $w(t) = 1$ ,  $t \in [a, b]$ , to get

$$(5.4) \quad \left| \frac{1}{b-a} \int_a^b (x-t)\varphi'(t) dt - \frac{1}{b-a} \int_a^b (x-t) dt \cdot \frac{1}{b-a} \int_a^b \varphi'(t) dt \right| \leq \frac{1}{2} (\Gamma - \gamma) \frac{1}{b-a} \int_a^b \left| (x-t) - \frac{1}{b-a} \int_a^b (x-s) ds \right| dt.$$

Since

$$\begin{aligned} \frac{1}{b-a} \int_a^b (x-t) dt &= \left(x - \frac{a+b}{2}\right), \\ \frac{1}{b-a} \int_a^b \varphi'(t) dt &= \frac{\varphi(b) - \varphi(a)}{b-a} = [\varphi; a, b] \end{aligned}$$

and

$$\begin{aligned} \frac{1}{b-a} \int_a^b \left| (x-t) - \frac{1}{b-a} \int_a^b (x-s) ds \right| dt &= \frac{1}{b-a} \int_a^b \left| x-t-x+\frac{a+b}{2} \right| dt \\ &= \frac{1}{b-a} \int_a^b \left| t-\frac{a+b}{2} \right| dt \\ &= \frac{b-a}{4}, \end{aligned}$$

from (5.4) we deduce the desired inequality (5.3).

The sharpness of the constant may be shown on choosing  $t = \frac{a+b}{2}$  and  $\varphi(t) = \left| t - \frac{a+b}{2} \right|$ ,  $t \in [a, b]$ . We omit the details. ■

#### REFERENCES

- [1] P. CERONE and S.S. DRAGOMIR, Midpoint-type rules from an inequalities point of view, *Handbook of Analytic-Computational Methods in Applied Mathematics*, Editor: G. Anastassiou, CRC Press, N.Y., (2000), 135-200.
- [2] P. CERONE and S.S. DRAGOMIR, Trapezoid-type rules from an inequalities point of view, *Handbook of Analytic-Computational Methods in Applied Mathematics*, Editor: G. Anastassiou, CRC Press, N.Y., (2000), 65-134.
- [3] X.L. CHENG, Improvement of some Ostrowski-Grüss type inequalities, *Computers Math. Applic.*, **42** (2001), 109-114.
- [4] X.L. CHENG and J. SUN, A note on the perturbed trapezoid inequality, *J. Ineq. Pure. & Appl. Math.*, **3**(2) (2002), Article 29. [ONLINE] [http://jipam.vu.edu.au/v3n2/046\\_01.html](http://jipam.vu.edu.au/v3n2/046_01.html)
- [5] J. ROUMELIOTIS, Improved weighted Ostrowski-Grüss type inequalities, *RGMA Research Report Collection*, **5**(1) (2002), Article 13. [ONLINE] <http://rgmia.vu.edu.au/v5n1.html>

SCHOOL OF COMMUNICATIONS AND INFORMATICS, VICTORIA UNIVERSITY OF TECHNOLOGY, PO BOX 14428, MCMC 8001, VICTORIA, AUSTRALIA

*E-mail address:* [pc@matilda.vu.edu.au](mailto:pc@matilda.vu.edu.au)

*URL:* <http://rgmia.vu.edu.au/cerone>

*E-mail address:* [sever@matilda.vu.edu.au](mailto:sever@matilda.vu.edu.au)

*URL:* <http://rgmia.vu.edu.au/SSDragomirWeb.html>