



**VICTORIA UNIVERSITY**  
MELBOURNE AUSTRALIA

*On Hadamard's Inequalities for the Convex Mappings Defined in Topological Groups and Connected Result*

This is the Published version of the following publication

Morassaei, Ali (2009) On Hadamard's Inequalities for the Convex Mappings Defined in Topological Groups and Connected Result. Research report collection, 12 (3).

The publisher's official version can be found at

Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/17756/>

# ON HADAMARD'S INEQUALITIES FOR THE CONVEX MAPPINGS DEFINED IN TOPOLOGICAL GROUPS AND CONNECTED RESULT

ALI MORASSAEI

*Department of Mathematics,  
Faculty of Sciences,  
University of Zanjan,  
Zanjan 45195-313, Iran.  
morassaei@znu.ac.ir*

## Abstract

In this paper, we study the Hadamard's inequality for midconvex and quasi-midconvex functions in topological groups. A mapping naturally connected with this inequality and related result is also pointed out. <sup>1</sup>

## 1 Introduction

Let  $f : I \rightarrow \mathbb{R}$  be a convex mapping defined on the interval  $I$  of real numbers and  $a, b \in I$  with  $a < b$ . The following double inequality:

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a)+f(b)}{2} \quad (1)$$

is known in the literature as Hadamard's inequality for convex mapping. Note that some of the classical inequalities for means can be derived from (1) for appropriate particular selections of the mapping  $f$ .

In the paper [4] (see also [5] and [6]) is considered the following mapping naturally connected with Hadamard's results:

$$H : [0, 1] \rightarrow \mathbb{R}, \quad H(t) = \frac{1}{b-a} \int_a^b f\left(tx + (1-t)\frac{a+b}{2}\right) dx.$$

The following properties of  $H$  hold:

- (i)  $H$  is convex and monotonic nondecreasing.

---

<sup>1</sup>2000 AMS Subject Classification: Primary 26A51, 22A10, Secondary 26D07, 26D15.

Key words and phrases: convex functions, Hadamard's inequality.

The author is supported in part by the Institute for Advanced Studies in Basic Sciences, Zanjan, Iran.

(ii) One has the bounds

$$\sup_{t \in [0,1]} H(t) = H(1) = \frac{1}{b-a} \int_a^b f(x) dx$$

and

$$\inf_{t \in [0,1]} H(t) = H(0) = f\left(\frac{a+b}{2}\right).$$

Another mapping also closely connected with Hadamard's inequality is the following one [5] (see also [6]):

$$F : [0, 1] \longrightarrow \mathbb{R}, \quad F(t) = \frac{1}{(b-a)^2} \int_a^b \int_a^b f(tx + (1-t)y) dx dy.$$

The properties of this mapping are the following ones:

- (i)  $F$  is convex and monotonic nonincreasing on  $[0, \frac{1}{2}]$  and nondecreasing on  $[\frac{1}{2}, 1]$ ;
- (ii)  $F$  is symmetrical relating the element  $\frac{1}{2}$ . That is,

$$F(t) = F(1-t) \quad \text{for all } t \in [0, 1];$$

(iii) One has the bounds

$$\sup_{t \in [0,1]} F(t) = F(0) = F(1) = \frac{1}{b-a} \int_a^b f(x) dx$$

and

$$\inf_{t \in [0,1]} F(t) = F\left(\frac{1}{2}\right) = \frac{1}{(b-a)^2} \int_a^b \int_a^b f\left(\frac{x+y}{2}\right) dx dy \geq f\left(\frac{a+b}{2}\right)$$

(iv) The following inequality holds:

$$F(t) \geq \max\{H(t), H(1-t)\} \quad \text{for all } t \in [0, 1].$$

Generalization of (1) for quasi convex functions defined on the real line is also well-known. It was established in [2] that for a quasi convex function  $f$  defined on  $[a, b]$  we have

$$f\left(\frac{a+b}{2}\right) \leq \frac{2}{b-a} \int_a^b f(x) dx.$$

In this paper we shall study generalizations of the left side of (1) inequality for some convex functions defined on an open subset of a topological group  $G$ .

## 2 A Secondary Result

Generalization of the left side of (1) for convex functions defined on a convex subset of  $\mathbb{R}^n$  is well-known. For example, if  $X \subset \mathbb{R}^n$  is a convex bounded symmetrical set (the latter means that  $x \in X \implies -x \in X$ ), than (c.f. [8])

$$f(0) \leq \frac{1}{\mu(X)} \int_X f(x) dx \tag{2}$$

for each lower semicontinuous convex function  $f : X \rightarrow \mathbb{R}$ , where  $\mu(X)$  is the volume of the set  $X$ . To show (2), consider the transformation of the  $\mathbb{R}^n$  in itself given by:

$$h : \mathbb{R}^n \rightarrow \mathbb{R}^n, \quad h = (h_1, h_2, \dots, h_n),$$

and

$$h_i(x_1, x_2, \dots, x_n) = -x_i, \quad i = 1, 2, \dots, n.$$

Then  $h(X) = X$  and since

$$\frac{D(h_1, h_2, \dots, h_n)}{D(x_1, x_2, \dots, x_n)} = \begin{vmatrix} -1 & 0 & 0 & \dots & 0 \\ 0 & -1 & 0 & \dots & 0 \\ 0 & 0 & -1 & \dots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & 0 & \dots & -1 \end{vmatrix} = (-1)^n.$$

we have the change variable:

$$\begin{aligned} & \int_X f(x_1, x_2, \dots, x_n) dx_1 dx_2 \dots dx_n \\ &= \int_X f(h_1(x_1, x_2, \dots, x_n), \dots, h_n(x_1, x_2, \dots, x_n)) \left| \frac{D(h_1, h_2, \dots, h_n)}{D(x_1, x_2, \dots, x_n)} \right| dx_1 dx_2 \dots dx_n \\ &= \int_X f(-x_1, -x_2, \dots, -x_n) dx_1 dx_2 \dots dx_n. \end{aligned}$$

Now, by the convexity of  $f$  on  $X$  we also have:

$$\begin{aligned} f(0, 0, \dots, 0) &= f\left(\frac{x_1 - x_1}{2}, \frac{x_2 - x_2}{2}, \dots, \frac{x_n - x_n}{2}\right) \\ &= f\left(\frac{(x_1, x_2, \dots, x_n) + (-x_1, -x_2, \dots, -x_n)}{2}\right) \\ &\leq \frac{1}{2} [f(x_1, x_2, \dots, x_n) + f(-x_1, -x_2, \dots, -x_n)] \end{aligned}$$

which gives, by integration of  $f$  on  $X$ , that:

$$\begin{aligned} & \int_X f(0, 0, \dots, 0) dx_1 dx_2 \dots dx_n \\ &\leq \frac{1}{2} \left[ \int_X f(x_1, x_2, \dots, x_n) dx_1 dx_2 \dots dx_n + \int_X f(-x_1, -x_2, \dots, -x_n) dx_1 dx_2 \dots dx_n \right] \\ &= \int_X f(x_1, x_2, \dots, x_n). \end{aligned}$$

Consequently, we get

$$f(0, 0, \dots, 0) \mu(X) \leq \int_X f(x_1, x_2, \dots, x_n) dx_1 dx_2 \dots dx_n$$

and thus

$$f(0, 0, \dots, 0) \leq \frac{1}{\mu(X)} \int_X f(x_1, x_2, \dots, x_n) dx_1 dx_2 \dots dx_n.$$

### 3 Hadamard's Inequality

In this section we prove the Hadamard's inequality for midconvex and quasi-midconvex functions in topological groups.

Let  $G$  be a topological group, and  $\Omega$  is a nonempty open subset of  $G$  and  $f$  is a real-valued function on  $\Omega$ . We say that  $f$  is *globally (right) midconvex* if

$$2f(a) \leq f(az) + f(az^{-1}) \quad (3)$$

for all  $a, z \in G$  such that  $a, az, az^{-1} \in \Omega$ . Also, we say that  $f$  is *locally (right) midconvex in  $a \in \Omega$*  if there exists an open symmetric set  $V = V^{-1}$  from  $e$  such that

$$2f(a) \leq f(az) + f(az^{-1}) \quad (4)$$

for all  $z \in G$  such that  $az, az^{-1} \in \Omega$  [1]. Also,  $f$  is called *quasi-(right)midconvex*, if

$$f(az) \leq \max\{f(a), f(az^2)\} \quad (5)$$

for every  $a, z \in G$  so that  $a, az, az^2 \in \Omega$  [7]. Note that  $a$  is midpoint of  $az^{-1}$  and  $az$ , and  $az$  is midpoint of  $a$  and  $az^2$ .

**Definition 1.** Let  $\Omega$  be an open subset of topological group  $G$ , and  $a \in G$ .  $\Omega$  is said to be *symmetric relative to  $a$* , if  $a^{-1}\Omega$  is symmetric and  $e \in a^{-1}\Omega$ .

**Definition 2.** Let  $G$  be a topological group and  $\Omega \subset G$  is an open set. A function  $\omega : \Omega \rightarrow \mathbb{R}$  is called *symmetric relative to  $a \in G$* , if

$$\forall z \in G; \quad az, az^{-1} \in \Omega \quad \omega(az) = \omega(az^{-1}).$$

The following theorems hold:

**Theorem 1.** Let  $G$  be a locally compact group and  $\Omega \subset G$  is an open symmetric set relative to  $a \in G$  with  $0 < \mu(\Omega) < \infty$ . If  $f : \Omega \rightarrow \mathbb{R}$  is measurable and locally midconvex in  $a$  and also,  $f \in L_1(\Omega)$ , and  $\omega : \Omega \rightarrow \mathbb{R}$  is nonnegative and symmetric to  $a$  and  $\omega \in L_1(\Omega)$  such that  $f\omega \in L_1(\Omega)$ , then

$$f(a) \int_{\Omega} \omega(az) d\mu(z) \leq \int_{\Omega} f(az)\omega(az) d\mu(z),$$

where that  $\mu$  is Haar measure on  $G$ .

*Proof.* Since  $f$  is locally midconvex in  $a$ , so

$$2f(a) \leq f(az) + f(az^{-1})$$

for any  $z \in \Omega$ , by (4). Since  $\omega$  is nonnegative and symmetric relative to  $a$ , thus

$$\begin{aligned} 2f(a)\omega(az) &\leq f(az)\omega(az) + f(az^{-1})\omega(az) \\ &= f(az)\omega(az) + f(az^{-1})\omega(az^{-1}). \end{aligned}$$

Integrating this inequality on  $\Omega$ , we get

$$\begin{aligned}
\int_{\Omega} 2f(a)\omega(az)d\mu(z) &\leq \int_{\Omega} f(az)\omega(az)d\mu(z) + \int_{\Omega} f(az^{-1})\omega(az^{-1})d\mu(z) \\
&= \int_{a^{-1}\Omega} f(z)\omega(z)d\mu(z) + \int_{a^{-1}\Omega} f(z^{-1})\omega(z^{-1})d\mu(z) \\
&= \int_G f(z)\omega(z)\chi_{a^{-1}\Omega}(z)d\mu(z) + \int_G f(z^{-1})\omega(z^{-1})\chi_{a^{-1}\Omega}(z)d\mu(z) \\
&= \int_G f(z)\omega(z)\chi_{a^{-1}\Omega}(z)d\mu(z) + \int_G f(z^{-1})\omega(z^{-1})\chi_{a^{-1}\Omega}(z^{-1})d\mu(z) \\
&= 2 \int_G f(z)\omega(z)\chi_{a^{-1}\Omega}(z)d\mu(z). \\
&= 2 \int_{a^{-1}\Omega} f(z)\omega(z)d\mu(z).
\end{aligned}$$

Consequently, we have

$$f(a) \int_{\Omega} \omega(az)d\mu(z) \leq \int_{\Omega} f(az)\omega(az)d\mu(z).$$

□

**Remark 1.** If in the above theorem, set  $a = e$  and  $\omega \equiv 1$  on  $\Omega$ , we have

$$f(e) \leq \frac{1}{\mu(\Omega)} \int_{\Omega} f(z)d\mu(z).$$

This result is similar result of section 2.

**Theorem 2.** Let  $G$  be a locally compact group and  $\Omega \subset G$  is an open symmetric set relative to  $a \in G$  with  $0 < \mu(\Omega) < \infty$  and  $e \in \Omega$ . If  $f$  is measurable and quasi-midconvex real-valued function on  $\Omega$  such that  $f \in L_2(\Omega)$  and also,  $\omega : \Omega \rightarrow \mathbb{R}$  is a nonnegative and symmetric to  $a$  and  $\omega \in L_2(\Omega)$ , then

$$f(a) \int_{\Omega} \omega(az)d\mu(z) \leq \int_{\Omega} f(az)\omega(az)d\mu(z) + I(a) \quad (6)$$

where

$$I(a) = \frac{1}{2} \int_{\Omega} |f(az) - f(az^{-1})|\omega(az)d\mu(z).$$

Furthermore,  $I(a)$  satisfies the inequalities:

$$0 \leq I(a) \leq \min \left\{ \frac{\int_{\Omega} |f(az)|\omega(az)d\mu(z)}{\sqrt{2} \left( \int_{\Omega} f^2(az)d\mu(z) - \int_{\Omega} f(az)f(az^{-1})d\mu(z) \right)^{\frac{1}{2}}}, \left( \int_{\Omega} \omega^2(az)d\mu(z) \right)^{\frac{1}{2}} \right\}. \quad (7)$$

*Proof.* Since  $\Omega$  is symmetric set relative to  $a$ , thus for  $z$  in  $G$ , by (5), we have

$$f(a) \leq \max\{f(az), f(az^{-1})\}$$

where  $\max\{f(az), f(az^{-1})\} = \frac{f(az)+f(az^{-1})+|f(az)-f(az^{-1})|}{2}$  and since  $\omega$  is nonnegative and symmetric relative to  $a$ , therefore

$$\begin{aligned}
\int_{\Omega} f(a)\omega(az)d\mu(z) &\leq \frac{1}{2} \int_{\Omega} f(az)\omega(az)d\mu(z) + \frac{1}{2} \int_{\Omega} f(az^{-1})\omega(az^{-1})d\mu(z) \\
&\quad + \frac{1}{2} \int_{\Omega} |f(az) - f(az^{-1})|\omega(az)d\mu(z).
\end{aligned}$$

So,

$$f(a) \int_{\Omega} \omega(az) d\mu(z) \leq \int_{\Omega} f(az) \omega(az) d\mu(z) + I(a)$$

and the inequality (6) is proved.

We now observe that, by the Cauchy-Schwartz inequality,

$$\begin{aligned} 0 &\leq I(a) = \frac{1}{2} \int_{\Omega} |f(az) - f(az^{-1})| \omega(az) d\mu(z) \\ &\leq \frac{1}{2} \left( \int_{\Omega} (f(az) - f(az^{-1}))^2 d\mu(z) \right)^{\frac{1}{2}} \left( \int_{\Omega} \omega^2(az) d\mu(z) \right)^{\frac{1}{2}} \\ &= \frac{1}{2} \left( \int_{\Omega} [f^2(az) - 2f(az)f(az^{-1}) + f^2(az^{-1})] d\mu(z) \right)^{\frac{1}{2}} \left( \int_{\Omega} \omega^2(az) d\mu(z) \right)^{\frac{1}{2}} \\ &= \frac{1}{2} \left( 2 \int_{\Omega} [f^2(az) - f(az)f(az^{-1})] d\mu(z) \right)^{\frac{1}{2}} \left( \int_{\Omega} \omega^2(az) d\mu(z) \right)^{\frac{1}{2}} \\ &= \frac{\sqrt{2}}{2} \left( \int_{\Omega} f^2(az) d\mu(z) - \int_{\Omega} f(az)f(az^{-1}) d\mu(z) \right)^{\frac{1}{2}} \left( \int_{\Omega} \omega^2(az) d\mu(z) \right)^{\frac{1}{2}}. \end{aligned}$$

On the other hands,

$$\begin{aligned} I(a) &\leq \frac{1}{2} \left( \int_{\Omega} |f(az)| \omega(az) d\mu(z) + \int_{\Omega} |f(az^{-1})| \omega(az) d\mu(z) \right) \\ &= \int_{\Omega} |f(az)| \omega(az) d\mu(z) \end{aligned}$$

and the inequality (7) is proved.  $\square$

**Definition 3.** The function  $f : \Omega \rightarrow \mathbb{R}$  is said to be *P-function* in  $\Omega$ , if

$$f(a) \leq f(az) + f(az^{-1}),$$

for all  $a \in \Omega$  and  $z \in G$  such that  $az, az^{-1} \in \Omega$ .

**Theorem 3.** Let  $G$  be a locally compact group and  $\Omega \subset G$  is an open symmetric set relative to  $a \in G$  with  $0 < \mu(\Omega) < \infty$ . If  $f$  is measurable and *P-function* real-valued on  $\Omega$  such that  $f \in L_1(\Omega)$  and also,  $\omega : \Omega \rightarrow \mathbb{R}$  is nonnegative and symmetric to  $a$  and  $\omega \in L_1(\Omega)$  such that  $f\omega \in L_1(\Omega)$ , then

$$f(a) \int_{\Omega} \omega(az) d\mu(z) \leq 2 \int_{\Omega} f(az) \omega(az) d\mu(z).$$

*Proof.* Since  $f$  is *P-function*, we have

$$f(a) \omega(az) \leq f(az) \omega(az) + f(az^{-1}) \omega(az).$$

Integrating this inequality on  $\Omega$ , we get

$$\begin{aligned} \int_{\Omega} f(a) \omega(az) d\mu(z) &\leq \int_{\Omega} f(az) \omega(az) d\mu(z) + \int_{\Omega} f(az^{-1}) \omega(az) d\mu(z) \\ &= \int_{\Omega} f(az) \omega(az) d\mu(z) + \int_{\Omega} f(az^{-1}) \omega(az^{-1}) d\mu(z) \\ &= 2 \int_{\Omega} f(az) \omega(az) d\mu(z) \end{aligned}$$

Thus,  $f(a) \int_{\Omega} \omega(az) d\mu(z) \leq 2 \int_{\Omega} f(az) \omega(az) d\mu(z)$ . □

Now, for a globally midconvex function  $f : G \rightarrow \mathbb{R}$  that  $e \in \Omega$ , we can define the mapping  $H : \Omega \rightarrow \mathbb{R}$ ,

$$H(x) = \frac{1}{\mu(\Omega)} \int_{\Omega} f(xz) d\mu(z).$$

The properties of this mapping are embodied in the following theorem:

**Theorem 4.** *Suppose that  $f : G \rightarrow \mathbb{R}$  is globally midconvex and  $\Omega$  is an open symmetric subset of  $G$  such that  $e \in \Omega$  and  $0 < \mu(\Omega) < \infty$ . Then*

- (i) *The mapping  $H$  is globally midconvex on  $G$ , if  $G$  is abelian.*
- (ii)  *$f(e) \leq H(e)$ .*

*Proof.* (i) Assume that  $a, x, z \in G$  such that  $ax, ax^{-1} \in \Omega$ , so

$$\begin{aligned} H(ax) + H(ax^{-1}) &= \frac{1}{\mu(\Omega)} \int_{\Omega} f(axz) d\mu(z) + \frac{1}{\mu(\Omega)} \int_{\Omega} f(ax^{-1}z) d\mu(z) \\ &= \frac{1}{\mu(\Omega)} \int_{\Omega} [f(azx) + f(azx^{-1})] d\mu(z) \\ &\geq \frac{2}{\mu(\Omega)} \int_{\Omega} f(az) d\mu(z) \\ &= 2H(a) \end{aligned}$$

that is,  $H$  is globally midconvex.

(ii) Since  $f$  is midconvex, by Theorem 1,  $f(e) \leq \frac{1}{\mu(\Omega)} \int_{\Omega} f(z) d\mu(z) = H(e)$ . □

## 4 Applications

In this section, we study special cases of theorem 1, 2, and 3.

Set  $G = \mathbb{R}$ . Since  $\mathbb{R}$  is an abelian additive group, thus, for all  $a, z \in \mathbb{R}$ ,  $a - z$  and  $a + z$  are points that  $a$  is its midpoint. Now, if let  $a - z = x$  and  $a + z = y$ , then  $a = \frac{x+y}{2}$ . Consequently, definitions of globally midconvex and quasi-midconvex functions are to be written as follows:

$$\begin{aligned} f\left(\frac{x+y}{2}\right) &\leq \frac{f(x) + f(y)}{2}, & (f \text{ globally midconvex}) \\ f\left(\frac{x+y}{2}\right) &\leq \max\{f(x), f(y)\}. & (f \text{ quasi midconvex}) \end{aligned}$$

**Application 1.** If in the Theorem 1 let  $G = \mathbb{R}$  and  $\Omega = [-a, a]$ , we have

$$f(0) \int_{-a}^a \omega(x) dx \leq \int_{-a}^a f(x) \omega(x) dx,$$

and if we set  $\omega \equiv 1$  on  $[-a, a]$ , we get

$$f(0) \leq \frac{1}{2a} \int_{-a}^a f(x) dx$$



that is special case of (2), where  $n = 1$ .

**Application 2.** If in the Theorem 1, let  $G = \mathbb{R}^n$  with an additive operation and  $\Omega = X$  is an open bounded symmetric and convex subset of  $\mathbb{R}^n$ , then the result of section 2 holds.

## References

- [1] A. CHADEMAN AND F. MIRZAPOUR, *Midconvex functions in locally compact groups*, Proc. Amer. Math. Soc. **127**(10), (1999) 2961-2968.
- [2] S. S. DRAGOMIR AND C. E. M. PEARCE, *Quasi-convex functions and Hadamard's inequalities*, Bull. Austral. Math. Soc. **57**, (1998) 377-385.
- [3] S. S. DRAGOMIR AND C. E. M. PEARCE, *Select topics on Hermite-Hadamard inequalities and applications*, Melbourne and Adelaide, 2000.
- [4] S. S. DRAGOMIR, *A mapping in connection to Hadamard's inequality*, An Ostro. Akad. Wiss. Math. -Natur (Wien) **128** (1991) 17-20.
- [5] S. S. DRAGOMIR, *Two mappings in connection to Hadamard's inequality*, J. Math. Anal. Appl., **167** (1992) 49-56.
- [6] S. S. DRAGOMIR, *On Hadamard's inequality for convex functions*, Math. Balkanica, **6** (1992) 215-222.
- [7] F. MIRZAPOUR AND A. MORASSAEI, *Quasi-convex functions in topological groups*, Int. J. Appl. Math. **16**(4) (2004), 281-290.
- [8] A. M. RUBINOV, *Abstract convexity and global optimization*, Kluwer Academic Publishers, Dordrecht, 2000.