



**VICTORIA UNIVERSITY**  
MELBOURNE AUSTRALIA

*Some Inequalities in Inner Product Spaces Related to the Generalized Triangle Inequality*

This is the Published version of the following publication

Dragomir, Sever S, Cho, Yeol Je and Kim, Seong Sik (2009) Some Inequalities in Inner Product Spaces Related to the Generalized Triangle Inequality. Research report collection, 12 (3).

The publisher's official version can be found at

Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/17758/>

# SOME INEQUALITIES IN INNER PRODUCT SPACES RELATED TO THE GENERALIZED TRIANGLE INEQUALITY

S.S. DRAGOMIR, Y.J. CHO\*, AND S.S. KIM

ABSTRACT. In this paper we obtain some inequalities related to the generalized triangle and quadratic triangle inequalities for vectors in inner product spaces. Some results that employ the Ostrowski discrete inequality for vectors in normed linear spaces are also obtained.

## 1. INTRODUCTION

In Mathematical Analysis one of the most important inequalities is the (generalized) *triangle inequality* which states that, in a normed linear space  $(X; \|\cdot\|)$  we have the inequality

$$(1.1) \quad \left\| \sum_{i=1}^n x_i \right\| \leq \sum_{i=1}^n \|x_i\|$$

for any vectors  $x_i \in X$ ,  $i \in \{1, \dots, n\}$ .

In 1966, J.B. Diaz and F.T. Metcalf [3] proved the following reverse of the triangle inequality in inner product spaces:

**Theorem 1** (Diaz-Metcalf, 1966, [3]). *Let  $a$  be a unit vector in the inner product space  $(H; \langle \cdot, \cdot \rangle)$  over the real or complex number field  $\mathbb{K}$ . Suppose that the vectors  $x_i \in H \setminus \{0\}$ ,  $i \in \{1, \dots, n\}$  satisfy*

$$(1.2) \quad 0 \leq r \leq \frac{\operatorname{Re} \langle x_i, a \rangle}{\|x_i\|}, \quad i \in \{1, \dots, n\}.$$

Then

$$(1.3) \quad r \sum_{i=1}^n \|x_i\| \leq \left\| \sum_{i=1}^n x_i \right\|,$$

where equality holds if and only if

$$(1.4) \quad \sum_{i=1}^n x_i = r \left( \sum_{i=1}^n \|x_i\| \right) a.$$

The following result with a natural geometrical meaning also holds:

---

*Date:* July 10, 2009.

*1991 Mathematics Subject Classification.* 46D05, 46B05.

*Key words and phrases.* Triangle inequality, Inner product spaces, Normed spaces, Ostrowski inequality.

The second author was supported by the Korea Research Foundation Grant funded by the Korean Government (KRF-2008-313-C00050).

\*Corresponding author.

**Theorem 2** (Dragomir, 2004, [5]). *Let  $a$  be a unit vector in the inner product space  $(H; \langle \cdot, \cdot \rangle)$  and  $\rho \in (0, 1)$ . If  $x_i \in H$ ,  $i \in \{1, \dots, n\}$  are such that*

$$(1.5) \quad \|x_i - a\| \leq \rho \quad \text{for each } i \in \{1, \dots, n\},$$

*then we have the inequality*

$$(1.6) \quad \sqrt{1 - \rho^2} \sum_{i=1}^n \|x_i\| \leq \left\| \sum_{i=1}^n x_i \right\|,$$

*with equality if and only if*

$$(1.7) \quad \sum_{i=1}^n x_i = \sqrt{1 - \rho^2} \left( \sum_{i=1}^n \|x_i\| \right) a.$$

The following result that provides a more intuitive condition for the vectors involved may be stated as well:

**Theorem 3** (Dragomir, 2004, [5]). *Let  $a$  be a unit vector in the inner product space  $(H; \langle \cdot, \cdot \rangle)$  and  $M \geq m > 0$ . If  $x_i \in H$ ,  $i \in \{1, \dots, n\}$  are such that either*

$$(1.8) \quad \operatorname{Re} \langle Ma - x_i, x_i - ma \rangle \geq 0$$

*or, equivalently,*

$$(1.9) \quad \left\| x_i - \frac{M+m}{2} \cdot a \right\| \leq \frac{1}{2} (M-m)$$

*holds for each  $i \in \{1, \dots, n\}$ , then we have the inequality*

$$(1.10) \quad \frac{2\sqrt{mM}}{m+M} \sum_{i=1}^n \|x_i\| \leq \left\| \sum_{i=1}^n x_i \right\|,$$

*or, equivalently,*

$$(1.11) \quad (0 \leq) \sum_{i=1}^n \|x_i\| - \left\| \sum_{i=1}^n x_i \right\| \leq \frac{(\sqrt{M} - \sqrt{m})^2}{2\sqrt{mM}} \left\| \sum_{i=1}^n x_i \right\|.$$

*The equality holds in (1.10) (or in (1.11)) if and only if*

$$(1.12) \quad \sum_{i=1}^n x_i = \frac{2\sqrt{mM}}{m+M} \left( \sum_{i=1}^n \|x_i\| \right) a.$$

In inner product spaces  $(H; \langle \cdot, \cdot \rangle)$  another important inequality for sequences of vectors is the following one

$$(1.13) \quad \left\| \sum_{i=1}^n p_i x_i \right\|^2 \leq \sum_{i=1}^n p_i \|x_i\|^2,$$

where  $x_i \in H$ ,  $i \in \{1, \dots, n\}$  and  $p_i \geq 0$  with  $\sum_{i=1}^n p_i = 1$ . This can be seen as the quadratic version of the generalized triangle inequality.

A reverse of this inequality is incorporated in the following result:

**Theorem 4** (Dragomir, 2000, [4]). *Consider an inner product space  $(H; \langle \cdot, \cdot \rangle)$  over the real or complex number field  $\mathbb{K}$ . Suppose that the vectors  $x_i, x, X \in H$ ,  $i \in \{1, \dots, n\}$  satisfy the condition*

$$(1.14) \quad \operatorname{Re} \langle X - x_i, x_i - x \rangle \geq 0 \quad \text{for all } i \in \{1, \dots, n\},$$

or, equivalently,

$$(1.15) \quad \left\| x_i - \frac{x + X}{2} \right\| \leq \frac{1}{2} \|X - x\| \quad \text{for all } i \in \{1, \dots, n\},$$

then we have the inequality

$$(1.16) \quad \sum_{i=1}^n p_i \|x_i\|^2 - \left\| \sum_{i=1}^n p_i x_i \right\|^2 \leq \frac{1}{4} \|X - x\|^2$$

for any  $p_i \geq 0$ ,  $i \in \{1, \dots, n\}$ , with  $\sum_{i=1}^n p_i = 1$ .

The constant  $\frac{1}{4}$  is best possible in (1.16).

For other results related to the generalised triangle inequality, see [1], [2], [7], [8] and [9], where further references are provided.

In this paper we obtain other inequalities related to the generalized triangle and quadratic triangle inequalities for vectors in inner product spaces. Some results that employ the Ostrowski discrete inequality for vectors in normed linear spaces are also obtained.

## 2. THE RESULTS

The following result that encompasses a condition depending on the probability distribution  $p_i \geq 0$ ,  $i \in \{1, \dots, n\}$  can be stated:

**Theorem 5.** *Let  $(H; \langle \cdot, \cdot \rangle)$  be an inner product space,  $x_i \in H$ ,  $i \in \{1, \dots, n\}$  and  $p_i \geq 0$  with  $\sum_{i=1}^n p_i = 1$ . If there exist constants  $r_i > 0$ ,  $i \in \{1, \dots, n\}$ , so that*

$$(2.1) \quad \left\| x_i - \sum_{j=1}^n p_j x_j \right\| \leq r_i \quad \text{for each } i \in \{1, \dots, n\},$$

then

$$(2.2) \quad \sum_{i=1}^n p_i \|x_i\|^2 - \left\| \sum_{i=1}^n p_i x_i \right\|^2 \leq \sum_{i=1}^n p_i r_i^2.$$

The inequality (2.2) is sharp.

*Proof.* Taking the square in (2.1) we get

$$(2.3) \quad \|x_i\|^2 - 2 \operatorname{Re} \left\langle x_i, \sum_{j=1}^n p_j x_j \right\rangle + \left\| \sum_{j=1}^n p_j x_j \right\|^2 \leq r_i^2$$

for each  $i \in \{1, \dots, n\}$ .

Now, if we multiply (2.3) by  $p_i \geq 0$  and sum over  $i$  from 1 to  $n$ , we obtain

$$(2.4) \quad \sum_{i=1}^n p_i \|x_i\|^2 - 2 \operatorname{Re} \left\langle \sum_{i=1}^n p_i x_i, \sum_{j=1}^n p_j x_j \right\rangle + \left\| \sum_{j=1}^n p_j x_j \right\|^2 \leq \sum_{i=1}^n p_i r_i^2$$

and since

$$\operatorname{Re} \left\langle \sum_{i=1}^n p_i x_i, \sum_{j=1}^n p_j x_j \right\rangle = \left\| \sum_{i=1}^n p_i x_i \right\|^2$$

we deduce from (2.4) the desired inequality (2.2).

Now, assume that  $p_1, p_2 \in (0, 1)$  with  $p_1 + p_2 = 1$  and  $x_1, x_2 \in H$  with  $x_1 \neq x_2$ . With this assumption, (2.1) becomes

$$\|x_1 - p_1x_1 - p_2x_2\| \leq r_1, \quad \|x_2 - p_1x_1 - p_2x_2\| \leq r_2$$

which obviously holds for  $r_1 = p_2 \|x_1 - x_2\|$  and  $r_2 = p_1 \|x_1 - x_2\|$ .

Now, assume that there exists a constant  $c > 0$  such that (2.2) holds with  $c$ , i.e.,

$$(2.5) \quad \sum_{i=1}^n p_i \|x_i\|^2 - \left\| \sum_{i=1}^n p_i x_i \right\|^2 \leq c \sum_{i=1}^n p_i r_i^2.$$

Since

$$\begin{aligned} & p_1 \|x_1\|^2 + p_2 \|x_2\|^2 - \|p_1x_1 + p_2x_2\|^2 \\ &= p_1(1-p_1)\|x_1\|^2 + p_2(1-p_2)\|x_2\|^2 - 2p_1p_2 \operatorname{Re} \langle x_1, x_2 \rangle \\ &= p_1p_2 \left( \|x_1\|^2 + \|x_2\|^2 - 2 \operatorname{Re} \langle x_1, x_2 \rangle \right) \\ &= p_1p_2 \|x_1 - x_2\|^2 \end{aligned}$$

and

$$p_1r_1^2 + p_2r_2^2 = (p_1p_2^2 + p_2p_1^2) \|x_1 - x_2\|^2 = p_1p_2 \|x_1 - x_2\|^2,$$

hence (2.5) becomes

$$p_1p_2 \|x_1 - x_2\|^2 \leq cp_1p_2 \|x_1 - x_2\|^2,$$

implying that  $c \geq 1$ . This proves the sharpness of the inequality (2.2). ■

The following reverse of the generalized triangle inequality also holds.

**Theorem 6.** *Let  $x_i, p_i$  and  $r_i, i \in \{1, \dots, n\}$  be as in the statement of Theorem 5. Then*

$$(2.6) \quad (0 \leq) \sum_{i=1}^n p_i \|x_i\| - \left\| \sum_{i=1}^n p_i x_i \right\| \leq \frac{1}{2} \cdot \frac{\sum_{i=1}^n p_i r_i^2}{\left\| \sum_{i=1}^n p_i x_i \right\|},$$

provided that  $\sum_{i=1}^n p_i x_i \neq 0$ .

*Proof.* From (2.3) we obviously have

$$(2.7) \quad \|x_i\|^2 + \left\| \sum_{j=1}^n p_j x_j \right\|^2 \leq 2 \operatorname{Re} \left\langle x_i, \sum_{j=1}^n p_j x_j \right\rangle + r_i^2$$

for any  $i \in \{1, \dots, n\}$ .

Utilising the elementary inequality  $\alpha^2 + \beta^2 \geq 2\alpha\beta$ , with  $\alpha, \beta > 0$ , we also have

$$(2.8) \quad 2 \|x_i\| \left\| \sum_{j=1}^n p_j x_j \right\| \leq \|x_i\|^2 + \left\| \sum_{j=1}^n p_j x_j \right\|^2$$

for any  $i \in \{1, \dots, n\}$ , which together with (2.7) produces

$$(2.9) \quad \|x_i\| \left\| \sum_{j=1}^n p_j x_j \right\| \leq \operatorname{Re} \left\langle x_i, \sum_{j=1}^n p_j x_j \right\rangle + \frac{1}{2} r_i^2$$

for any  $i \in \{1, \dots, n\}$ .

Now, if we multiply (2.9) by  $p_i > 0$  and sum over  $i$  from 1 to  $n$ , we deduce

$$(2.10) \quad \sum_{i=1}^n p_i \|x_i\| \left\| \sum_{j=1}^n p_j x_j \right\| \leq \left\| \sum_{i=1}^n p_i x_i \right\|^2 + \frac{1}{2} \sum_{i=1}^n p_i r_i^2,$$

which, for  $\sum_{i=1}^n p_i x_i \neq 0$ , is equivalent to the desired result (2.6). ■

Another result is as follows.

**Theorem 7.** *Let  $x_i, p_i$  and  $r_i$  be as in the statement of Theorem 5. Then*

$$(2.11) \quad 0 \leq \left( \sum_{i=1}^n p_i \|x_i\|^2 \right)^{\frac{1}{2}} - \left\| \sum_{i=1}^n p_i x_i \right\| \leq \frac{1}{2} \cdot \frac{\sum_{i=1}^n p_i r_i^2}{\left\| \sum_{i=1}^n p_i x_i \right\|}.$$

*Proof.* If we multiply the inequality (2.7) by  $p_i \geq 0$  and sum over  $i$  from 1 to  $n$ , we deduce

$$(2.12) \quad \sum_{i=1}^n p_i \|x_i\|^2 + \left\| \sum_{j=1}^n p_j x_j \right\|^2 \leq 2 \left\| \sum_{j=1}^n p_j x_j \right\|^2 + \sum_{i=1}^n p_i r_i^2$$

for any  $i \in \{1, \dots, n\}$ .

However,

$$(2.13) \quad 2 \left( \sum_{i=1}^n p_i \|x_i\|^2 \right)^{\frac{1}{2}} \left\| \sum_{j=1}^n p_j x_j \right\| \leq \sum_{i=1}^n p_i \|x_i\|^2 + \left\| \sum_{j=1}^n p_j x_j \right\|^2$$

and by (2.12) and (2.13), we deduce the desired result (2.11). ■

**Remark 1.** *It is an open problem of whether the constant  $\frac{1}{2}$  is best possible in (2.6) or in (2.11). Also, since*

$$\sum_{i=1}^n p_i \|x_i\| \leq \left( \sum_{i=1}^n p_i \|x_i\|^2 \right)^{\frac{1}{2}},$$

*we observe that the inequality (2.11) is better than (2.6).*

The following result also holds.

**Theorem 8.** *With the assumptions of Theorem 5 and if, in addition, we have*

$$(0 <) \max_{i=1, n} \{r_i\} \leq \left\| \sum_{j=1}^n p_j x_j \right\| \quad \text{and} \quad x_i \neq 0 \quad \text{for all } i \in \{1, \dots, n\},$$

*then we obtain*

$$(2.14) \quad \left\| \sum_{j=1}^n p_j x_j \right\|^2 \leq \sum_{i=1}^n p_i r_i^2 + \operatorname{Re}^2 \left\langle \sum_{i=1}^n p_i \frac{x_i}{\|x_i\|}, \sum_{j=1}^n p_j x_j \right\rangle.$$

*This implies the inequality*

$$(2.15) \quad (0 \leq) 1 - \left\| \sum_{i=1}^n p_i \frac{x_i}{\|x_i\|} \right\|^2 \leq \frac{\sum_{i=1}^n p_i r_i^2}{\left\| \sum_{i=1}^n p_i x_i \right\|^2}.$$

*Proof.* From (2.3), we have

$$(2.16) \quad \|x_i\|^2 + \left[ \left( \left\| \sum_{j=1}^n p_j x_j \right\|^2 - r_i^2 \right)^{\frac{1}{2}} \right]^2 \leq 2 \operatorname{Re} \left\langle x_i, \sum_{j=1}^n p_j x_j \right\rangle$$

for each  $i \in \{1, \dots, n\}$  and since

$$(2.17) \quad 2 \|x_i\| \sqrt{\left\| \sum_{j=1}^n p_j x_j \right\|^2 - r_i^2} \leq \|x_i\|^2 + \left[ \left( \left\| \sum_{j=1}^n p_j x_j \right\|^2 - r_i^2 \right)^{\frac{1}{2}} \right]^2,$$

then by (2.16) and (2.17) we get

$$(2.18) \quad \|x_i\| \left( \left\| \sum_{j=1}^n p_j x_j \right\|^2 - r_i^2 \right)^{\frac{1}{2}} \leq \operatorname{Re} \left\langle x_i, \sum_{j=1}^n p_j x_j \right\rangle$$

for each  $i \in \{1, \dots, n\}$ . If we multiply (2.14) by  $\frac{p_i}{\|x_i\|} \geq 0$  and sum over  $i$  from 1 to  $n$ , we obtain

$$(2.19) \quad \sum_{i=1}^n \left( \left\| \sum_{j=1}^n p_j x_j \right\|^2 - r_i^2 \right)^{\frac{1}{2}} p_i \leq \operatorname{Re} \left\langle \sum_{i=1}^n p_i \frac{x_i}{\|x_i\|}, \sum_{j=1}^n p_j x_j \right\rangle.$$

However,

$$(2.20) \quad \sum_{i=1}^n p_i \left( \left\| \sum_{j=1}^n p_j x_j \right\|^2 - r_i^2 \right)^{\frac{1}{2}} \geq \left( \left\| \sum_{j=1}^n p_j x_j \right\|^2 - \sum_{i=1}^n p_i r_i^2 \right)^{\frac{1}{2}}$$

and by (2.19) and (2.20), we then have

$$\left\| \sum_{j=1}^n p_j x_j \right\|^2 \leq \sum_{i=1}^n p_i r_i^2 + \operatorname{Re}^2 \left\langle \sum_{i=1}^n p_i \frac{x_i}{\|x_i\|}, \sum_{j=1}^n p_j x_j \right\rangle,$$

which proves the desired inequality (2.14).

Now, on making use of the Schwarz inequality, we have

$$\operatorname{Re}^2 \left\langle \sum_{i=1}^n p_i \frac{x_i}{\|x_i\|}, \sum_{j=1}^n p_j x_j \right\rangle \leq \left\| \sum_{i=1}^n p_i \frac{x_i}{\|x_i\|} \right\|^2 \left\| \sum_{j=1}^n p_j x_j \right\|^2,$$

which together with (2.14) produces the desired result (2.15). ■

**Theorem 9.** *With the assumptions of Theorem 5 and if, in addition, we have*

$$(0 <) \max_{i=1, n} \{r_i\} \leq \left\| \sum_{j=1}^n p_j x_j \right\|,$$

then

$$(2.21) \quad (0 \leq) \sum_{i=1}^n p_i \|x_i\|^2 - \left\| \sum_{i=1}^n p_i x_i \right\|^2 \leq \frac{\sum_{i=1}^n p_i r_i^2 \|x_i\|^2}{\left\| \sum_{j=1}^n p_j x_j \right\|^2} \left( \leq \max_{i=1, n} r_i^2 \frac{\sum_{i=1}^n p_i \|x_i\|^2}{\left\| \sum_{j=1}^n p_j x_j \right\|^2} \right).$$

*Proof.* From (2.18) we have that

$$(2.22) \quad \sum_{i=1}^n p_i \|x_i\| \left( \left\| \sum_{j=1}^n p_j x_j \right\|^2 - r_i^2 \right)^{\frac{1}{2}} \leq \left\| \sum_{j=1}^n p_j x_j \right\|^2.$$

However,

$$(2.23) \quad \left[ \sum_{i=1}^n p_i \|x_i\|^2 \left( \left\| \sum_{j=1}^n p_j x_j \right\|^2 - r_i^2 \right) \right]^{\frac{1}{2}} \leq \sum_{i=1}^n p_i \|x_i\| \left( \left\| \sum_{j=1}^n p_j x_j \right\|^2 - r_i^2 \right)^{\frac{1}{2}}$$

and since

$$\sum_{i=1}^n p_i \|x_i\|^2 \left( \left\| \sum_{j=1}^n p_j x_j \right\|^2 - r_i^2 \right) = \sum_{i=1}^n p_i \|x_i\|^2 \left\| \sum_{j=1}^n p_j x_j \right\|^2 - \sum_{i=1}^n p_i \|x_i\|^2 r_i^2,$$

hence by (2.22) and (2.23) we obtain (2.21). ■

### 3. SOME RESULTS VIA OSTROWSKI'S INEQUALITY

In order to provide more convenient inequalities, the condition (2.1), namely

$$(3.1) \quad \left\| x_i - \sum_{j=1}^n p_j x_j \right\| \leq r_i \quad \text{for all } i \in \{1, \dots, n\},$$

where  $x_i \in H$ ,  $p_i \geq 0$  with  $i \in \{1, \dots, n\}$  and  $\sum_{i=1}^n p_i = 1$  can be replaced with a number of discrete Ostrowski inequalities such as those obtained by Dragomir in [6].

For the sake of completeness, we recall them here in a single theorem.

**Theorem 10** (Dragomir, 2002, [6]). *Let  $(X, \|\cdot\|)$  be a normed linear space,  $x_i \in X$  ( $i \in \{1, \dots, n\}$ ) and  $p_i \geq 0$ ,  $i \in \{1, \dots, n\}$  with  $\sum_{i=1}^n p_i = 1$ . Then we have the*



inequalities:

$$(3.2) \quad \left\| x_i - \sum_{j=1}^n p_j x_j \right\| \leq \max_{k \in \{1, \dots, n-1\}} \|\Delta x_k\| \sum_{j=1}^n p_j |j - i|$$

$$\leq \max_{k \in \{1, \dots, n-1\}} \|\Delta x_k\| \times \begin{cases} \frac{n-1}{2} + |i - \frac{n+1}{2}|; \\ \left( \sum_{j=1}^n |j - i|^p \right)^{\frac{1}{p}} \left( \sum_{j=1}^n p_j^q \right)^{\frac{1}{q}}, \\ p > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \left[ \frac{n^2-1}{4} + \left( i - \frac{n+1}{2} \right)^2 \right] \max_{j \in \{1, \dots, n\}} \{p_j\} \end{cases}$$

and, for  $\alpha > 1$ ,  $\frac{1}{\alpha} + \frac{1}{\beta} = 1$ , the inequalities:

$$(3.3) \quad \left\| x_i - \sum_{j=1}^n p_j x_j \right\| \leq \left( \sum_{k=1}^{n-1} \|\Delta x_k\|^\alpha \right)^{\frac{1}{\alpha}} \sum_{j=1}^n |i - j|^{\frac{1}{\beta}} p_j$$

$$\leq \left( \sum_{k=1}^{n-1} \|\Delta x_k\|^\alpha \right)^{\frac{1}{\alpha}} \begin{cases} \left[ \frac{n-1}{2} + |i - \frac{n+1}{2}| \right]^{\frac{1}{\beta}}; \\ \left( \sum_{j=1}^n |i - j|^{\frac{\delta}{\beta}} \right)^{\frac{1}{\delta}} \left( \sum_{j=1}^n p_j^\gamma \right)^{\frac{1}{\gamma}}, \\ \gamma > 1, \frac{1}{\gamma} + \frac{1}{\delta} = 1; \\ \sum_{j=1}^n |i - j|^{\frac{1}{\beta}} \max_{j \in \{1, \dots, n\}} \{p_j\} \end{cases}$$

and

$$(3.4) \quad \left\| x_i - \sum_{j=1}^n p_j x_j \right\| \leq \begin{cases} \max \{P_{i-1}, 1 - P_i\} \sum_{k=1}^{n-1} \|\Delta x_k\| \\ (1 - p_i) \max \left\{ \sum_{k=1}^{i-1} \|\Delta x_k\|, \sum_{k=i}^{n-1} \|\Delta x_k\| \right\} \end{cases}$$

$$\leq (1 - p_i) \sum_{k=1}^{n-1} \|\Delta x_k\|$$

for each  $i \in \{1, \dots, n\}$ , where  $\Delta x_k := x_{k+1} - x_k$  and  $P_m := \sum_{i=1}^m p_i$ ,  $k \in \{1, \dots, n-1\}$ ,  $m \in \{1, \dots, n\}$  and  $P_0 := 0$ .

Now, if we use Theorems 5 and 10 we can state the following result that provides upper bounds for the quantity

$$\sum_{i=1}^n p_i \|x_i\|^2 - \left\| \sum_{i=1}^n p_i x_i \right\|^2.$$

**Proposition 1.** Let  $(H; \langle \cdot, \cdot \rangle)$  be an inner product space,  $x_i \in H$ ,  $i \in \{1, \dots, n\}$  and  $p_i \geq 0$  with  $\sum_{i=1}^n p_i = 1$ . Then we have the inequalities:

$$(3.5) \quad \sum_{i=1}^n p_i \|x_i\|^2 - \left\| \sum_{i=1}^n p_i x_i \right\|^2 \leq \max_{k \in \{1, \dots, n-1\}} \|\Delta x_k\|^2 \sum_{i=1}^n p_i \left( \sum_{j=1}^n p_j |j-i| \right)^2 \leq \max_{k \in \{1, \dots, n-1\}} \|\Delta x_k\|^2 \begin{cases} \sum_{i=1}^n p_i \left( \frac{n-1}{2} + |i - \frac{n+1}{2}| \right)^2; \\ \left( \sum_{j=1}^n p_j^q \right)^{\frac{2}{q}} \sum_{i=1}^n \left( \sum_{j=1}^n |j-i|^p \right)^{\frac{2}{p}}, \\ p > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \max_{j \in \{1, \dots, n\}} \{p_j^2\} \sum_{i=1}^n p_i \left[ \frac{n^2-1}{4} + \left( i - \frac{n+1}{2} \right)^2 \right]^2 \end{cases}$$

and, for  $\alpha > 1$ ,  $\frac{1}{\alpha} + \frac{1}{\beta} = 1$ , the inequalities:

$$(3.6) \quad \sum_{i=1}^n p_i \|x_i\|^2 - \left\| \sum_{i=1}^n p_i x_i \right\|^2 \leq \left( \sum_{k=1}^{n-1} \|\Delta x_k\|^\alpha \right)^{\frac{2}{\alpha}} \sum_{i=1}^n p_i \left( \sum_{j=1}^n |i-j|^{\frac{1}{\beta}} p_j \right)^2 \leq \left( \sum_{k=1}^{n-1} \|\Delta x_k\|^\alpha \right)^{\frac{2}{\alpha}} \begin{cases} \sum_{i=1}^n p_i \left[ \frac{n-1}{2} + |i - \frac{n+1}{2}| \right]^{\frac{2}{\beta}}; \\ \left( \sum_{j=1}^n p_j^\gamma \right)^{\frac{2}{\gamma}} \sum_{i=1}^n p_i \left( \sum_{j=1}^n |i-j|^{\frac{\delta}{\beta}} \right)^{\frac{2}{\delta}}, \\ \gamma > 1, \frac{1}{\gamma} + \frac{1}{\delta} = 1; \\ \max_{j \in \{1, \dots, n\}} \{p_j^2\} \sum_{i=1}^n p_i \left( \sum_{j=1}^n |i-j|^{\frac{1}{\beta}} \right)^2 \end{cases}$$

and

$$(3.7) \quad \sum_{i=1}^n p_i \|x_i\|^2 - \left\| \sum_{i=1}^n p_i x_i \right\|^2 \leq \begin{cases} \left( \sum_{k=1}^{n-1} \|\Delta x_k\| \right)^2 \sum_{i=1}^n p_i \max \{P_{i-1}^2, (1-P_i)^2\} \\ \sum_{i=1}^n p_i (1-p_i)^2 \left( \max \left\{ \sum_{k=1}^{i-1} \|\Delta x_k\|, \sum_{k=i}^{n-1} \|\Delta x_k\| \right\} \right)^2 \end{cases} \leq \left( \sum_{k=1}^{n-1} \|\Delta x_k\| \right)^2 \sum_{i=1}^n p_i (1-p_i)^2.$$

**Remark 2.** If one uses the other inequalities obtained in Theorems 6 – 9, then other similar results can be stated. However the details are not presented here.

## REFERENCES

- [1] A. H. ANSARI and M. S. MOSLEHIAN, More on reverse triangle inequality in inner product spaces, *Int. J. Math. Math. Sci.*, **18**(2005), 2883–2893.
- [2] A. H. ANSARI and M. S. MOSLEHIAN, Refinements of reverse triangle inequalities in inner product spaces, *J. Inequal. Pure Appl. Math.*, **6** (2005), no. 3, Article 64, 12 pp.
- [3] J.B. DIAZ and F.T. METCALF, A complementary triangle inequality in Hilbert and Banach spaces, *Proceedings Amer. Math. Soc.*, **17**(1) (1966), 88-97.
- [4] S.S. DRAGOMIR, A Grüss' type discrete inequality in inner product spaces and applications, *J. Math. Anal. Appl.*, **250** (2000), 494-511.
- [5] S.S. DRAGOMIR, Reverses of the triangle inequality in inner product spaces, *Austral. J. Math. Anal Appl.*, **1**(2004), No. 2, Article 7, pp. 1-14. Preprint *RGMA Res. Rep. Coll.*, **7**(2004), Supplement, Article 7, [ONLINE: [http://rgmia.vu.edu.au/v7\(E\).html](http://rgmia.vu.edu.au/v7(E).html)].
- [6] S.S. DRAGOMIR, The discrete version of Ostrowski's inequality in normed linear spaces, *J. Inequal. Pure Appl. Math.*, **3**(2002), No. 1, Article 2, [Online [http://jipam.vu.edu.au/images/042\\_01\\_JIPAM/042\\_01.pdf](http://jipam.vu.edu.au/images/042_01_JIPAM/042_01.pdf)].
- [7] S.S. DRAGOMIR, A survey of recent reverses for the generalized triangle inequality in inner product spaces, *Kragujevac J. Math.*, **27** (2005), 101–143.
- [8] S. SAITOH, Generalizations of the triangle inequality, *J. Inequal. Pure Appl. Math.*, **4** (2003), no. 3, Article 62, 5 pp.
- [9] C. WU and Y. LI, On the triangle inequality in quasi-Banach spaces, *J. Inequal. Pure Appl. Math.*, **9** (2008), no. 2, Article 41, 4 pp.

MATHEMATICS, SCHOOL OF ENGINEERING & SCIENCE, VICTORIA UNIVERSITY, PO Box 14428, MELBOURNE CITY, MC 8001, AUSTRALIA.

*E-mail address:* [sever.dragomir@vu.edu.au](mailto:sever.dragomir@vu.edu.au)

*URL:* <http://www.staff.vu.edu.au/rgmia/dragomir/>

DEPARTMENT OF MATHEMATICAL EDUCATION AND THE RINS, GYEONGSANG NATIONAL UNIVERSITY, CHINJU 660-701, KOREA

*E-mail address:* [yjcho@nongae.gsnu.ac.kr](mailto:yjcho@nongae.gsnu.ac.kr)

DEPARTMENT OF MATHEMATICS, DONGEU UNIVERSITY, BUSAN 614-714, KOREA

*E-mail address:* [sskim@deu.ac.kr](mailto:sskim@deu.ac.kr)