



**VICTORIA UNIVERSITY**  
MELBOURNE AUSTRALIA

## *Three Inequalities Involving Hyperbolically Trigonometric Functions*

This is the Published version of the following publication

Chen, Chao-Ping, Zhao, Jian-Wei and Qi, Feng (2003) Three Inequalities Involving Hyperbolically Trigonometric Functions. RGMIA research report collection, 6 (3).

The publisher's official version can be found at

Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/17850/>

# THREE INEQUALITIES INVOLVING HYPERBOLICALLY TRIGONOMETRIC FUNCTIONS

CHAO-PING CHEN, JIAN-WEI ZHAO, AND FENG QI

ABSTRACT. In the short note, by using mathematical induction and infinite product representations of the cosine function, hyperbolic sine function and hyperbolic cosine function, three inequalities for the cosine function, hyperbolic sine function and hyperbolic cosine function are established.

## 1. INTRODUCTION

It is well-known that the following

$$\frac{2}{\pi} \leq \frac{\sin x}{x} \leq 1, \quad |x| \in \left(0, \frac{\pi}{2}\right]; \quad (1)$$

is called Jordan's inequality [8, p. 42].

Kober's inequality is given in [5] and [6, p. 317] as follows:

$$\cos x \geq 1 - \frac{2}{\pi}x, \quad x \in \left[0, \frac{\pi}{2}\right]. \quad (2)$$

For  $\frac{\pi}{2} < x < \pi$ , inequality (2) reverses.

These two inequalities are basic inequalities in calculus and in trigonometry.

In [12], R. Redheffer established that

$$\frac{\sin x}{x} \geq \frac{\pi^2 - x^2}{\pi^2 + x^2}, \quad x \in (-\infty, +\infty). \quad (3)$$

The inequality (3) and Jordan's inequality (1) do not imply each other.

The study of Jordan's and Kober's inequality and inequalities of trigonometric functions has a rich literature, for example, [1, 2, 4, 6, 7, 8, 9, 10, 11] and references therein. There are many refinements, extensions, and variants of them, each based on a different principle, or at least using a different device. A much complete list of references in recent years can be found in [2].

In this note, by using mathematical induction and infinite product representations of  $\cos x$ ,  $\sinh x$  and  $\cosh x$ , three inequalities for the cosine function, hyperbolic sine function and hyperbolic cosine function, which are similar to Redheffer's inequality (3), are established.

---

2000 *Mathematics Subject Classification.* Primary 26D05, 26D07.

*Key words and phrases.* inequality, cosine function, hyperbolic sine function, hyperbolic cosine function, infinite product representation, mathematical induction.

The first and third authors were supported in part by NNSF (#10001016) of China, SF for the Prominent Youth of Henan Province (#0112000200), SF of Henan Innovation Talents at Universities, Doctor Fund of Jiaozuo Institute of Technology, CHINA.

This paper was typeset using  $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{L}\mathcal{A}\mathcal{T}\mathcal{E}\mathcal{X}$ .

**Theorem 1.** *If  $|x| \leq \frac{1}{2}$ , then*

$$\cos(\pi x) \geq \frac{1 - 4x^2}{1 + 4x^2}, \quad (4)$$

$$\cosh(\pi x) \leq \frac{1 + 4x^2}{1 - 4x^2}. \quad (5)$$

*If  $0 < |x| < 1$ , then*

$$\frac{\sinh(\pi x)}{\pi x} \leq \frac{1 + x^2}{1 - x^2}. \quad (6)$$

## 2. PROOF OF THEOREM 1

*Proof of inequality (4).* It is sufficient to prove inequality (4) for  $0 < x < \frac{1}{2}$ .

In [3, p. 193], the following product representation is given

$$\cos(\pi x) = \prod_{n=1}^{\infty} \left(1 - \frac{4x^2}{(2n-1)^2}\right). \quad (7)$$

Set

$$F_n = \prod_{k=2}^n \left(1 - \frac{4x^2}{(2k-1)^2}\right), \quad n = 2, 3, \dots \quad (8)$$

Then we have

$$\cos(\pi x) = \frac{1 - 4x^2}{1 + 4x^2} \left[ (1 + 4x^2) \lim_{n \rightarrow \infty} F_n \right] \quad (9)$$

and

$$F_{n+1} = F_n \left(1 - \frac{4x^2}{(2n+1)^2}\right), \quad n = 2, 3, \dots \quad (10)$$

Using mathematical induction, we can prove the following

$$(1 + 4x^2)F_n > 1 + \frac{4x^2}{2n-1}, \quad n = 2, 3, \dots \quad (11)$$

In fact, for  $n = 2$ , we have

$$\begin{aligned} & (1 + 4x^2)F_2 - \left(1 + \frac{4}{3}x^2\right) \\ &= (1 + 4x^2) \left(1 - \frac{4}{9}x^2\right) - \left(1 + \frac{4}{3}x^2\right) \\ &= \frac{20}{9}x^2 - \frac{16}{9}x^4 \\ &> 0, \end{aligned} \quad (12)$$

that is

$$(1 + 4x^2)F_2 > 1 + \frac{4}{3}x^2. \quad (13)$$

Therefore, inequality (11) holds for  $n = 2$ .

Suppose inequality (11) holds for some  $m \geq 2$ , that is

$$(1 + 4x^2)F_m > 1 + \frac{4x^2}{2m-1}. \quad (14)$$

Then we have

$$\begin{aligned}
& (1 + 4x^2)F_{m+1} - \left(1 + \frac{4x^2}{2m+1}\right) \\
&= (1 + 4x^2)F_m \left(1 - \frac{4x^2}{(2m+1)^2}\right) - \left(1 + \frac{4x^2}{2m+1}\right) \\
&> \left(1 + \frac{4x^2}{2m-1}\right) \left(1 - \frac{4x^2}{(2m+1)^2}\right) - \left(1 + \frac{4x^2}{2m+1}\right) \\
&= \frac{4(2m+3-4x^2)x^2}{(2m-1)(2m+1)^2} \\
&> 0,
\end{aligned} \tag{15}$$

that is

$$(1 + 4x^2)F_{m+1} > 1 + \frac{4x^2}{2m+1}. \tag{16}$$

By induction, inequality (11) follows.

Further, since

$$\lim_{n \rightarrow \infty} (1 + 4x^2)F_n \geq 1, \tag{17}$$

combining (9) with (17) yields (4). The proof is complete.  $\square$

*Proof of inequality (5).* It suffices to prove inequality (5) holds for  $0 < x < \frac{1}{2}$ .

It is well-known [3, p. 193] that

$$\cosh(\pi x) = \prod_{n=1}^{\infty} \left(1 + \frac{4x^2}{(2n-1)^2}\right). \tag{18}$$

Let

$$Q_n = \prod_{k=2}^n \left(1 + \frac{4x^2}{(2k-1)^2}\right), \quad n = 2, 3, \dots \tag{19}$$

Then we have

$$\cosh(\pi x) = \frac{1 + 4x^2}{1 - 4x^2} \left[ (1 - 4x^2) \lim_{n \rightarrow \infty} Q_n \right], \tag{20}$$

and

$$Q_{n+1} = Q_n \left(1 + \frac{4x^2}{(2n+1)^2}\right), \quad n = 2, 3, \dots \tag{21}$$

Using mathematical induction, we can prove the following

$$(1 - 4x^2)Q_n < 1 - \frac{4x^2}{2n-1}, \quad n = 2, 3, \dots \tag{22}$$

In fact, for  $n = 2$ , we have

$$\begin{aligned}
& (1 - 4x^2)Q_2 - \left(1 - \frac{4}{3}x^2\right) \\
&= (1 - 4x^2) \left(1 + \frac{4}{9}x^2\right) - \left(1 - \frac{4}{3}x^2\right) \\
&= -\frac{20}{9}x^2 - \frac{16}{9}x^4 \\
&< 0,
\end{aligned} \tag{23}$$

that is,

$$(1 - 4x^2)Q_2 < 1 - \frac{4}{3}x^2. \quad (24)$$

Therefore, inequality (22) holds for  $n = 2$ .

Suppose inequality (22) holds for some  $m \geq 2$ , that is

$$(1 - 4x^2)Q_m < 1 - \frac{4x^2}{2m - 1}. \quad (25)$$

Then we have

$$\begin{aligned} & (1 - 4x^2)Q_{m+1} - \left(1 - \frac{4x^2}{2m + 1}\right) \\ &= (1 - 4x^2)Q_m \left(1 + \frac{4x^2}{(2m + 1)^2}\right) - \left(1 - \frac{4x^2}{2m + 1}\right) \\ &< \left(1 - \frac{4x^2}{2m - 1}\right) \left(1 + \frac{4x^2}{(2m + 1)^2}\right) - \left(1 - \frac{4x^2}{2m + 1}\right) \\ &= -4 \left(\frac{2}{(2m - 1)(2m + 1)} - \frac{1}{(2m + 1)^2}\right) x^2 - \frac{16x^2}{(2m - 1)(2m + 1)^2} \\ &< 0, \end{aligned} \quad (26)$$

that is

$$(1 - 4x^2)Q_{m+1} < 1 - \frac{4x^2}{2m + 1}. \quad (27)$$

By induction, inequality (22) follows.

It is easy to see that

$$\lim_{n \rightarrow \infty} (1 - 4x^2)Q_n \leq 1. \quad (28)$$

Combining (20) with (28) yields inequality (5). The proof is complete.  $\square$

*Proof of inequality (6).* It is sufficient to prove that inequality (6) holds for  $0 < x < 1$ .

It is well-known [3, p. 193] that

$$\frac{\sinh(\pi x)}{\pi x} = \prod_{n=1}^{\infty} \left(1 + \frac{x^2}{n^2}\right). \quad (29)$$

Setting

$$P_n = \prod_{k=2}^n \left(1 + \frac{x^2}{k^2}\right), \quad n = 2, 3, \dots, \quad (30)$$

then we have

$$\frac{\sinh(\pi x)}{\pi x} = \frac{1 + x^2}{1 - x^2} \left[ (1 - x^2) \lim_{n \rightarrow \infty} P_n \right], \quad (31)$$

and

$$P_{n+1} = P_n \left[ 1 + \frac{x^2}{(n+1)^2} \right], \quad n = 2, 3, \dots. \quad (32)$$

Using mathematical induction, it is easy to prove that

$$(1 - x^2)P_n < 1 - \frac{x^2}{n}, \quad n = 2, 3, \dots. \quad (33)$$

In fact, for  $n = 2$ , we have

$$\begin{aligned} (1-x^2)P_2 - \left(1 - \frac{x^2}{2}\right) &= (1-x^2) \left(1 + \frac{x^2}{4}\right) - \left(1 - \frac{x^2}{2}\right) \\ &= -\frac{x^2}{4} - \frac{x^4}{4} \\ &< 0, \end{aligned} \quad (34)$$

that is,

$$(1-x^2)P_2 < 1 - \frac{x^2}{2}. \quad (35)$$

Therefore, inequality (33) holds for  $n = 2$ .

Suppose inequality (33) holds for some  $m \geq 2$ , that is

$$(1-x^2)P_m < 1 - \frac{x^2}{m}. \quad (36)$$

Then we have

$$\begin{aligned} &(1-x^2)P_{m+1} - \left(1 - \frac{x^2}{m+1}\right) \\ &= (1-x^2)P_m \left(1 + \frac{x^2}{(m+1)^2}\right) - \left(1 - \frac{x^2}{m+1}\right) \\ &< \left(1 - \frac{x^2}{m}\right) \left(1 + \frac{x^2}{(m+1)^2}\right) - \left(1 - \frac{x^2}{m+1}\right) \\ &= -\left(\frac{1}{m(m+1)} - \frac{1}{(m+1)^2}\right)x^2 - \frac{x^4}{m(m+1)^2} \\ &< 0, \end{aligned} \quad (37)$$

that is,

$$(1-x^2)P_{m+1} < 1 - \frac{x^2}{m+1}. \quad (38)$$

By induction, inequality (33) holds.

Further, since

$$\lim_{n \rightarrow \infty} (1-x^2)P_n \leq 1, \quad (39)$$

from (31), inequality (6) follows.  $\square$

## REFERENCES

- [1] Ch.-P. Chen and F. Qi, *A double inequality for remainder of power series of tangent function*, Tamkang J. Math. **34** (2003), no. 3, accepted. RGMIA Res. Rep. Coll. **5** (2002), suppl., Art. 2. Available online at [http://rgmia.vu.edu.au/v5\(E\).html](http://rgmia.vu.edu.au/v5(E).html).
- [2] Ch.-P. Chen and F. Qi, *Inequalities of some trigonometric functions*, RGMIA Res. Rep. Coll. **6** (2003). Available online at <http://rgmia.vu.edu.au>.
- [3] Group of compilation, *Shùxué Shǒucè (Handbook of Mathematics)*, The People's Education Press, Beijing, China, 1979. (Chinese)
- [4] G. Klambauer, *Problems and Propositions in Analysis*, Marcel Dekker, New York and Basel, 1979.
- [5] H. Kober, *Approximation by integral functions in the complex domain*, Trans. Amer. Math. Soc. **56** (1944), 7–31.
- [6] J.-Ch. Kuang, *Chángyòng Bùděngshì (Applied Inequalities)*, 2nd ed., Hunan Education Press, Changsha, China, 1993. (Chinese)
- [7] A. McD. Mercer, Ulrich and Donald Caccia, *A sharpening of Jordan's inequality*, Amer. Math. Monthly **93** (1986), 568–569.

- [8] D. S. Mitrinović, Translated by X.-P. Zhang and L. Wang, *Jiěxī Bùdēngshì (Analytic Inequalities)*, Chinese Ed., Science Press, Beijing, China, 1987. English Ed., Springer, New York, 1970.
- [9] F. Qi, *Extensions and sharpenings of Jordan's and Kober's inequality*, Gōngkē Shùxué (J. Math. Tech.) **12** (1996), no. 4, 98–102. (Chinese)
- [10] F. Qi, L.-H. Cui and S.-L. Xu, *Some inequalities constructed by Tchebysheff's integral inequality*, Math. Inequal. Appl. **2** (1999), no. 4, 517–528.
- [11] F. Qi and Q.-D. Hao, *Refinements and sharpenings of Jordan's and Kober's inequality*, Math. Inform. Quart. **8** (1998), no. 3, 116–120.
- [12] R. Redheffer, *Problem 5642*, Amer. Math. Monthly **76** (1969), 422.

(Ch.-P. Chen) DEPARTMENT OF APPLIED MATHEMATICS AND INFORMATICS, JIAOZUO INSTITUTE OF TECHNOLOGY, JIAOZUO CITY, HENAN 454000, CHINA

(J.-W. Zhao) INFORMATION CENTER, EDUCATION DEPARTMENT OF HENAN PROVINCE, ZHENGZHOU CITY, HENAN 450003, CHINA

(F. Qi) DEPARTMENT OF APPLIED MATHEMATICS AND INFORMATICS, JIAOZUO INSTITUTE OF TECHNOLOGY, JIAOZUO CITY, HENAN 454000, CHINA

*E-mail address:* qifeng@jzit.edu.cn, fengqi618@member.ams.org

*URL:* <http://rgmia.vu.edu.au/qi.html>