



VICTORIA UNIVERSITY
MELBOURNE AUSTRALIA

Self Sharpening Inequality

This is the Published version of the following publication

Hassani, Mehdi (2003) Self Sharpening Inequality. RGMIA research report collection, 6 (3).

The publisher's official version can be found at

Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/17852/>

SELF SHARPENING INEQUALITY

MEHDI HASSANI

ABSTRACT. In this note we analyze the following inequality that is observed in the study of the density of n^{th} -power free integers

$$\left| \frac{x}{\zeta(n)} - f_n(x) \right| < \frac{\sqrt[n]{x}}{n-1} + \left(\frac{\sqrt[n]{x}}{\zeta(2)} - 1 \right) + \left(1 + \frac{1}{\zeta(2)} \right) \sum_{i=1}^M x^{\frac{1}{n2^i}} + 2x^{\frac{1}{n2^{M+1}}} \quad (M \in \mathbb{N}),$$

in which $f_n(x)$ is the number of n^{th} -power free positive integers $\leq x$. We show that the above inequality is *self sharpening* and then we compute the order of its sharpening.

1. INTRODUCTION AND MOTIVATION

Let \mathbb{P} be the set of all primes and suppose N is a positive integer, with the following prime factoring:

$$N = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_k^{\alpha_k} \quad (p_1, p_2, \dots, p_k \in \mathbb{P}).$$

We say that N , is n^{th} -power free if all α_i 's are less than n . Let $f_n(x)$ be the number of n^{th} -power free integers $\leq x$. It is well-known that the density of n^{th} -power free integers is $\frac{1}{\zeta(n)}$, or equivalently

$$f_n(x) \sim \frac{x}{\zeta(n)} \quad (x \rightarrow \infty),$$

such that $\zeta(s) = \sum_{m=1}^{\infty} \frac{1}{m^s}$. In fact, we have

$$f_n(x) = \frac{x}{\zeta(n)} + O(\sqrt[n]{x}),$$

and the stronger result

$$(1.1) \quad \left| \frac{x}{\zeta(n)} - f_n(x) \right| < \frac{n}{n-1} \sqrt[n]{x} - 1.$$

In particular,

$$f_2(x) < \frac{x}{\zeta(2)} + \sqrt{x} - 1.$$

As you will see soon, during the proof of (1.1) we use $\left| \frac{x}{\zeta(n)} - f_n(x) \right| < \sum_{1 < k \leq \sqrt[n]{x}} |\mu(k)| + \frac{\sqrt[n]{x}}{n-1}$ and $\sum_{1 < k \leq \sqrt[n]{x}} |\mu(k)| < \sqrt[n]{x} - 1$, in which μ is the well-known *Mobius Function* and defined by

$$\mu(m) = \begin{cases} 1 & m = 1; \\ (-1)^k & m = p_1 p_2 \cdots p_k; \\ 0 & \text{otherwise.} \end{cases}$$

1991 *Mathematics Subject Classification*. Primary 39B72, 26D15, 26D20.

Key words and phrases. Self sharpening inequality, Square free integer, n^{th} -power free integer, Density.

Since the function $\mu(k)$ vanishes if k is not square free, we can claim that

$$\sum_{1 < k \leq \sqrt[n]{x}} |\mu(k)| \leq f_2(\sqrt[n]{x}) < \frac{\sqrt[n]{x}}{\zeta(2)} + {}^2\sqrt[n]{x} - 1.$$

The obtained bound for $\sum_{1 < k \leq \sqrt[n]{x}} |\mu(k)|$ is sharper than $\sqrt[n]{x} - 1$. Thus, (1.1) becomes sharper as follows:

$$(1.2) \quad \left| \frac{x}{\zeta(n)} - f_n(x) \right| < \frac{\sqrt[n]{x}}{n-1} + \frac{\sqrt[n]{x}}{\zeta(2)} + 2 {}^2\sqrt[n]{x} - 1.$$

By using (1.2) we have

$$f_2(x) < \frac{x}{\zeta(2)} + \left(1 + \frac{1}{\zeta(2)}\right) \sqrt{x} + 2 {}^4\sqrt{x} - 1,$$

and repeating above method we obtain,

$$\left| \frac{x}{\zeta(n)} - f_n(x) \right| < \frac{\sqrt[n]{x}}{n-1} + \frac{\sqrt[n]{x}}{\zeta(2)} + \left(1 + \frac{1}{\zeta(2)}\right) {}^2\sqrt[n]{x} + 2 {}^4\sqrt[n]{x} - 1.$$

This process leads us to the following definition:

Definition 1 (Self Sharpening Inequality). *We say that an inequality is self sharpening, if it sharp itself!*

In the next section we continue above method to show that (1.1) is self sharpening.

2. PROOFS AND SELF SHARPENING INEQUALITY

For a proof of (1.1), we note that

$$\sum_{m=1}^{\infty} \frac{\mu(m)}{m^s} = \frac{1}{\zeta(s)}.$$

Now, by a counting we have

$$f_n(x) = x - \sum_{p \in \mathbb{P}} \left\lfloor \frac{x}{p^n} \right\rfloor + \sum_{p, q \in \mathbb{P}, p \neq q} \left\lfloor \frac{x}{(pq)^n} \right\rfloor - \dots = \sum_{k \leq \sqrt[n]{x}} \mu(k) \left\lfloor \frac{x}{k^n} \right\rfloor.$$

and so,

$$\begin{aligned} \left| \frac{x}{\zeta(n)} - f_n(x) \right| &= \left| \sum_{1 < k \leq \sqrt[n]{x}} \mu(k) \left(\frac{x}{k^n} - \left\lfloor \frac{x}{k^n} \right\rfloor \right) + \sum_{k > \sqrt[n]{x}} \mu(k) \frac{x}{k^n} \right| \\ &< \sum_{1 < k \leq \sqrt[n]{x}} |\mu(k)| + x \sum_{k > \sqrt[n]{x}} \frac{1}{k^n} \\ &< \sqrt[n]{x} - 1 + x \int_{\sqrt[n]{x}}^{\infty} \frac{ds}{s^n} = \frac{n}{n-1} \sqrt[n]{x} - 1. \end{aligned}$$

Now, we show that (1.1) is self sharpening.

Theorem 1. *For all $M \in \mathbb{N}$ the inequality*

$$(2.1) \quad \left| \frac{x}{\zeta(n)} - f_n(x) \right| < \frac{\sqrt[n]{x}}{n-1} + \left(\frac{\sqrt[n]{x}}{\zeta(2)} - 1 \right) + \left(1 + \frac{1}{\zeta(2)} \right) \sum_{i=1}^M x^{\frac{1}{n2^i}} + 2x^{\frac{1}{n2^{M+1}}},$$

is self sharpening and the order of its sharpening is

$$O\left(x^{\frac{1}{n2^{M+1}}}\right).$$

Proof. The inequality (2.1) yields by induction on M and using (1.1). Now, let $U(M)$ be the right side of (2.1);

$$U(M) = \frac{\sqrt[n]{x}}{n-1} + \left(\frac{\sqrt[n]{x}}{\zeta(2)} - 1\right) + \left(1 + \frac{1}{\zeta(2)}\right) \sum_{i=1}^M x^{\frac{1}{n2^i}} + 2x^{\frac{1}{n2^{M+1}}}.$$

Now, we have

$$(2.2) \quad \Delta U(M) := U(M) - U(M+1) = \left(1 - \frac{1}{\zeta(2)}\right) x^{\frac{1}{n2^{M+1}}} - 2x^{\frac{1}{n2^{M+2}}}.$$

For $x > \left(\frac{2}{1-\frac{1}{\zeta(2)}}\right)^{n2^{M+1}}$ we easily have $\Delta U(M) > 0$; which means that (2.1) is self sharpening. Also, according to (2.2), we have

$$\Delta U(M) = O\left(x^{\frac{1}{n2^{M+1}}}\right),$$

which is the order of sharpening. \square

As you see, the idea of sharpening (1.1) and (2.1) comes from noting the definition of the Mobius function and applying them for $n = 2$.

3. ACKNOWLEDGEMENTS

I would like to express my gratitude to Dr J. Roojin for his valuable comments.

REFERENCES

- [1] Razvan, Hassani and Pirayesh, *DENSITY OF n^{th} -POWER FREES*, Submitted.

DEPARTMENT OF MATHEMATICS, INSTITUTE FOR ADVANCED STUDIES IN BASIC SCIENCES, ZANJAN, IRAN
E-mail address: mhassani@iasbs.ac.ir