



**VICTORIA UNIVERSITY**  
MELBOURNE AUSTRALIA

*Monotonicity and convexity of the function  $x\sqrt{\Gamma(x+1)}/(x+\alpha)\sqrt{\Gamma(x+\alpha+1)}$*

This is the Published version of the following publication

Qi, Feng and Guo, Bai-Ni (2003) Monotonicity and convexity of the function  $x\sqrt{\Gamma(x+1)}/(x+\alpha)\sqrt{\Gamma(x+\alpha+1)}$ . RGMIA research report collection, 6 (4).

The publisher's official version can be found at

Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/17868/>

# MONOTONICITY AND CONVEXITY OF THE FUNCTION

$$\sqrt[x]{\Gamma(x+1)} / \sqrt[x+\alpha]{\Gamma(x+\alpha+1)}$$

FENG QI AND BAI-NI GUO

ABSTRACT. For  $\alpha > 0$  a real number, the function  $\frac{\sqrt[x]{\Gamma(x+1)}}{x+\sqrt[x+\alpha]{\Gamma(x+\alpha+1)}}$  is increasing with  $x \in (x_0, \infty)$  and logarithmically concave with  $x \in [1, \infty)$ , where  $x_0 \in (0, 1)$  is a constant. Moreover, some monotonicity and convexity results and inequalities of functions involving the gamma function and polygamma functions are obtained as corollaries and by-products.

## 1. INTRODUCTION

In this section, we first state some known results: monotonicity of the geometric mean sequence and some sequences involving geometric means, inequalities of ratio between geometric means and ratio between power means, and monotonicity and convexity of ratio between two gamma functions and functions involving the gamma function.

**1.1. Inequalities of ratio between power means.** H. Minc and L. Sathre in [27] gave the lower and upper bounds of ratio between two geometric means of natural numbers:

$$\frac{n-1}{n} < \frac{\sqrt[n-1]{(n-1)!}}{\sqrt[n]{n!}} < 1. \quad (1)$$

The right hand side inequality in (1) also reveals that the geometric mean sequence  $\{\sqrt[n]{n!}\}_{n \in \mathbb{N}}$  is strictly increasing and the sequence  $\{\frac{\sqrt[n]{n!}}{n}\}_{n \in \mathbb{N}}$  is strictly decreasing. Note that  $\Gamma(n+1) = n!$ , where the gamma function is usually defined [19, 54] for  $\text{Re } z > 0$  by

$$\Gamma(z) = \int_0^\infty t^{z-1} e^{-t} dt. \quad (2)$$

In [2, 26], H. Alzer and J. S. Martins refined the left hand side inequality in (1) and showed that, if  $n$  is a positive integer, then, for all positive real numbers  $r$ ,

$$\frac{n}{n+1} < \left( \frac{\frac{1}{n} \sum_{i=1}^n i^r}{\frac{1}{n+1} \sum_{i=1}^{n+1} i^r} \right)^{1/r} < \frac{\sqrt[n]{n!}}{\sqrt[n+1]{(n+1)!}}. \quad (3)$$

---

2000 *Mathematics Subject Classification.* Primary 33B15, 33B20; Secondary 26D07, 26D15, 26D20, 60E05.

*Key words and phrases.* gamma function, polygamma function, monotonicity, convexity, mean value, inequality, ratio.

The authors were supported in part by NNSF (#10001016) of China, SF for the Prominent Youth of Henan Province (#0112000200), SF of Henan Innovation Talents at Universities, Doctor Fund of Jiaozuo Institute of Technology, CHINA.

This paper was typeset using  $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{L}\mathcal{A}\mathcal{T}\mathcal{E}\mathcal{X}$ .

Both bounds in (3) are the best possible. The middle term in (3) is indeed a ratio between the power means  $(\frac{1}{n} \sum_{i=1}^n i^r)^{1/r}$  and  $(\frac{1}{n+1} \sum_{i=1}^{n+1} i^r)^{1/r}$ . The inequality (3) implies that the sequence  $\{\frac{1}{n}(\frac{1}{n} \sum_{i=1}^n i^r)^{1/r}\}_{n \in \mathbb{N}}$  is decreasing strictly and the sequence  $\{\frac{1}{\sqrt[n]{n!}}(\frac{1}{n} \sum_{i=1}^n i^r)^{1/r}\}_{n \in \mathbb{N}}$  is increasing strictly for given  $r > 0$ .

The integral form of inequality (3) was established in [29, 41] by the authors: Let  $b > a > 0$  and  $\delta > 0$  be real numbers. Then, for any positive  $r \in \mathbb{R}$ , we have

$$\begin{aligned} \frac{b}{b+\delta} &< \left( \frac{b+\delta-a}{b-a} \cdot \frac{b^{r+1}-a^{r+1}}{(b+\delta)^{r+1}-a^{r+1}} \right)^{1/r} \\ &= \left( \frac{\frac{1}{b-a} \int_a^b x^r dx}{\frac{1}{b+\delta-a} \int_a^{b+\delta} x^r dx} \right)^{1/r} < \frac{[b^b/a^a]^{1/(b-a)}}{[(b+\delta)^{b+\delta}/a^a]^{1/(b+\delta-a)}}. \end{aligned} \quad (4)$$

The lower and upper bounds in (4) are the best possible. The inequality (4) can be restated as monotonicity results: The function  $\frac{1}{x} \left( \frac{x^{r+1}-a^{r+1}}{x-a} \right)^{1/r}$  is decreasing and the function  $\frac{1}{[x^x/a^a]^{1/(x-a)}} \left( \frac{x^{r+1}-a^{r+1}}{x-a} \right)^{1/r}$  is increasing with  $x > 0$  for given  $r > 0$ . Notice that  $\frac{1}{e} \left[ \frac{x^x}{a^a} \right]^{1/(x-a)}$  is called the identric or exponential mean.

After obtaining the following generalization of the left hand side inequality of (3):

$$\frac{n+k}{n+m+k} < \left( \frac{\frac{1}{n} \sum_{i=k+1}^{n+k} i^r}{\frac{1}{n+m} \sum_{i=k+1}^{n+m+k} i^r} \right)^{1/r}, \quad (5)$$

where  $r$  is a given positive real number,  $n$  and  $m$  are natural numbers and  $k$  is a nonnegative integer, the first author in [31] asked as an open problem the validity of an inequality below:

$$\left( \frac{\frac{1}{n} \sum_{i=1}^n a_i^r}{\frac{1}{n+m} \sum_{i=1}^{n+m} a_i^r} \right)^{1/r} < \frac{\sqrt[n]{a_n!}}{n+m \sqrt[n+m]{a_{n+m!}}}, \quad (6)$$

where  $r$  is a positive number,  $a_n!$  denotes the sequence factorial defined by  $\prod_{i=1}^n a_i$ . The upper bound in (6) is the best possible. Inequality (5) means that the sequence  $\{\frac{1}{n+k}(\frac{1}{n} \sum_{i=k+1}^{n+k} i^r)^{1/r}\}_{n \in \mathbb{N}}$  is decreasing for given  $r > 0$  and nonnegative integer  $k$ .

Inequality (6) has been researched in [3, 36, 55], some sufficient conditions are found. The first author in [36] obtained: Let  $n, m \in \mathbb{N}$  and  $\{a_i\}_{i=1}^{n+m}$  be an increasing, logarithmically concave, positive, and nonconstant sequence such that the sequence  $\{i[\frac{a_{i+1}}{a_i} - 1]\}_{i=1}^{n+m-1}$  is increasing, then inequality (6) holds. In particular, let  $a$  be a positive real numbers,  $b$  a nonnegative real number,  $k$  a nonnegative integer, and  $m, n \in \mathbb{N}$ , then, for any real number  $r > 0$ , we have

$$\left( \frac{\frac{1}{n} \sum_{i=k+1}^{n+k} (ai+b)^r}{\frac{1}{n+m} \sum_{i=k+1}^{n+m+k} (ai+b)^r} \right)^{1/r} < \frac{\sqrt[n]{\prod_{i=k+1}^{n+k} (ai+b)}}{n+m \sqrt[n+m]{\prod_{i=k+1}^{n+m+k} (ai+b)}}. \quad (7)$$

The authors and a coworker in [47] give a lower bound for ratio between two power means: Let  $n$  and  $m$  be natural numbers, suppose  $\{a_i\}_{i=1}^{n+m}$  is an increasing,

logarithmically convex, and positive sequence, then

$$\left( \frac{\frac{1}{n} \sum_{i=1}^n a_i^r}{\frac{1}{n+m} \sum_{i=1}^{n+m} a_i^r} \right)^{1/r} \geq \frac{a_n}{a_{n+m}}. \quad (8)$$

The lower bound in (8) is the best possible.

*Remark 1.* Indeed, the inequalities (3) to (7) are also valid for negative power  $r$ . For more information, please refer to [4, 5] and some unpublished papers.

**1.2. Inequalities of ratio between geometric means.** The inequalities in (1) were also refined and generalized in [33, 45, 48] and the following inequalities were obtained:

$$\frac{n+k+1}{n+m+k+1} < \frac{(\prod_{i=k+1}^{n+k} i)^{1/n}}{(\prod_{i=k+1}^{n+m+k} i)^{1/(n+m)}} \leq \sqrt{\frac{n+k}{n+m+k}}, \quad (9)$$

where  $k$  is a nonnegative integer,  $n$  and  $m$  are natural numbers. For  $n = m = 1$ , the equality in (9) is valid.

In [15, 34], inequalities in (9) were generalized and obtained the following inequalities on the ratio for the geometric means of a positive arithmetic sequence:

$$\frac{a(n+k+1)+b}{a(n+m+k+1)+b} \leq \frac{[\prod_{i=k+1}^{n+k} (ai+b)]^{\frac{1}{n}}}{[\prod_{i=k+1}^{n+m+k} (ai+b)]^{\frac{1}{n+m}}} \leq \sqrt{\frac{a(n+k)+b}{a(n+m+k)+b}}, \quad (10)$$

where  $a$  and  $b$  are positive constants,  $k$  is a nonnegative integer,  $n$  and  $m$  are natural numbers.

In [43, 44], the following general monotonicity properties are established: Let  $f$  be a positive function defined on  $[1, \infty)$  such that  $\frac{f(x+2)}{f(x+1)} \geq \frac{x+2}{x+1} \left[ \frac{x(x+2)}{(x+1)^2} \right]^{\frac{x}{2}}$  for  $x \geq 0$ , then the sequence  $\left\{ \frac{\sqrt[n]{\prod_{i=1}^n f(i)}}{\sqrt{n}} \right\}_{n \in \mathbb{N}}$  is increasing; if  $\frac{f(x+2)}{f(x+1)} \leq \left( \frac{x+3}{x+2} \right)^2 \left[ \frac{(x+1)(x+3)}{(x+2)^2} \right]^x$  holds on  $[0, \infty)$ , then the sequence  $\left\{ \frac{\sqrt[n]{\prod_{i=1}^n f(i)}}{(n+1)} \right\}_{n \in \mathbb{N}}$  is decreasing. Let  $f$  be a positive function such that  $x \left[ \frac{f(x+1)}{f(x)} - 1 \right]$  is increasing on  $[1, \infty)$ , then the sequence  $\left\{ \frac{\sqrt[n]{\prod_{i=1}^n f(i)}}{f(n+1)} \right\}_{n=1}^{\infty}$  is decreasing; if  $f$  is a logarithmically concave and positive function defined on  $[1, \infty)$ , then the sequence  $\left\{ \frac{\sqrt[n]{\prod_{i=1}^n f(i)}}{\sqrt{f(n)}} \right\}_{n=1}^{\infty}$  is increasing. As consequences of these monotonicities, the lower and upper bounds for the ratio  $\frac{\sqrt[n]{\prod_{i=k+1}^{n+k} f(i)}}{\sqrt[n+m]{\prod_{i=k+1}^{n+m+k} f(i)}}$  are obtained, where  $k$  is a nonnegative integer and  $m$  a natural number.

As lemmas in [36], the following results were showed: Let  $n, m \in \mathbb{N}$ , and  $\{a_i\}_{i=1}^{n+m+1}$  a nonconstant positive sequence such that the sequence  $\left\{ i \left[ \frac{a_{i+1}}{a_i} - 1 \right] \right\}_{i=1}^{n+m}$  is increasing, then the sequence  $\left\{ \frac{\sqrt[i]{a_i!}}{a_{i+1}} \right\}_{i=1}^{n+m}$  is decreasing, and then  $\frac{\sqrt[n]{a_n!}}{n+m \sqrt[n+m]{a_{n+m}!}} > \frac{a_{n+1}}{a_{n+m+1}}$ . Let  $n > 1$  be a positive integer and  $\{a_i\}_{i=1}^n$  an increasing nonconstant positive sequence such that  $\left\{ i \left[ \frac{a_{i+1}}{a_i} - 1 \right] \right\}_{i=1}^{n-1}$  is increasing, then the sequence

$\left\{\frac{a_i}{(a_i!)^{1/i}}\right\}_{i=1}^n$  is increasing, and then  $\frac{a_\ell}{a_n} < \frac{\sqrt[\ell]{a_\ell!}}{\sqrt[n]{a_n!}}$  for any positive integer  $\ell$  satisfying  $1 \leq \ell < n$ , where  $a_n!$  denotes the sequence factorial  $\prod_{i=1}^n a_i$ .

In [42] and a subsequent paper [7], some inequalities for ratios of geometric means of positive sequence are obtained as applications: If  $\{a_i\}_{i \in \mathbb{N}}$  is an increasing, positive sequence such that  $\left\{i\left(\frac{a_{i+1}}{a_i} - 1\right)\right\}_{i \in \mathbb{N}}$  increases, then we have

$$\frac{a_n}{a_{n+1}} \leq \frac{\sqrt[n]{\prod_{i=1}^n (a_i + a_n)}}{\sqrt[n+1]{\prod_{i=1}^{n+1} (a_i + a_{n+1})}} \leq \frac{\sqrt[n]{\prod_{i=1}^n a_i}}{\sqrt[n+1]{\prod_{i=1}^{n+1} a_i}}. \quad (11)$$

If  $\varphi$  is increasing, convex, positive and defined on  $(0, \infty)$  with  $\left\{\varphi(i)\left[\frac{\varphi(i)}{\varphi(i+1)} - 1\right]\right\}_{i \in \mathbb{N}}$  decreases, then

$$\frac{[\varphi(n)]^{n/\varphi(n)}}{[\varphi(n+1)]^{(n+1)/\varphi(n+1)}} \leq \frac{\sqrt[\varphi(n)]{\prod_{i=1}^n [\varphi(i) + \varphi(n)]}}{\sqrt[\varphi(n+1)]{\prod_{i=1}^{n+1} [\varphi(i) + \varphi(n+1)]}}. \quad (12)$$

There are much literature devoted to research of ratios of mean values, for example [50]. For more detailed information, please refer to references in this paper and references therein.

**1.3. Monotonicity and convexity of functions involving gamma functions and ratio of gamma functions.** It is well-known that the incomplete gamma function is defined and denoted for  $\operatorname{Re} z > 0$  by

$$\Gamma(z, x) = \int_x^\infty t^{z-1} e^{-t} dt, \quad \gamma(z, x) = \int_0^x t^{z-1} e^{-t} dt, \quad (13)$$

with  $\Gamma(z, 0) = \Gamma(z)$  and  $\Gamma(0, x) = E_1(x)$  is called the exponential integral.

In [18], the following monotonicity results for the gamma function were established: The function  $[\Gamma(1 + \frac{1}{x})]^x$  decreases with  $x > 0$  and  $x[\Gamma(1 + \frac{1}{x})]^x$  increases with  $x > 0$ , which recover the inequalities (1), which refer to integer values of  $n$ .

These are equivalent to the function  $[\Gamma(1+x)]^{\frac{1}{x}}$  being increasing and  $\frac{[\Gamma(1+x)]^{\frac{1}{x}}}{x}$  being decreasing on  $(0, \infty)$ , respectively. In addition, it was proved that the function  $x^{1-\gamma}[\Gamma(1 + \frac{1}{x})]^x$  decreases for  $0 < x < 1$ , where  $\gamma = 0.57721566 \dots$  denotes the Euler-Mascheroni constant, which is equivalent to  $\frac{[\Gamma(1+x)]^{\frac{1}{x}}}{x^{1-\gamma}}$  being increasing on  $(1, \infty)$ .

In [6, 39], it is proved that the function  $f(x) = \frac{[\Gamma(x+1)]^{1/x}}{x+1}$  is strictly decreasing and strictly logarithmically convex in  $(0, \infty)$  and the function  $g(x) = \frac{[\Gamma(x+1)]^{1/x}}{\sqrt{x+1}}$  is strictly increasing and strictly logarithmically concave in  $(0, \infty)$ . Moreover, if  $s$  is a positive real number, then for all real numbers  $x > 0$ ,

$$\frac{e^{-\gamma}}{[\Gamma(s+1)]^{1/s}} < \frac{[\Gamma(x+1)]^{1/x}}{[\Gamma(x+s+1)]^{1/(x+s)}} < 1, \quad (14)$$

$\lim_{x \rightarrow 0} f(x) = e^{-\gamma}$  and  $\lim_{x \rightarrow \infty} f(x) = e^{-1}$ .

Using monotonicity properties and inequalities of the generalized weighted mean values (see [13, 30, 32, 37, 51]), the first author proved [35] that the functions  $\left[\frac{\Gamma(s)}{\Gamma(r)}\right]^{1/(s-r)}$ ,  $\left[\frac{\Gamma(s,x)}{\Gamma(r,x)}\right]^{1/(s-r)}$  and  $\left[\frac{\gamma(s,x)}{\gamma(r,x)}\right]^{1/(s-r)}$  are increasing in  $r > 0$ ,  $s > 0$  and  $x > 0$ ; for any given  $x > 0$ , the function  $\frac{s\gamma(s,x)}{x^s}$  is decreasing in  $s > 0$ . These generalize and extend the related results in [10, 11, 17, 18, 27] for the range of the argument. For more inequalities of quotients between gamma functions can be found in [16], [28, p. 526] and [52, pp. 442–443].

Using the approach by A. Laforgia and S. Sismondi in [20], some more general inequalities of the functions  $\int_0^x e^{pt} dt$  and  $\int_0^x e^{-pt} dt = \frac{\Gamma(1/p) - \Gamma(1/p, x^p)}{p}$  for  $p > 0$  and  $x > 0$  are obtained in [49]. These two functions are also been investigated by utilizing Tchebysheff integral inequality and Hermite-Hadamard integral inequality in [40, 46] by the first author and coworkers. For more information, please refer to [8].

In [9], Elezović, Giordana and Pečarić, among others, verified the convexity with respect to variable  $x$  of the function  $\left[\frac{\Gamma(x+t)}{\Gamma(x+s)}\right]^{1/(t-s)}$  for  $|t-s| < 1$ , obtained the best bounds for  $\frac{\Gamma(x+1)}{\Gamma(x+s)}$ , where  $s \in (0, 1)$  and  $x \geq 1$ , given some different approach from Gautschi's in [11], proved several new simple inequalities for digamma function, and improved related results by D. Kershaw in [17].

In [53], it is shown that the function  $1 + \frac{1}{x} \ln \Gamma(x+1) - \ln(x+1)$  is strictly completely monotone on  $(-1, \infty)$  and tends to 1 as  $x \rightarrow -1$  and to 0 as  $x \rightarrow \infty$ .

In [14], the following monotonicity result was obtained: The function

$$\frac{[\Gamma(x+y+1)/\Gamma(y+1)]^{1/x}}{x+y+1} \quad (15)$$

is decreasing in  $x \geq 1$  for fixed  $y \geq 0$ . Then, for positive real numbers  $x$  and  $y$ , we have

$$\frac{x+y+1}{x+y+2} \leq \frac{[\Gamma(x+y+1)/\Gamma(y+1)]^{1/x}}{[\Gamma(x+y+2)/\Gamma(y+1)]^{1/(x+1)}}. \quad (16)$$

Inequality (16) extends and generalizes inequality (9), since  $\Gamma(n+1) = n!$ .

In this article, we are about to prove monotonicity and convexity properties of ratio between  $\sqrt[x]{\Gamma(x+1)}$  and  $\sqrt[x+\alpha]{\Gamma(x+\alpha+1)}$  which are generalizations of the geometric means. Our main results are as follows.

**Theorem 1.** For  $\alpha > 0$  a real number, the function  $\frac{\sqrt[x]{\Gamma(x+1)}}{\sqrt[x+\alpha]{\Gamma(x+\alpha+1)}}$  is increasing with  $x \in (x_0, \infty)$ , where  $x_0 \in (0, 1)$  is a constant.

**Theorem 2.** For  $\alpha > 0$  a real number, the function  $\frac{\sqrt[x]{\Gamma(x+1)}}{\sqrt[x+\alpha]{\Gamma(x+\alpha+1)}}$  is logarithmically concave with  $x \in [1, \infty)$ .

*Remark 2.* Basing on the graph of  $\frac{\sqrt[x]{\Gamma(x+1)}}{\sqrt[x+\alpha]{\Gamma(x+\alpha+1)}}$  pictured by Mathematica, we conjecture that the function  $\frac{\sqrt[x]{\Gamma(x+1)}}{\sqrt[x+\alpha]{\Gamma(x+\alpha+1)}}$  is increasing and logarithmically concave with  $x \in (-1, \infty)$  for a given  $\alpha > 0$ .

## 2. LEMMA

It is well known that the Bernoulli numbers  $B_n$  is defined ([1] and [54, p. 1]) in general by

$$\frac{1}{e^t - 1} + \frac{1}{2} - \frac{1}{t} = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{t^{2n}}{(2n)!} B_n. \quad (17)$$

In particular, we have the following

$$B_1 = \frac{1}{6}, \quad B_2 = \frac{1}{30}, \quad B_3 = \frac{1}{42}, \quad B_4 = \frac{1}{30}. \quad (18)$$

In [54, p. 45], the following summation formula is given

$$\sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)^{2k+1}} = \frac{\pi^{2k+1} E_k}{2^{2k+2} (2k)!} \quad (19)$$

for nonnegative integer  $k$ , where  $E_k$  denotes the Euler number, which implies

$$B_n = \frac{2(2n)!}{(2\pi)^{2n}} \sum_{m=1}^{\infty} \frac{1}{m^{2n}}, \quad n \in \mathbb{N}. \quad (20)$$

*Remark 3.* Recently, the Bernoulli and Euler numbers and polynomials are generalized in [12, 21, 22, 23, 24, 25] and some unpublished papers by the authors and coworkers.

**Lemma 1.** *For real number  $x > 0$  and natural number  $m$ , we have*

$$\begin{aligned} \ln \Gamma(x) &= \frac{1}{2} \ln(2\pi) + \left(x - \frac{1}{2}\right) \ln x - x + \sum_{n=1}^m (-1)^{n-1} \frac{B_n}{2(2n-1)n} \cdot \frac{1}{x^{2n-1}} \\ &\quad + (-1)^m \theta_1 \cdot \frac{B_{m+1}}{(2m+1)(2m+2)} \cdot \frac{1}{x^{2m+1}}, \quad 0 < \theta_1 < 1; \end{aligned} \quad (21)$$

$$\begin{aligned} \psi(x) &= \ln x - \frac{1}{2x} + \sum_{n=1}^m (-1)^n \frac{B_n}{2n} \cdot \frac{1}{x^{2n}} + (-1)^{m+1} \theta_2 \cdot \frac{B_{m+1}}{2m+2} \cdot \frac{1}{x^{2m+2}}; \\ 0 < \theta_2 < 1, \end{aligned} \quad (22)$$

$$\psi'(x) = \frac{1}{x} + \frac{1}{2x^2} + \sum_{n=1}^m (-1)^{n-1} \frac{B_n}{x^{2n+1}} + (-1)^m \theta_3 \cdot \frac{B_{m+1}}{x^{2m+3}}, \quad 0 < \theta_3 < 1; \quad (23)$$

$$\begin{aligned} \psi''(x) &= -\frac{1}{x^2} - \frac{1}{x^3} + \sum_{n=1}^m (-1)^n (2n+1) \frac{B_n}{x^{2n+3}} \\ &\quad + (-1)^{m+1} (2m+3) \theta_4 \cdot \frac{B_{m+1}}{x^{2m+4}}, \quad 0 < \theta_4 < 1. \end{aligned} \quad (24)$$

*Proof.* Let  $x > 0$ , then we have

$$\begin{aligned} \ln \Gamma(x) &= \frac{1}{2} \ln(2\pi) + \left(x - \frac{1}{2}\right) \ln x - x - \int_{-\infty}^0 \left(\frac{1}{2} - \frac{1}{t} - \frac{1}{1-e^t}\right) \frac{e^{xt}}{t} dt \\ &\triangleq \frac{1}{2} \ln(2\pi) + \left(x - \frac{1}{2}\right) \ln x - x - \omega(x), \end{aligned} \quad (25)$$

which is called the first Binet's formula. See [1] and [54, p. 106].

It is well-known that

$$\coth x = \frac{1}{x} + \sum_{k=1}^{\infty} \frac{2x}{x^2 + k^2\pi^2}$$

for  $x \neq 0$ , and

$$\frac{x}{e^x - 1} + \frac{x}{2} = \frac{x}{2} \coth \frac{x}{2},$$

therefore

$$\frac{1}{x} \left( \frac{1}{e^x - 1} - \frac{1}{x} + \frac{1}{2} \right) = 2 \sum_{k=1}^{\infty} \frac{1}{x^2 + 4\pi^2 k^2}. \quad (26)$$

For any given natural number  $m$ , we have

$$\frac{1}{x^2 + 4\pi^2 k^2} = \sum_{n=1}^m (-1)^{n-1} \frac{x^{2(n-1)}}{(4\pi^2 k^2)^n} + (-1)^m \frac{x^{2m}}{(4\pi^2 k^2)^{m+1}} \cdot \frac{1}{1 + \frac{x^2}{4\pi^2 k^2}}, \quad (27)$$

for  $1 \leq n \leq m$ , we have

$$\sum_{k=1}^{\infty} (-1)^{n-1} \frac{x^{2(n-1)}}{(4\pi^2 k^2)^n} = (-1)^{n-1} \frac{x^{2(n-1)}}{(2\pi)^{2n}} \sum_{k=1}^{\infty} \frac{1}{k^{2n}}. \quad (28)$$

Substituting (20) into (28) leads to

$$\sum_{k=1}^{\infty} (-1)^{n-1} \frac{x^{2(n-1)}}{(4\pi^2 k^2)^n} = (-1)^{n-1} \frac{B_n}{2(2n)!} x^{2(n-1)}, \quad 1 \leq n \leq m. \quad (29)$$

Summing up on both sides of (27) over  $k \in \mathbb{N}$  yields

$$\sum_{k=1}^{\infty} (-1)^m \frac{x^{2m}}{(4\pi^2 k^2)^{m+1}} \cdot \frac{1}{1 + \frac{x^2}{4\pi^2 k^2}} = (-1)^m \tilde{\theta} \frac{B_{m+1}}{2(2m+2)!} x^{2m}, \quad (30)$$

where  $\tilde{\theta}$  is a positive proper fraction (This means that  $0 < \tilde{\theta} < 1$ ) and dependent on  $x$ . Hence, we have the following

$$\frac{1}{e^x - 1} - \frac{1}{x} + \frac{1}{2} = \sum_{n=1}^m (-1)^{n-1} \frac{B_n}{(2n)!} x^{2n-1} + (-1)^m \tilde{\theta} \frac{B_{m+1}}{(2m+2)!} x^{2m+1}. \quad (31)$$

It is easy to see that

$$\int_{-\infty}^0 e^{xt} t^{2n-2} dt = \int_0^{\infty} e^{-xt} t^{2n-2} dt = \frac{(2n-2)!}{x^{2n-1}}, \quad (32)$$

$$\int_{-\infty}^0 \tilde{\theta} e^{xt} t^{2m} dt = \theta_1 \int_{-\infty}^0 e^{xt} t^{2m} dt = \theta_1 \frac{(2m)!}{x^{2m+1}}, \quad (33)$$

where  $0 < \theta_1 < 1$  and  $\theta_1$  is independent on  $x$ .

Substituting (32) and (33) into  $\omega(x)$  reveals

$$\omega(x) = \sum_{n=1}^m (-1)^{n-1} \frac{B_n}{2n(2n-1)} \cdot \frac{1}{x^{2n-1}} + (-1)^m \theta_1 \frac{B_{m+1}}{(2m+1)(2m+2)} \cdot \frac{1}{x^{2m+1}}, \quad (34)$$

and then formula (21) follows.

Differentiating on both sides of (25) yields

$$\frac{d}{dx} \ln \Gamma(x) = \ln x - \frac{1}{2x} + \omega'(x), \quad (35)$$



Easy computation gives

$$\int_{-\infty}^0 e^{xt} t^{2n-1} dt = -\frac{1}{x^{2n}} \int_0^{\infty} e^{-t} t^{2n-1} dt = -\frac{(2n-1)!}{x^{2n}}, \quad (36)$$

$$\int_{-\infty}^0 \tilde{\theta} e^{xt} t^{2m+1} dt = \theta_2 \int_{-\infty}^0 e^{xt} t^{2m+1} dt = -\theta_2 \frac{(2m+1)!}{x^{2m+2}}, \quad (37)$$

where  $\theta_2$  is independent of  $x$  and  $0 < \theta_2 < 1$ .

Substituting (31) into  $\omega'(x)$  and utilizing (36) and (37) shows

$$\omega'(x) = \sum_{n=1}^m (-1)^n \frac{B_n}{2n} \cdot \frac{1}{x^{2n}} + (-1)^{m+1} \theta_2 \frac{B_{m+1}}{2m+2} \cdot \frac{1}{x^{2m+2}}, \quad 0 < \theta_2 < 1. \quad (38)$$

Formula (22) follows from combining of (38) with  $\omega'(x)$ .

Differentiating with  $x$  on both sides of (35) yields

$$\frac{d^2}{dx^2} \ln \Gamma(x) = \frac{1}{x} + \frac{1}{2x^2} + \omega''(x), \quad (39)$$

substituting (31) into  $\omega''(x)$  and integrating directly produces

$$\begin{aligned} \omega''(x) &= \sum_{n=1}^m (-1)^{n-1} \frac{B_n}{(2n)!} \int_{-\infty}^0 t^{2n} e^{xt} dt + (-1)^m \frac{B_{m+1}}{(2m+2)!} \int_{-\infty}^0 \tilde{\theta} t^{2m+2} e^{xt} dt \\ &= \sum_{n=1}^m (-1)^{n-1} \frac{B_n}{x^{2n+1}} + (-1)^m \frac{B_{m+1}}{(2m+2)!} \theta_3 \int_{-\infty}^0 t^{2m+2} e^{xt} dt \\ &= \sum_{n=1}^m (-1)^{n-1} \frac{B_n}{x^{2n+1}} + (-1)^m \theta_3 \frac{B_{m+1}}{x^{2m+3}}, \end{aligned}$$

where  $\theta_3$  is independent of  $x$  and  $0 < \theta_3 < 1$ . Formula (23) follows.

By the same argument as above, we obtain

$$\omega'''(x) = \sum_{n=1}^m \frac{(-1)^n (2n+1) B_n}{x^{2n+3}} + \frac{(-1)^{m+1} \theta_4 (2m+3) B_{m+1}}{x^{2m+4}}, \quad (40)$$

where  $\theta_4$  is independent of  $x$  and  $0 < \theta_4 < 1$ . Then formula (24) is proved.  $\square$

### 3. PROOFS OF THEOREM 1 AND 2

*Proof of Theorem 1.* For  $\alpha > 0$ , let

$$f_\alpha(x) = \frac{\sqrt[x]{\Gamma(x+1)}}{x + \alpha \sqrt[x]{\Gamma(x+\alpha+1)}} \quad (41)$$

for  $x > -1$ . By direct calculation, we obtain

$$\ln f_\alpha(x) = \frac{\ln \Gamma(x+1)}{x} - \frac{\ln \Gamma(x+\alpha+1)}{x+\alpha}, \quad (42)$$

$$[\ln f_\alpha(x)]' = \left[ \frac{\psi(x+1)}{x} - \frac{\ln \Gamma(x+1)}{x^2} \right] - \left[ \frac{\psi(x+\alpha+1)}{x+\alpha} - \frac{\ln \Gamma(x+\alpha+1)}{(x+\alpha)^2} \right] \quad (43)$$

$$\triangleq g(x) - g(x+\alpha), \quad (44)$$

and

$$g'(x) = \frac{2 \ln \Gamma(x+1) - 2x\psi(x+1) + x^2\psi'(x+1)}{x^3} \triangleq \frac{h(x)}{x^3}, \quad (45)$$

$$g''(x) = \frac{x^3\psi''(x+1) - 3x^2\psi'(x+1) + 6x\psi(x+1) - 6\ln\Gamma(x+1)}{x^4} \triangleq \frac{p(x)}{x^4}, \quad (46)$$

where  $\psi(x) = \frac{\Gamma(x)'}{\Gamma(x)}$  is known as the digamma function, the logarithmic derivative of  $\Gamma(x)$ . Therefore, it is sufficient to verify  $h(x) < 0$  for  $x > 0$  and  $h(x) > 0$  for  $-1 < x < 0$ .

Using the inequality

$$\ln(1+t) \leq \frac{t(2+t)}{2(1+t)} \quad (47)$$

for  $t \geq 0$  in [38] and the special cases  $m = 2$  of formulas (21), (22) and (23), we have

$$\begin{aligned} h(x) &= 2\ln\Gamma(x+1) - 2x\psi(x+1) + x^2\psi'(x+1) \\ &< 2\left[\frac{1}{2}\ln(2\pi) + \left(x + \frac{1}{2}\right)\ln(x+1) - (x+1) + \frac{1}{12(x+1)}\right. \\ &\quad \left. - \frac{1}{360(x+1)^3} + \frac{1}{1260(x+1)^5}\right] - 2x\left[\ln(x+1) - \frac{1}{2(x+1)}\right. \\ &\quad \left. - \frac{1}{12(x+1)^2} + \frac{1}{120(x+1)^4} - \frac{1}{252(x+1)^6}\right] + x^2\left[\frac{1}{x+1}\right. \\ &\quad \left. + \frac{1}{2(x+1)^2} + \frac{1}{6(x+1)^3} - \frac{1}{30(x+1)^5} + \frac{1}{42(x+1)^7}\right] \\ &= \ln(2\pi) - 2 + \ln(x+1) - 2x + \frac{6x^2 + 6x + 1}{6(x+1)} \\ &\quad + \frac{3x^2 + x}{6(x+1)^2} + \frac{30x^2 - 1}{180(x+1)^3} - \frac{x}{60(x+1)^4} \\ &\quad + \frac{1 - 21x^2}{630(x+1)^5} + \frac{x}{126(x+1)^6} + \frac{x^2}{42(x+1)^7} \\ &< \ln(2\pi) - 2 - 2x + \frac{9x^2 + 12x + 1}{6(x+1)} + \frac{3x^2 + x}{6(x+1)^2} + \frac{30x^2 - 1}{180(x+1)^3} \\ &\quad - \frac{x}{60(x+1)^4} + \frac{1 - 21x^2}{630(x+1)^5} + \frac{x}{126(x+1)^6} + \frac{x^2}{42(x+1)^7} \\ &= \ln(2\pi) - 1 - \frac{x}{2} - \frac{1}{x+1} + \frac{1}{9(x+1)^3} \\ &\quad + \frac{1}{12(x+1)^4} - \frac{1}{18(x+1)^6} + \frac{1}{42(x+1)^7} \\ &\triangleq \ln(2\pi) - \frac{1}{2} + \phi\left(\frac{1}{x+1}\right), \end{aligned} \quad (48)$$

and, for  $y \in (0, 1]$ ,

$$\begin{aligned} \phi'(y) &= -1 + \frac{1}{2y^2} + \frac{y^2}{3} + \frac{y^3}{3} - \frac{y^5}{3} + \frac{y^6}{6}, \\ \phi''(y) &= -\frac{1}{y^3} + \frac{2y}{3} + y^2 - \frac{5y^4}{3} + y^5, \\ \phi^{(3)}(y) &= \frac{2}{3} + \frac{3}{y^4} + 2y - \frac{20y^3}{3} + 5y^4, \end{aligned}$$

$$\begin{aligned}\phi^{(4)}(y) &= 2 - \frac{12}{y^5} - 20y^2 + 20y^3, \\ \phi^{(5)}(y) &= \frac{60}{y^6} - 40y + 60y^2, \\ \phi^{(6)}(y) &= -40 - \frac{360}{y^7} + 120y, \\ \phi^{(7)}(y) &= 120 + \frac{2520}{y^8}.\end{aligned}$$

It is clear that  $\phi^{(7)}(y) > 0$  and  $\phi^{(6)}(y)$  is increasing. Since  $\phi^{(6)}(1) = -280$  and  $\lim_{y \rightarrow 0} \phi^{(6)}(y) = -\infty$ , we have  $\phi^{(6)}(y) < 0$  and  $\phi^{(5)}(y)$  is decreasing. It is easy to see that  $\lim_{y \rightarrow 0} \phi^{(5)}(y) = \infty$  and  $\phi^{(5)}(1) = 80$ , thus  $\phi^{(5)}(y) > 0$  and then  $\phi^{(4)}(y)$  is increasing. From  $\lim_{y \rightarrow 0} \phi^{(4)}(y) = -\infty$  and  $\phi^{(4)}(1) = -10$ , it is deduced that  $\phi^{(4)}(y) < 0$ , hence  $\phi^{(3)}(y)$  decreases. From  $\lim_{y \rightarrow 0} \phi^{(3)}(y) = \infty$  and  $\phi^{(3)}(1) = 4$ , it is concluded that  $\phi^{(3)}(y) > 0$ , therefore  $\phi''(y)$  increases. Because of  $\lim_{y \rightarrow 0} \phi''(y) = -\infty$  and  $\phi''(1) = 0$ , we obtain  $\phi''(y) \leq 0$ , then  $\phi'(y)$  decreases. By  $\lim_{y \rightarrow 0} \phi'(y) = \infty$  and  $\phi'(1) = 0$ , it follows that  $\phi'(y) \geq 0$ , and then  $\phi(y)$  is increasing in  $(0, 1]$ .

Utilizing monotonicity property of  $\phi(y)$  and the inequality

$$h(x) < \ln(2\pi) - \frac{1}{2} + \phi\left(\frac{1}{x+1}\right) \quad (49)$$

with

$$\ln(2\pi) - \frac{1}{2} + \phi\left(\frac{1}{0+1}\right) = \ln(2\pi) - \frac{1}{2} - \frac{337}{252} = \ln(2\pi) - \frac{463}{252} > 0 \quad (50)$$

and

$$\ln(2\pi) - \frac{1}{2} + \phi\left(\frac{1}{1+1}\right) = \ln(2\pi) - \frac{1}{2} - \frac{2655}{1792} = \ln(2\pi) - \frac{3551}{1792} < 0, \quad (51)$$

we conclude that there exists a point  $x_0 \in (0, 1)$  such that  $h(x) < 0$  in  $x \in (x_0, \infty)$ . This implies  $g'(x) < 0$  for  $x \in (x_0, \infty)$  and  $g(x)$  is decreasing in  $(x_0, \infty)$ . Hence  $[\ln f_\alpha(x)]' > 0$  in  $(x_0, \infty)$ , and then  $\ln f_\alpha(x)$  is increasing in  $(x_0, \infty)$ , that is,  $f_\alpha(x)$  is increasing in  $(x_0, \infty)$ . The proof is complete.  $\square$

*Proof of Theorem 2.* Using the inequality (47) for  $t \geq 0$  in [38] and the special cases  $m = 2$  of formulas (21), (22), (23) and (24), we obtain

$$\begin{aligned}p(x) &= x^3\psi''(x+1) - 3x^2\psi'(x+1) + 6x\psi(x+1) - 6\ln\Gamma(x+1) \\ &> x^3 \left[ -\frac{1}{(x+1)^2} - \frac{1}{(x+1)^3} - \frac{1}{2(x+1)^5} + \frac{1}{6(x+1)^7} - \frac{1}{6(x+1)^8} \right] \\ &\quad - 3x^2 \left[ \frac{1}{x+1} + \frac{1}{2(x+1)^2} + \frac{1}{6(x+1)^3} - \frac{1}{30(x+1)^5} + \frac{1}{42(x+1)^7} \right] \\ &\quad + 6x \left[ \ln(x+1) - \frac{1}{2(x+1)} - \frac{1}{12(x+1)^2} + \frac{1}{120(x+1)^4} - \frac{1}{252(x+1)^6} \right] \\ &\quad - 6 \left[ \frac{1}{2} \ln(2\pi) + \left( x + \frac{1}{2} \right) \ln(x+1) - (x+1) + \frac{1}{12(x+1)} \right. \\ &\quad \left. - \frac{1}{360(x+1)^3} + \frac{1}{1260(x+1)^5} \right]\end{aligned}$$

$$\begin{aligned}
 &> x^3 \left[ -\frac{1}{(x+1)^2} - \frac{1}{(x+1)^3} - \frac{1}{2(x+1)^5} + \frac{1}{6(x+1)^7} - \frac{1}{6(x+1)^8} \right] \\
 &\quad - 3x^2 \left[ \frac{1}{x+1} + \frac{1}{2(x+1)^2} + \frac{1}{6(x+1)^3} - \frac{1}{30(x+1)^5} + \frac{1}{42(x+1)^7} \right] \\
 &\quad + 6x \left[ -\frac{1}{2(x+1)} - \frac{1}{12(x+1)^2} + \frac{1}{120(x+1)^4} - \frac{1}{252(x+1)^6} \right] \\
 &\quad - 6 \left[ \frac{1}{2} \ln(2\pi) - (x+1) + \frac{1}{12(x+1)} - \frac{1}{360(x+1)^3} + \frac{1}{1260(x+1)^5} \right] \\
 &\quad - \frac{3x(2+x)}{2(1+x)} \\
 &= 4 - 3 \ln(2\pi) + \frac{x}{2} + \frac{3}{x+1} - \frac{5}{2(x+1)^2} + \frac{13}{6(x+1)^3} - \frac{19}{12(x+1)^4} \\
 &\quad - \frac{1}{6(x+1)^5} + \frac{7}{6(x+1)^6} - \frac{31}{42(x+1)^7} + \frac{1}{6(x+1)^8} \\
 &\triangleq (x+1)q\left(\frac{1}{x+1}\right),
 \end{aligned}$$

and, for  $t \in [0, \frac{1}{2}]$ ,

$$\begin{aligned}
 q'(t) &= \frac{7}{2} - 3 \ln(2\pi) + 6t - \frac{15t^2}{2} + \frac{26t^3}{3} - \frac{95t^4}{12} - t^5 + \frac{49t^6}{6} - \frac{124t^7}{21} + \frac{3t^8}{2}, \\
 q''(t) &= 6 - 15t + 26t^2 - \frac{95t^3}{3} - 5t^4 + 49t^5 - \frac{124t^6}{3} + 12t^7, \\
 q^{(3)}(t) &= -15 + 52t - 95t^2 - 20t^3 + 245t^4 - 248t^5 + 84t^6, \\
 q^{(4)}(t) &= 52 - 190t - 60t^2 + 980t^3 - 1240t^4 + 504t^5, \\
 q^{(5)}(t) &= -190 - 120t + 2940t^2 - 4960t^3 + 2520t^4, \\
 q^{(6)}(t) &= -120 + 5880t - 14880t^2 + 10080t^3, \\
 q^{(7)}(t) &= 5880 - 29760t + 30240t^2, \\
 q^{(8)}(t) &= -29760 + 60480t.
 \end{aligned}$$

It is easy to see that  $t = \frac{31}{63}$  is an unique minimum point of  $q^{(7)}(t)$  on  $[0, \frac{1}{2}]$ . Since  $q^{(7)}(0) = 5880$  and  $q^{(7)}(\frac{1}{2}) = -1440$ , thus  $q^{(6)}(t)$  has an unique maximum on  $[0, \frac{1}{2}]$ . Since  $q^{(6)}(0) = -120$  and  $q^{(6)}(\frac{1}{2}) = 360$ , then  $q^{(5)}(t)$  has an unique minimum on  $[0, \frac{1}{2}]$ . Because of  $q^{(5)}(0) = -190$  and  $q^{(5)}(\frac{1}{2}) = \frac{45}{2}$ , therefore  $q^{(4)}(t)$  has an unique minimum on  $[0, \frac{1}{2}]$ . The unique zero point of  $q^{(5)}(t)$  in  $[0, \frac{1}{2}]$  is

$$\begin{aligned}
 t_0 &= \frac{31}{63} + \frac{1}{756} \sqrt{\frac{1}{7} \left[ 381528 - 1323\kappa - 7938\mu + \frac{294848}{\nu} \right]} - \frac{\nu}{2} \quad (52) \\
 &= 0.4437889482188733 \dots,
 \end{aligned}$$

where

$$\kappa = \sqrt[3]{5600664 - 1296\sqrt{17855817}}, \quad (53)$$

$$\mu = \sqrt[3]{3(8643 + 2\sqrt{17855817})}, \quad (54)$$

$$\nu = \sqrt{\frac{757}{3969} + \frac{1}{756} \sqrt[3]{5600664 - 1296\sqrt{17855817}} + \frac{\sqrt[3]{8643 + 2\sqrt{17855817}}}{42\sqrt[3]{9}}}, \quad (55)$$

the minimum  $q^{(4)}(t_0) = 0.03717920\dots$ . Hence  $q^{(4)}(t) > 0$  and  $q^{(3)}(t)$  is increasing on  $[0, \frac{1}{2}]$ . From  $q^{(3)}(\frac{1}{2}) = -\frac{51}{8}$ , it follows that  $q^{(3)}(t) < 0$  and  $q''(t)$  is decreasing on  $[0, \frac{1}{2}]$ . From  $q''(\frac{1}{2}) = \frac{41}{24}$ , it is deduced that  $q''(t) > 0$  and  $q'(t)$  is increasing on  $[0, \frac{1}{2}]$ . Since  $q'(\frac{1}{2}) = \frac{56659}{10752} - 3\ln(2\pi) < 0$ , we have  $q'(t) < 0$  and  $q(t)$  is decreasing on  $[0, \frac{1}{2}]$ . From  $q(\frac{1}{2}) = \frac{22093}{21504} + \frac{1}{2}(\frac{7}{2} - 3\ln(2\pi)) > 0$ , it is concluded that the function  $q(t) > 0$  on  $[0, \frac{1}{2}]$ .

Note that  $q(t) > 0$  with  $t \in (0, \frac{1}{2}]$  is equivalent to  $q(\frac{1}{x+1}) > 0$  with  $x \in [1, \infty)$ . This implies that  $p(x) > 0$  and  $g''(x) > 0$  with  $x \in [1, \infty)$ , then  $g'(x)$  is increasing and  $g(x)$  is convex with  $x \in [1, \infty)$ . Therefore  $[\ln f_\alpha(x)]'' = g'(x) - g'(x+\alpha) < 0$ , that is, the function  $f_\alpha(x)$  is logarithmically concave on  $[1, \infty)$ . The proof is complete.  $\square$

*Remark 4.* To prove that  $\frac{\sqrt[3]{\Gamma(x+1)}}{x+\alpha\sqrt[3]{\Gamma(x+\alpha+1)}}$  is increasing and logarithmically concave with  $x \in (-1, \infty)$  for a given  $\alpha > 0$ , it is sufficient to verify

$$h(x) = x^3 \left( \tau''(x) - \frac{1}{x^2 + 1} \right) \leq 0, \quad (56)$$

$$p(x) = x^4 \tau'''(x) - 12 + 5x + \frac{2}{(1+x)^3} - \frac{11}{(1+x)^2} + \frac{21}{1+x} \geq 0, \quad (57)$$

where

$$\tau(x) = \frac{1}{x} \int_0^\infty \left( \frac{1}{t} - \frac{1}{e^t - 1} \right) e^{-t} \frac{1 - e^{-xt}}{t} dt. \quad (58)$$

We will give proofs of (56) and (57) in a subsequent article.

#### 4. COROLLARIES

As by-products, from Theorem 1 and 2, the following corollaries are deduced.

**Corollary 1.** *The function  $\frac{\psi(x+1)}{x} - \frac{\ln \Gamma(x+1)}{x^2}$  is decreasing and convex on  $[1, \infty)$ .*

**Corollary 2.** *For  $x \in [1, \infty)$ , we have*

$$\ln \Gamma(x+1) < x\psi(x+1) - \frac{x^2}{2}\psi'(x+1), \quad (59)$$

$$\ln \Gamma(x+1) < x\psi(x+1) - \frac{x^2}{2}\psi'(x+1) + \frac{x^3}{6}\psi''(x+1). \quad (60)$$

*Remark 5.* It is conjectured that Corollary 1 and 2 are valid on the interval  $(-1, \infty)$ .

**Corollary 3.** *Let  $n$  be natural number. Then the sequence*

$$\frac{\sqrt[n]{n!}}{n+k\sqrt[n+k]{(n+k+1)!}} \quad (61)$$

are increasing with  $k \in \mathbb{N}$ .

## 5. OPEN PROBLEMS

The function  $\frac{\sqrt[x]{\Gamma(x+1)}}{\sqrt[x+\alpha]{\Gamma(x+\alpha+1)}}$  can be expressed as

$$\frac{\sqrt[x]{\int_0^\infty t^x e^{-t} dt}}{\sqrt[x+\alpha]{\int_0^\infty t^{x+\alpha} e^{-t} dt}}, \quad (62)$$

where  $\int_0^\infty e^{-t} dt = 1$ . Then we propose the following

**Open Problem 1.** Let  $w(x) \geq 0$  be a nonnegative weight defined on a domain  $\Omega$  with  $\int_\Omega w(x) dx = 1$ . Find conditions about  $w(x)$  and  $f(x) \geq 0$  such that the ratio between two power means

$$\mathcal{Q}(t) = \frac{[\int_\Omega w(x) f^t(x) dx]^{1/t}}{[\int_\Omega w(x) f^{t+\alpha}(x) dx]^{1/(t+\alpha)}} \quad (63)$$

is monotonic or convex with  $t \in \mathbb{R}$  for a given  $\alpha > 0$ .

**Open Problem 2.** Find conditions about the positive sequence  $\{a_i\}_{n \in \mathbb{N}}$  such that the function

$$\mathcal{F}(r) = \left( \frac{\frac{1}{n} \sum_{i=1}^n a_i^r}{\frac{1}{n+m} \sum_{i=1}^{n+m} a_i^r} \right)^{1/r} \quad (64)$$

is monotonic or convex with  $r \in \mathbb{R}$ , where  $n$  and  $m$  are two given natural numbers. In particular, for  $\{a_i\}_{n \in \mathbb{N}}$  being the natural number sequence (that is,  $a_i = i$ ), show that the function  $\mathcal{F}(r)$  is decreasing strictly with  $r \in \mathbb{R}$ .

## REFERENCES

- [1] M. Abramowitz and I. A. Stegun (Eds.), *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, National Bureau of Standards, Applied Mathematics Series **55**, 4th printing, with corrections, Washington, 1965.
- [2] H. Alzer, *On an inequality of H. Minc and L. Sathre*, J. Math. Anal. Appl. **179** (1993), 396–402.
- [3] T. H. Chan, P. Gao and F. Qi, *On a generalization of Martins' inequality*, Monatsh. Math. **138** (2003), no. 3, 179–187. RGMIA Res. Rep. Coll. **4** (2001), no. 1, Art. 12, 93–101. Available online at <http://rgmia.vu.edu.au/v4n1.html>.
- [4] Ch.-P. Chen and F. Qi, *Extension of H. Alzer's inequality for negative powers*, Tamkang J. Math. **36** (2005), no. 1, accepted.
- [5] Ch.-P. Chen and F. Qi, *The inequality of Alzer for negative powers*, Octagon Math. Mag. **12** (2004), accepted.
- [6] Ch.-P. Chen and F. Qi, *Monotonicity results for the gamma function*, J. Inequal. Pure Appl. Math. **3** (2003), no. 2, Art. 44. Available online at [http://jipam.vu.edu.au/v4n2/065\\_02.html](http://jipam.vu.edu.au/v4n2/065_02.html). RGMIA Res. Rep. Coll. **5** (2002), suppl., Art. 16. Available online at [http://rgmia.vu.edu.au/v5\(E\).html](http://rgmia.vu.edu.au/v5(E).html).
- [7] Ch.-P. Chen, F. Qi, P. Cerone, and S. S. Dragomir, *Monotonicity of sequences involving convex and concave functions*, Math. Inequal. Appl. **6** (2003), no. 2, 229–239. RGMIA Res. Rep. Coll. **5** (2002), no. 1, Art. 1, 3–13. Available online at <http://rgmia.vu.edu.au/v5n1.html>.
- [8] Á. Elbert and A. Laforgia, *An inequality for the product of two integrals relating to the incomplete Gamma function*, J. Inequal. Appl. **5** (2000), 39–51.
- [9] N. Elezović, C. Giordano and J. Pečarić, *The best bounds in Gautschi's inequality*, Math. Inequal. Appl. **3** (2000), 239–252.
- [10] T. Erber, *The gamma function inequalities of Gurland and Gautschi*, Scand. Actuar. J. **1960** (1961), 27–28.

- [11] W. Gautschi, *Some elementary inequalities relating to the gamma and incomplete gamma function*, J. Math. Phys. **38** (1959), 77–81.
- [12] B.-N. Guo and F. Qi, *Generalization of Bernoulli polynomials*, Internat. J. Math. Ed. Sci. Tech. **33** (2002), no. 3, 428–431. RGMIA Res. Rep. Coll. **4** (2001), no. 4, Art. 10, 691–695. Available online at <http://rgmia.vu.edu.au/v4n4.html>.
- [13] B.-N. Guo and F. Qi, *Inequalities for generalized weighted mean values of convex function*, Math. Inequal. Appl. **4** (2001), no. 2, 195–202. RGMIA Res. Rep. Coll. **2** (1999), no. 7, Art. 11, 1059–1065. Available online at <http://rgmia.vu.edu.au/v2n7.html>.
- [14] B.-N. Guo and F. Qi, *Inequalities and monotonicity for the ratio of gamma functions*, Taiwanese J. Math. **7** (2003), no. 2, 239–247.
- [15] B.-N. Guo and F. Qi, *Inequalities and monotonicity of the ratio for the geometric means of a positive arithmetic sequence with arbitrary difference*, Tamkang. J. Math. **34** (2003), no. 3, 261–270.
- [16] J. D. Kečlić and P. M. Vasić, *Some inequalities for the gamma function*, Publ. Inst. Math. Beograd N. S. **11** (1971), 107–114.
- [17] D. Kershaw, *Some extensions of W. Gautschi's inequalities for the gamma function*, Math. Comp. **41** (1983), 607–611.
- [18] D. Kershaw and A. Laforgia, *Monotonicity results for the gamma function*, Atti Accad. Sci. Torino Cl. Sci. Fis. Mat. Natur. **119** (1985), 127–133.
- [19] J.-Ch. Kuang, *Chángyòng Bùděngshì (Applied Inequalities)*, 2nd ed., Hunan Education Press, Changsha, China, 1993. (Chinese)
- [20] A. Laforgia and S. Sismondi, *Monotonicity results and inequalities for the gamma and error functions*, J. Comp. Appl. Math. **23** (1988), 25–33.
- [21] Q.-M. Luo, B.-N. Guo, F. Qi, and L. Debnath, *Generalizations of Bernoulli numbers and polynomials*, Internat. J. Math. Math. Sci. **2003** (2003), no. 59, 3769–3776. RGMIA Res. Rep. Coll. **5** (2002), no. 2, Art. 12, 353–359. Available online at <http://rgmia.vu.edu.au/v5n2.html>.
- [22] Q.-M. Luo, T.-F. Guo, and F. Qi, *Relations of Bernoulli numbers and Euler numbers*, J. Henan Normal Univ. (Nat. Sci.) **31** (2003), no. 2, 9–11. (Chinese)
- [23] Q.-M. Luo and F. Qi, *Relationships between generalized Bernoulli numbers and polynomials and generalized Euler numbers and polynomials*, Adv. Stud. Contemp. Math. (Kyungshang) **7** (2003), no. 1, 11–18. RGMIA Res. Rep. Coll. **5** (2002), no. 3, Art. 1, 405–412. Available online at <http://rgmia.vu.edu.au/v5n3.html>.
- [24] Q.-M. Luo, F. Qi, and L. Debnath, *Generalizations of Euler numbers and polynomials*, Internat. J. Math. Math. Sci. **2003** (2003), no. 61, 3893–3901. RGMIA Res. Rep. Coll. **5** (2002), suppl., Art. 4. Available online at [http://rgmia.vu.edu.au/v5\(E\).html](http://rgmia.vu.edu.au/v5(E).html).
- [25] Q.-M. Luo, Y.-M. Zheng, and F. Qi, *The higher order Euler numbers and higher order Euler polynomials*, Henan Sci. **21** (2003), no. 1, 1–6. (Chinese)
- [26] J. S. Martins, *Arithmetic and geometric means, an applications to Lorentz sequence spaces*, Math Nachr. **139** (1988), 281–288.
- [27] H. Minc and L. Sathre, *Some inequalities involving  $(r!)^{1/r}$* , Proc. Edinburgh Math. Soc. **14** (1964/65), 41–46.
- [28] D. S. Mitrinović, J. E. Pečarić and A. M. Fink, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, Dordrecht/Boston/London, 1993.
- [29] F. Qi, *An algebraic inequality*, J. Inequal. Pure Appl. Math. **2** (2001), no. 1, Art. 13. Available online at [http://jipam.vu.edu.au/v2n1/006\\_00.html](http://jipam.vu.edu.au/v2n1/006_00.html). RGMIA Res. Rep. Coll. **2** (1999), no. 1, Art. 8, 81–83. Available online at <http://rgmia.vu.edu.au/v2n1.html>.
- [30] F. Qi, *Generalized abstracted mean values*, J. Inequal. Pure Appl. Math. **1** (2000), no. 1, Art. 4. Available online at [http://jipam.vu.edu.au/v1n1/013\\_99.html](http://jipam.vu.edu.au/v1n1/013_99.html). RGMIA Res. Rep. Coll. **2** (1999), no. 5, Art. 4, 633–642. Available online at <http://rgmia.vu.edu.au/v2n5.html>.
- [31] F. Qi, *Generalization of H. Alzer's Inequality*, J. Math. Anal. Appl. **240** (1999), 294–297.
- [32] F. Qi, *Generalized weighted mean values with two parameters*, R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci. **454** (1998), no. 1978, 2723–2732.
- [33] F. Qi, *Inequalities and monotonicity of sequences involving  $\sqrt[n]{(n+k)!/k!}$* , Soochow J. Math. **29** (2004), no. 4, 353–361. RGMIA Res. Rep. Coll. **2** (1999), no. 5, Art. 8, 685–692. Available online at <http://rgmia.vu.edu.au/v2n5.html>.

- [34] F. Qi, *Inequalities and monotonicity of the ratio for the geometric means of a positive arithmetic sequence with unit difference*, Internat. J. Math. Edu. Sci. Tech. **34** (2003), no. 4, 601–607. Australian Math. Soc. Gaz. **30** (2003), no. 3, 142–147. RGMIA Res. Rep. Coll. **6** (2003), suppl., Art. 2. Available online at [http://rgmia.vu.edu.au/v6\(E\).html](http://rgmia.vu.edu.au/v6(E).html).
- [35] F. Qi, *Monotonicity results and inequalities for the gamma and incomplete gamma functions*, Math. Inequal. Appl. **5** (2002), no. 1, 61–67. RGMIA Res. Rep. Coll. **2** (1999), no. 7, Art. 7, 1027–1034. Available online at <http://rgmia.vu.edu.au/v2n7.html>.
- [36] F. Qi, *On a new generalization of Martins' inequality*, RGMIA Res. Rep. Coll. **5** (2002), no. 3, Art. 13, 527–538. Available online at <http://rgmia.vu.edu.au/v5n3.html>.
- [37] F. Qi, *The extended mean values: definition, properties, monotonicities, comparison, convexities, generalizations, and applications*, Cubo Mathematica Educational **6** (2004), no. 1, in press. RGMIA Res. Rep. Coll. **5** (2002), no. 1, Art. 5, 57–80. Available online at <http://rgmia.vu.edu.au/v5n1.html>.
- [38] F. Qi and Ch.-P. Chen, *Monotonicities of two sequences*, Math. Inform. Quart. **9** (1999), no. 4, 136–139.
- [39] F. Qi and Ch.-P. Chen, *Monotonicity and convexity results for functions involving the gamma function*, RGMIA Res. Rep. Coll. **6** (2003). Available online at <http://rgmia.vu.edu.au>.
- [40] F. Qi, L.-H. Cui, and S.-L. Xu, *Some inequalities constructed by Tchebysheff's integral inequality*, Math. Inequal. Appl. **2** (1999), no. 4, 517–528.
- [41] F. Qi and B.-N. Guo, *An inequality between ratio of the extended logarithmic means and ratio of the exponential means*, Taiwanese J. Math. **7** (2003), no. 2, 229–237. RGMIA Res. Rep. Coll. **4** (2001), no. 1, Art. 8, 55–61. Available online at <http://rgmia.vu.edu.au/v4n1.html>.
- [42] F. Qi and B.-N. Guo, *Monotonicity of sequences involving convex function and sequence*, RGMIA Res. Rep. Coll. **3** (2000), no. 2, Art. 14, 321–329. Available online at <http://rgmia.vu.edu.au/v3n2.html>.
- [43] F. Qi and B.-N. Guo, *Monotonicity of sequences involving geometric mean of positive sequences*, Nonlinear Funct. Anal. Appl. **8** (2003), no. 4, in press.
- [44] F. Qi and B.-N. Guo, *Monotonicity of sequences involving geometric means of positive sequences with logarithmical convexity*, RGMIA Res. Rep. Coll. **5** (2002), no. 3, Art. 10, 497–507. Available online at <http://rgmia.vu.edu.au/v5n3.html>.
- [45] F. Qi and B.-N. Guo, *Some inequalities involving the geometric mean of natural numbers and the ratio of gamma functions*, RGMIA Res. Rep. Coll. **4** (2001), no. 1, Art. 6, 41–48. Available online at <http://rgmia.vu.edu.au/v4n1.html>.
- [46] F. Qi and S.-L. Guo, *Inequalities for the incomplete gamma and related functions*, Math. Inequal. Appl. **2** (1999), no. 1, 47–53.
- [47] F. Qi, B.-N. Guo, and L. Debnath, *A lower bound for ratio of power means*, Internat. J. Math. Math. Sci. **2003** (2003), in press. RGMIA Res. Rep. Coll. **5** (2002), no. 4, Art. 2. Available online at <http://rgmia.vu.edu.au/v5n4.html>.
- [48] F. Qi and Q.-M. Luo, *Generalization of H. Minc and J. Sathre's inequality*, Tamkang J. Math. **31** (2000), no. 2, 145–148. RGMIA Res. Rep. Coll. **2** (1999), no. 6, Art. 14, 909–912. Available online at <http://rgmia.vu.edu.au/v2n6.html>.
- [49] F. Qi and J.-Q. Mei, *Some inequalities for the incomplete gamma and related functions*, Z. Anal. Anwendungen **18** (1999), no. 3, 793–799.
- [50] F. Qi and N. Towghi, *Inequalities for the ratios of the mean values of functions*, Nonlinear Funct. Anal. Appl. **9** (2004), no. 1, in press.
- [51] F. Qi and Sh.-Q. Zhang, *Note on monotonicity of generalized weighted mean values*, R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci. **455** (1999), no. 1989, 3259–3260.
- [52] H. R. van der Vaart, *Some extensions of the idea of Bias*, Ann. Math. Statist. **32** (1961), 436–447.
- [53] H. Vogt and J. Voigt, *A monotonicity property of the  $\Gamma$ -function*, J. Inequal. Pure Appl. Math. **3** (2002), no. 5, Art. 73. Available online at [http://jipam.vu.edu.au/v3n5/007\\_01.html](http://jipam.vu.edu.au/v3n5/007_01.html).
- [54] Zh.-X. Wang and D.-R. Guo, *Tèshū Hánshù Gàilùn (Introduction to Special Function)*, The Series of Advanced Physics of Peking University, Peking University Press, Beijing, CHINA, 2000. (Chinese)
- [55] Z.-K. Xu and D.-P. Xu, *A general form of Alzer's inequality*, Comput. Math. Appl. **44** (2002), 365–373.



(F. Qi) DEPARTMENT OF APPLIED MATHEMATICS AND INFORMATICS, JIAOZUO INSTITUTE OF TECHNOLOGY, JIAOZUO CITY, HENAN 454000, CHINA

*E-mail address:* [qifeng@jzit.edu.cn](mailto:qifeng@jzit.edu.cn), [fengqi618@member.ams.org](mailto:fengqi618@member.ams.org)

*URL:* <http://rgmia.vu.edu.au/qi.html>

(B.-N. Guo) DEPARTMENT OF APPLIED MATHEMATICS AND INFORMATICS, JIAOZUO INSTITUTE OF TECHNOLOGY, JIAOZUO CITY, HENAN 454000, CHINA

*E-mail address:* [guobaini@jzit.edu.cn](mailto:guobaini@jzit.edu.cn)