



**VICTORIA UNIVERSITY**  
MELBOURNE AUSTRALIA

*On Some Weighted Integral Inequalities for Convex Functions Related to Fejér's Result*

This is the Published version of the following publication

Tseng, Kuei-Lin, Hwang, Shioh-Ru and Dragomir, Sever S (2009) On Some Weighted Integral Inequalities for Convex Functions Related to Fejér's Result. Research report collection, 12 (Supp).

The publisher's official version can be found at

Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/17953/>

# ON SOME WEIGHTED INTEGRAL INEQUALITIES FOR CONVEX FUNCTIONS RELATED TO FEJÉR'S RESULT

K.-L. TSENG, SHIOW-RU HWANG, AND S.S. DRAGOMIR

ABSTRACT. In this paper, we introduce some functionals associated with weighted integral means for convex functions. Some new Fejér-type inequalities are obtained as well.

## 1. INTRODUCTION

Throughout this paper, let  $f : [a, b] \rightarrow \mathbb{R}$  be convex,  $g : [a, b] \rightarrow [0, \infty)$  be integrable and symmetric to  $\frac{a+b}{2}$ . We define the following mappings on  $[0, 1]$  that are associated with the well known *Hermite-Hadamard inequality* [1]

$$(1.1) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2},$$

namely

$$G(t) = \frac{1}{2} \left[ f\left(ta + (1-t)\frac{a+b}{2}\right) + f\left(tb + (1-t)\frac{a+b}{2}\right) \right];$$

$$Q(t) = \frac{1}{2} [f(ta + (1-t)b) + f(tb + (1-t)a)];$$

$$H(t) = \frac{1}{b-a} \int_a^b f\left(tx + (1-t)\frac{a+b}{2}\right) dx;$$

$$H_g(t) = \int_a^b f\left(tx + (1-t)\frac{a+b}{2}\right) g(x) dx;$$

$$I(t) = \int_a^b \frac{1}{2} \left[ f\left(t\frac{x+a}{2} + (1-t)\frac{a+b}{2}\right) + f\left(t\frac{x+b}{2} + (1-t)\frac{a+b}{2}\right) \right] g(x) dx;$$

$$P(t) = \frac{1}{2(b-a)} \int_a^b \left[ f\left(\left(\frac{1+t}{2}\right)a + \left(\frac{1-t}{2}\right)x\right) + f\left(\left(\frac{1+t}{2}\right)b + \left(\frac{1-t}{2}\right)x\right) \right] dx;$$

---

1991 *Mathematics Subject Classification.* 26D15.

*Key words and phrases.* Hermite-Hadamard inequality, Fejér inequality, Convex function.

This research was partially supported by grant NSC 98-2115-M-156-004.

$$\begin{aligned}
P_g(t) &= \int_a^b \frac{1}{2} \left[ f \left( \left( \frac{1+t}{2} \right) a + \left( \frac{1-t}{2} \right) x \right) g \left( \frac{x+a}{2} \right) \right. \\
&\quad \left. + f \left( \left( \frac{1+t}{2} \right) b + \left( \frac{1-t}{2} \right) x \right) g \left( \frac{x+b}{2} \right) \right] dx; \\
N(t) &= \int_a^b \frac{1}{2} \left[ f \left( ta + (1-t) \frac{x+a}{2} \right) + f \left( tb + (1-t) \frac{x+b}{2} \right) \right] g(x) dx; \\
L(t) &= \frac{1}{2(b-a)} \int_a^b [f(ta + (1-t)x) + f(tb + (1-t)x)] dx; \\
L_g(t) &= \frac{1}{2} \int_a^b [f(ta + (1-t)x) + f(tb + (1-t)x)] g(x) dx
\end{aligned}$$

and

$$\begin{aligned}
S_g(t) &= \frac{1}{4} \int_a^b \left[ f \left( ta + (1-t) \frac{x+a}{2} \right) + f \left( ta + (1-t) \frac{x+b}{2} \right) \right. \\
&\quad \left. + f \left( tb + (1-t) \frac{x+a}{2} \right) + f \left( tb + (1-t) \frac{x+b}{2} \right) \right] g(x) dx.
\end{aligned}$$

**Remark 1.** We note that  $H = H_g = I, P = P_g = N$  and  $L = L_g = S_g$  on  $[0, 1]$  as  $g(x) = \frac{1}{b-a}$  ( $x \in [a, b]$ ).

For some results which generalize, improve, and extend the famous Hermite-Hadamard integral inequality, see [2] – [19].

In [8], Fejér established the following weighted generalization of the Hermite-Hadamard inequality (1.1) :

**Theorem A.** Let  $f, g$  be defined as above. Then

$$(1.2) \quad f \left( \frac{a+b}{2} \right) \int_a^b g(x) dx \leq \int_a^b f(x) g(x) dx \leq \frac{f(a) + f(b)}{2} \int_a^b g(x) dx.$$

In [11], Tseng et al. established the following Fejér-type inequalities.

**Theorem B.** Let  $f, g$  be defined as above. Then we have

$$\begin{aligned}
(1.3) \quad f \left( \frac{a+b}{2} \right) \int_a^b g(x) dx &\leq \frac{f \left( \frac{3a+b}{4} \right) + f \left( \frac{a+3b}{4} \right)}{2} \int_a^b g(x) dx \\
&\leq \int_a^b \frac{1}{2} \left[ f \left( \frac{x+a}{2} \right) + f \left( \frac{x+b}{2} \right) \right] g(x) dx \\
&\leq \frac{1}{2} \left[ f \left( \frac{a+b}{2} \right) + \frac{f(a) + f(b)}{2} \right] \int_a^b g(x) dx \\
&\leq \frac{f(a) + f(b)}{2} \int_a^b g(x) dx.
\end{aligned}$$

In [2], Dragomir established the following Hermite-Hadamard-type inequality which refines the first inequality of (1.1).

**Theorem C.** Let  $f, H$  be defined as above. Then  $H$  is convex, increasing on  $[0, 1]$ , and for all  $t \in [0, 1]$ , we have

$$(1.4) \quad f \left( \frac{a+b}{2} \right) = H(0) \leq H(t) \leq H(1) = \frac{1}{b-a} \int_a^b f(x) dx.$$

In [15], Yang and Hong obtained the following Hermite-Hadamard-type inequality which is a refinement of the second inequality in (1.1).

**Theorem D.** *Let  $f, P$  be defined as above. Then  $P$  is convex, increasing on  $[0, 1]$ , and for all  $t \in [0, 1]$ , we have*

$$(1.5) \quad \frac{1}{b-a} \int_a^b f(x) dx = P(0) \leq P(t) \leq P(1) = \frac{f(a) + f(b)}{2}.$$

Yang and Tseng [16] and Tseng et al. [11] established the following Fejér-type inequalities which are weighted generalizations of Theorems C – D.

**Theorem E** ([16]). *Let  $f, g, H_g, P_g$  be defined as above. Then  $H_g, P_g$  are convex, increasing on  $[0, 1]$ , and for all  $t \in [0, 1]$ , we have*

$$(1.6) \quad \begin{aligned} f\left(\frac{a+b}{2}\right) \int_a^b g(x) dx &= H_g(0) \leq H_g(t) \leq H_g(1) \\ &= \int_a^b f(x) g(x) dx \\ &= P_g(0) \leq P_g(t) \leq P_g(1) \\ &= \frac{f(a) + f(b)}{2} \int_a^b g(x) dx. \end{aligned}$$

**Theorem F** ([11]). *Let  $f, g, I, N$  be defined as above. Then  $I, N$  are convex, increasing on  $[0, 1]$ , and for all  $t \in [0, 1]$ , we have*

$$(1.7) \quad \begin{aligned} f\left(\frac{a+b}{2}\right) \int_a^b g(x) dx &= I(0) \leq I(t) \leq I(1) \\ &= \int_a^b \frac{1}{2} \left[ f\left(\frac{x+a}{2}\right) + f\left(\frac{x+b}{2}\right) \right] g(x) dx \\ &= N(0) \leq N(t) \leq N(1) \\ &= \frac{f(a) + f(b)}{2} \int_a^b g(x) dx. \end{aligned}$$

In [7], Dragomir et al. established the following Hermite-Hadamard-type inequality.

**Theorem G.** *Let  $f, H, G, L$  be defined as above. Then  $G$  is convex, increasing on  $[0, 1]$ ,  $L$  is convex on  $[0, 1]$ , and for all  $t \in [0, 1]$ , we have*

$$(1.8) \quad H(t) \leq G(t) \leq L(t) \leq \frac{1-t}{b-a} \int_a^b f(x) dx + t \cdot \frac{f(a) + f(b)}{2} \leq \frac{f(a) + f(b)}{2}.$$

In [12] – [13], Tseng et al. obtained the following theorems related to Fejér's result which in their turn are weighted generalizations of the inequality (1.8).

**Theorem H** ([12]). *Let  $f, g, G, H_g, L_g$  be defined as above. Then  $L_g$  is convex, increasing on  $[0, 1]$ , and for all  $t \in [0, 1]$ , we have*

$$\begin{aligned}
 (1.9) \quad H_g(t) &\leq G(t) \int_a^b g(x) dx \\
 &\leq L_g(t) \\
 &\leq (1-t) \int_a^b f(x) g(x) dx + t \cdot \frac{f(a) + f(b)}{2} \int_a^b g(x) dx \\
 &\leq \frac{f(a) + f(b)}{2} \int_a^b g(x) dx.
 \end{aligned}$$

**Theorem I** ([13]). *Let  $f, g, G, I, S_g$  be defined as above. Then  $S_g$  is convex, increasing on  $[0, 1]$ , and for all  $t \in [0, 1]$ , we have*

$$\begin{aligned}
 (1.10) \quad I(t) &\leq G(t) \int_a^b g(x) dx \leq S_g(t) \\
 &\leq (1-t) \int_a^b \frac{1}{2} \left[ f\left(\frac{x+a}{2}\right) + f\left(\frac{x+b}{2}\right) \right] g(x) dx \\
 &\quad + t \cdot \frac{f(a) + f(b)}{2} \int_a^b g(x) dx \\
 &\leq \frac{f(a) + f(b)}{2} \int_a^b g(x) dx.
 \end{aligned}$$

In this paper, we provide some new Fejér-type inequalities related to the mappings  $G, Q, H_g, P_g, I, N, L_g, S_g$  defined above. They generalize known results obtained in relation with the Hermite-Hadamard inequality and therefore are useful in obtaining various results for means when the convex function and the weight take particular forms.

## 2. MAIN RESULTS

The following lemmatae are needed in the proofs of our main results:

**Lemma 2** (see [9]). *Let  $f$  be defined as above and let  $a \leq A \leq C \leq D \leq B \leq b$  with  $A + B = C + D$ . Then*

$$f(C) + f(D) \leq f(A) + f(B).$$

The assumptions in Lemma 2 can be weakened as in the following lemma:

**Lemma 3.** *Let  $f$  be defined as above and let  $a \leq A \leq C \leq B \leq b$  and  $a \leq A \leq D \leq B \leq b$  with  $A + B = C + D$ . Then*

$$f(C) + f(D) \leq f(A) + f(B).$$

**Lemma 4** (see [14]). *Let  $f, G, Q$  be defined as above. Then  $Q$  is symmetric about  $\frac{1}{2}$ ,  $Q$  is decreasing on  $[0, \frac{1}{2}]$  and increasing on  $[\frac{1}{2}, 1]$ ,*

$$\begin{aligned}
 G(2t) &\leq Q(t) && \left( t \in \left[0, \frac{1}{4}\right] \right), \\
 G(2t) &\geq Q(t) && \left( t \in \left[\frac{1}{4}, \frac{1}{2}\right] \right),
 \end{aligned}$$

$$G(2(1-t)) \geq Q(t) \quad \left( t \in \left[ \frac{1}{2}, \frac{3}{4} \right] \right)$$

and

$$G(2(1-t)) \leq Q(t) \quad \left( t \in \left[ \frac{3}{4}, 1 \right] \right).$$

Now, we are ready to state and prove our results.

**Theorem 5.** *Let  $f, g, G, H_g, P_g, L_g, S_g$  be defined as above. Then:*

(1) *The inequality*

$$(2.1) \quad \int_a^b f(x)g(x) dx \leq 2 \left[ \int_a^{\frac{3a+b}{4}} f(x)g(2x-a) dx + \int_{\frac{a+3b}{4}}^b f(x)g(2x-b) dx \right] \\ \leq \int_0^1 P_g(t) dt \\ \leq \frac{1}{2} \left[ \int_a^b f(x)g(x) dx + \frac{f(a)+f(b)}{2} \int_a^b g(x) dx \right]$$

*holds.*

(2) *The inequalities*

$$(2.2) \quad L_g(t) \leq P_g(t) \\ \leq (1-t) \int_a^b f(x)g(x) dx + t \cdot \frac{f(a)+f(b)}{2} \int_a^b g(x) dx \\ \leq \frac{f(a)+f(b)}{2} \int_a^b g(x) dx$$

and

$$(2.3) \quad 0 \leq N(t) - G(t) \int_a^b g(x) dx \leq \frac{f(a)+f(b)}{2} \int_a^b g(x) dx - N(t)$$

*hold for all  $t \in [0, 1]$ .*

(3) *If  $f$  is differentiable on  $[a, b]$ , then we have the inequalities*

$$(2.4) \quad 0 \leq t \left[ \frac{1}{b-a} \int_a^b f(x) dx - f\left(\frac{a+b}{2}\right) \right] \cdot \inf_{x \in [a,b]} g(x) \\ \leq P_g(t) - \int_a^b f(x)g(x) dx;$$

$$(2.5) \quad 0 \leq P_g(t) - f\left(\frac{a+b}{2}\right) \int_a^b g(x) dx \\ \leq \frac{(f'(b) - f'(a))(b-a)}{4} \int_a^b g(x) dx;$$

$$(2.6) \quad 0 \leq L_g(t) - H_g(t) \leq \frac{(f'(b) - f'(a))(b-a)}{4} \int_a^b g(x) dx;$$

$$(2.7) \quad 0 \leq P_g(t) - L_g(t) \leq \frac{(f'(b) - f'(a))(b-a)}{4} \int_a^b g(x) dx;$$

$$(2.8) \quad 0 \leq P_g(t) - H_g(t) \leq \frac{(f'(b) - f'(a))(b-a)}{4} \int_a^b g(x) dx;$$

$$(2.9) \quad 0 \leq N(t) - I(t) \leq \frac{(f'(b) - f'(a))(b-a)}{4} \int_a^b g(x) dx$$

and

$$(2.10) \quad 0 \leq S_g(t) - I(t) \leq \frac{(f'(b) - f'(a))(b-a)}{4} \int_a^b g(x) dx$$

for all  $t \in [0, 1]$ .

*Proof.* (1) By using simple integration techniques and the hypothesis of  $g$ , we have the following identities

$$(2.11) \quad \int_a^b f(x) g(x) dx = 2 \int_a^{\frac{a+b}{2}} \int_0^{\frac{1}{2}} [f(x) + f(a+b-x)] g(x) dt dx;$$

$$(2.12) \quad \begin{aligned} & 2 \left[ \int_a^{\frac{3a+b}{4}} f(x) g(2x-a) dx + \int_{\frac{a+3b}{4}}^b f(x) g(2x-b) dx \right] \\ &= 2 \int_a^{\frac{3a+b}{4}} [f(x) + f(a+b-x)] g(2x-a) dx \\ &= 2 \int_a^{\frac{a+b}{2}} \int_0^{\frac{1}{2}} \left[ f\left(\frac{a+x}{2}\right) + f\left(\frac{a+2b-x}{2}\right) \right] g(x) dt dx; \end{aligned}$$

$$(2.13) \quad \begin{aligned} \int_0^1 P_g(t) dt &= \int_a^{\frac{a+b}{2}} \int_0^1 f(ta + (1-t)x) g(x) dt dx \\ &\quad + \int_{\frac{a+b}{2}}^b \int_0^1 f(tb + (1-t)x) g(x) dt dx \\ &= \int_a^{\frac{a+b}{2}} \int_0^1 f(ta + (1-t)x) g(x) dt dx \\ &\quad + \int_a^{\frac{a+b}{2}} \int_0^1 f(tb + (1-t)(a+b+x)) g(x) dt dx \\ &= \int_a^{\frac{a+b}{2}} \int_0^{\frac{1}{2}} [f(tx + (1-t)a) + f(ta + (1-t)x)] g(x) dt dx \\ &\quad + \int_a^{\frac{a+b}{2}} \int_0^{\frac{1}{2}} [f(tb + (1-t)(a+b-x)) \\ &\quad + f(t(a+b-x) + (1-t)b)] g(x) dt dx \end{aligned}$$

and

$$\begin{aligned}
 (2.14) \quad & \frac{1}{2} \left[ \int_a^b f(x) g(x) dx + \frac{f(a) + f(b)}{2} \int_a^b g(x) dx \right] \\
 &= \int_a^{\frac{a+b}{2}} \int_0^{\frac{1}{2}} [f(a) + f(x)] g(x) dt dx \\
 & \quad + \int_a^{\frac{a+b}{2}} \int_0^{\frac{1}{2}} [f(a+b-x) + f(b)] g(x) dt dx.
 \end{aligned}$$

By Lemma 2, the following inequalities hold for all  $t \in [0, \frac{1}{2}]$  and  $x \in [a, \frac{a+b}{2}]$ .

$$(2.15) \quad f(x) + f(a+b-x) \leq f\left(\frac{a+x}{2}\right) + f\left(\frac{a+2b-x}{2}\right)$$

holds when  $A = \frac{a+x}{2}$ ,  $C = x$ ,  $D = a+b-x$  and  $B = \frac{a+2b-x}{2}$  in Lemma 2.

$$(2.16) \quad f\left(\frac{a+x}{2}\right) \leq \frac{1}{2} [f(tx + (1-t)a) + f(ta + (1-t)x)]$$

holds when  $A = tx + (1-t)a$ ,  $C = D = \frac{a+x}{2}$  and  $B = ta + (1-t)x$  in Lemma 2.

$$\begin{aligned}
 (2.17) \quad & f\left(\frac{a+2b-x}{2}\right) \\
 & \leq \frac{1}{2} [f(tb + (1-t)(a+b-x)) + f(t(a+b-x) + (1-t)b)]
 \end{aligned}$$

holds when  $A = tb + (1-t)(a+b-x)$ ,  $C = D = \frac{a+2b-x}{2}$  and  $B = t(a+b-x) + (1-t)b$  in Lemma 2.

$$(2.18) \quad \frac{1}{2} [f(tx + (1-t)a) + f(ta + (1-t)x)] \leq \frac{f(a) + f(x)}{2}$$

holds when  $A = a$ ,  $C = tx + (1-t)a$ ,  $D = ta + (1-t)x$  and  $B = x$  in Lemma 2.

$$\begin{aligned}
 (2.19) \quad & \frac{1}{2} [f(tb + (1-t)(a+b-x)) + f(t(a+b-x) + (1-t)b)] \\
 & \leq \frac{f(a+b-x) + f(b)}{2}
 \end{aligned}$$

holds as  $A = a+b-x$ ,  $C = tb + (1-t)(a+b-x)$ ,  $D = t(a+b-x) + (1-t)b$  and  $B = b$  in Lemma 2. Multiplying the inequalities (2.15) – (2.19) by  $g(x)$  and integrating them over  $t$  on  $[0, \frac{1}{2}]$ , over  $x$  on  $[a, \frac{a+b}{2}]$  and using identities (2.11) – (2.14), we derive (2.1).



(2) Using substitution rules for integration and the hypothesis of  $g$ , we have the following identities

$$\begin{aligned}
 (2.20) \quad P_g(t) &= \int_a^{\frac{a+b}{2}} f(ta + (1-t)x) g(x) dx \\
 &\quad + \int_{\frac{a+b}{2}}^b f(tb + (1-t)x) g(x) dx \\
 &= \int_a^{\frac{a+b}{2}} [f(ta + (1-t)x) \\
 &\quad + f(tb + (1-t)(a+b-x))] g(x) dx
 \end{aligned}$$

and

$$\begin{aligned}
 (2.21) \quad L_g(t) &= \frac{1}{2} \left[ \int_a^{\frac{a+b}{2}} f(ta + (1-t)x) g(x) dx \right. \\
 &\quad \left. + \int_{\frac{a+b}{2}}^b f(tb + (1-t)x) g(x) dx \right] \\
 &\quad + \frac{1}{2} \left[ \int_{\frac{a+b}{2}}^b f(ta + (1-t)x) g(x) dx \right. \\
 &\quad \left. + \int_a^{\frac{a+b}{2}} f(tb + (1-t)x) g(x) dx \right] \\
 &= \frac{1}{2} P_g(t) + \frac{1}{2} \int_a^{\frac{a+b}{2}} [f(ta + (1-t)(a+b-x)) \\
 &\quad + f(tb + (1-t)x)] g(x) dx
 \end{aligned}$$

for all  $t \in [0, 1]$ .

If we choose  $A = ta + (1-t)x$ ,  $C = ta + (1-t)(a+b-x)$ ,  $D = tb + (1-t)x$  and  $B = tb + (1-t)(a+b-x)$  in Lemma 3, then the inequality

$$\begin{aligned}
 (2.22) \quad &f(ta + (1-t)(a+b-x)) + f(tb + (1-t)x) \\
 &\leq f(ta + (1-t)x) + f(tb + (1-t)(a+b-x))
 \end{aligned}$$

holds for all  $t \in [0, 1]$  and  $x \in [a, \frac{a+b}{2}]$ . Multiplying the inequality (2.22) by  $g(x)$ , integrating both sides over  $x$  on  $[a, \frac{a+b}{2}]$  and using identities (2.20) – (2.21), we derive the first inequality of (2.2). The second and third inequalities of (2.2) can be obtained by the convexity of  $f$  and (1.2). This proves (2.2).

Again, using substitution rules for integration and the hypothesis of  $g$ , we have the following identity

$$\begin{aligned}
 N(t) &= \int_a^b \frac{1}{2} \left[ f\left(ta + (1-t)\frac{x+a}{2}\right) \right. \\
 &\quad \left. + f\left(tb + (1-t)\frac{a+2b-x}{2}\right) \right] g(x) dx
 \end{aligned}$$

$$\begin{aligned}
(2.23) \quad &= \int_a^{\frac{a+b}{2}} [f(ta + (1-t)x) \\
&\quad + f(tb + (1-t)(a+b-x))] g(2x-a) dx \\
(2.24) \quad &= \int_a^{\frac{3a+b}{4}} [f(ta + (1-t)x) \\
&\quad + f\left(ta + (1-t)\left(\frac{3a+b}{2} - x\right)\right) \\
&\quad + f\left(tb + (1-t)\left(\frac{b-a}{2} + x\right)\right) \\
&\quad + f(tb + (1-t)(a+b-x))] g(2x-a) dx
\end{aligned}$$

for all  $t \in [0, 1]$ . By Lemma 2, the following inequalities hold for all  $t \in [0, 1]$  and  $x \in [a, \frac{3a+b}{4}]$ .

$$\begin{aligned}
(2.25) \quad &f(ta + (1-t)x) + f\left(ta + (1-t)\left(\frac{3a+b}{2} - x\right)\right) \\
&\leq f(a) + f\left(ta + (1-t)\frac{a+b}{2}\right)
\end{aligned}$$

holds when  $A = a$ ,  $C = ta + (1-t)x$ ,  $D = ta + (1-t)\left(\frac{3a+b}{2} - x\right)$  and  $B = ta + (1-t)\frac{a+b}{2}$  in Lemma 2.

$$\begin{aligned}
(2.26) \quad &f\left(tb + (1-t)\left(\frac{b-a}{2} + x\right)\right) + f(tb + (1-t)(a+b-x)) \\
&\leq f\left(tb + (1-t)\frac{a+b}{2}\right) + f(b).
\end{aligned}$$

holds when  $A = tb + (1-t)\frac{a+b}{2}$ ,  $C = tb + (1-t)\left(\frac{b-a}{2} + x\right)$ ,  $D = tb + (1-t)(a+b-x)$  and  $B = b$  in Lemma 2. Multiplying the inequalities (2.25) – (2.26) by  $g(2x-a)$  and integrating them over  $x$  on  $[a, \frac{3a+b}{4}]$  and using (2.24), we have

$$(2.27) \quad N(t) \leq \frac{1}{2} \left[ \frac{f(a) + f(b)}{2} + G(t) \right] \int_a^b g(x) dx$$

for all  $t \in [0, 1]$ . Using (2.27), we derive the second inequality of (2.3).

Again, using Lemma 2, we have

$$\begin{aligned}
(2.28) \quad &f\left(ta + (1-t)\frac{a+b}{2}\right) + f\left(tb + (1-t)\frac{a+b}{2}\right) \\
&\leq f(ta + (1-t)x) + f(tb + (1-t)(a+b-x))
\end{aligned}$$

for all  $t \in [0, 1]$  and  $x \in [a, \frac{a+b}{2}]$ . Multiplying the inequality (2.28) by  $g(2x-a)$ , integrating both sides over  $x$  on  $[a, \frac{a+b}{2}]$  and using (2.23), we derive the first inequality of (2.3).

This proves (2.3).

(3) Integrating by parts, we have

$$(2.29) \quad \frac{1}{b-a} \int_a^{\frac{a+b}{2}} [(a-x)f'(x) + (x-a)f'(a+b-x)] dx \\ = \frac{1}{b-a} \int_a^b f(x) dx - f\left(\frac{a+b}{2}\right).$$

Using substitution rules for integration, we have the following identity

$$(2.30) \quad \frac{1}{b-a} \int_a^b f(x) dx = \frac{1}{b-a} \int_a^{\frac{a+b}{2}} [f(x) + f(a+b-x)] dx.$$

Now, using the convexity of  $f$  and  $g(x) \geq 0$  on  $[a, b]$ , the inequality

$$\begin{aligned} & [f(ta + (1-t)x) - f(x)]g(x) \\ & \quad + [f(tb + (1-t)(a+b-x)) - f(a+b-x)]g(x) \\ & \geq t(a-x)f'(x)g(x) + t(x-a)f'(a+b-x)g(x) \\ & = t(x-a)[f'(a+b-x) - f'(x)]g(x) \\ & \geq t(x-a)[f'(a+b-x) - f'(x)] \inf_{x \in [a,b]} g(x) \end{aligned}$$

holds for all  $t \in [0, 1]$  and  $x \in [a, \frac{a+b}{2}]$ . Integrating the above inequality over  $x$  on  $[a, \frac{a+b}{2}]$ , dividing both sides by  $(b-a)$  and using (1.1), (2.20), (2.29) and (2.30), we derive (2.4).

On the other hand, we have

$$\begin{aligned} \frac{f(a) - f\left(\frac{a+b}{2}\right)}{2} \int_a^b g(x) dx & \leq \frac{1}{2} \left(a - \frac{a+b}{2}\right) f'(a) \int_a^b g(x) dx \\ & = \frac{a-b}{4} f'(a) \int_a^b g(x) dx \end{aligned}$$

and

$$\begin{aligned} \frac{f(b) - f\left(\frac{a+b}{2}\right)}{2} \int_a^b g(x) dx & \leq \frac{1}{2} \left(b - \frac{a+b}{2}\right) f'(b) \int_a^b g(x) dx \\ & = \frac{b-a}{4} f'(b) \int_a^b g(x) dx \end{aligned}$$

and taking their sum we obtain:

$$(2.31) \quad \left[ \frac{f(a) + f(b)}{2} - f\left(\frac{a+b}{2}\right) \right] \int_a^b g(x) dx \\ \leq \frac{(f'(b) - f'(a))(b-a)}{4} \int_a^b g(x) dx.$$

Finally, (2.5) – (2.10) follow from (1.6), (1.7), (1.9), (1.10), (2.2) and (2.31).

This completes the proof.  $\square$

Let  $g(x) = \frac{1}{b-a}$  ( $x \in [a, b]$ ). Then the following Hermite-Hadamard-type inequalities, which are also given in [14], are natural consequences of Theorem 5.

**Corollary 6.** *Let  $f, G, H, L, P$  be defined as above. Then:*

(1) *The inequality*

$$\begin{aligned} \frac{1}{b-a} \int_a^b f(x) dx &\leq \frac{2}{b-a} \int_{[a, \frac{3a+b}{4}] \cup [\frac{a+3b}{4}, b]} f(x) dx \\ &\leq \int_0^1 P(t) dt \\ &\leq \frac{1}{2} \left[ \frac{1}{b-a} \int_a^b f(x) dx + \frac{f(a) + f(b)}{2} \right] \end{aligned}$$

*holds.*

(2) *The inequalities*

$$L(t) \leq P(t) \leq \frac{1-t}{b-a} \int_a^b f(x) dx + t \cdot \frac{f(a) + f(b)}{2} \leq \frac{f(a) + f(b)}{2}$$

*and*

$$0 \leq P(t) - G(t) \leq \frac{f(a) + f(b)}{2} - P(t)$$

*hold for all  $t \in [0, 1]$ .*

(3) *If  $f$  is differentiable on  $[a, b]$ , then we have the inequalities*

$$\begin{aligned} 0 &\leq t \left[ \frac{1}{b-a} \int_a^b f(x) dx - f\left(\frac{a+b}{2}\right) \right] \\ &\leq P(t) - \frac{1}{b-a} \int_a^b f(x) dx; \end{aligned}$$

$$0 \leq P(t) - f\left(\frac{a+b}{2}\right) \leq \frac{(f'(b) - f'(a))(b-a)}{4};$$

$$0 \leq L(t) - H(t) \leq \frac{(f'(b) - f'(a))(b-a)}{4};$$

$$0 \leq P(t) - L(t) \leq \frac{(f'(b) - f'(a))(b-a)}{4}$$

*and*

$$0 \leq P(t) - H(t) \leq \frac{(f'(b) - f'(a))(b-a)}{4}$$

*for all  $t \in [0, 1]$ .*

**Remark 7.** *In Theorem 5, the inequality (2.1) gives a new refinement of the Fejér inequality (1.2).*

**Remark 8.** *In Theorem 5, the inequality (2.2) refines the Fejér-type inequality (1.9).*

In the next theorem, we point out some inequalities for the functions  $G, Q, H_g, P_g, S_g$  considered above:

**Theorem 9.** *Let  $f, g, G, Q, H_g, P_g, S_g$  be defined as above. Then:*

(1) *The inequalities*

$$(2.32) \quad \begin{aligned} H_g(t) &\leq Q(t) \int_a^b g(x) dx \\ &\leq \frac{f(a) + f(b)}{2} \int_a^b g(x) dx \quad \left( t \in \left[0, \frac{1}{3}\right] \right) \end{aligned}$$

and

$$(2.33) \quad \begin{aligned} f\left(\frac{a+b}{2}\right) \int_a^b g(x) dx &\leq Q(t) \int_a^b g(x) dx \\ &\leq P_g(t) \quad \left( t \in \left[\frac{1}{3}, 1\right] \right) \end{aligned}$$

hold for all  $t \in [0, 1]$ .

(2) *The inequality*

$$(2.34) \quad \begin{aligned} 0 &\leq S_g(t) - G(t) \int_a^b g(x) dx \\ &\leq \frac{1}{2} \left[ \frac{f(a) + f(b)}{2} + Q(t) \right] \int_a^b g(x) dx - S_g(t) \end{aligned}$$

holds for all  $t \in [0, 1]$ .

*Proof.* (1) We discuss the following two cases.

**Case 1.**  $t \in \left[0, \frac{1}{3}\right]$ .

Using substitution rules for integration and the hypothesis of  $g$ , we have the following identity

$$(2.35) \quad \begin{aligned} H(t) = \int_a^{\frac{a+b}{2}} &\left[ f\left(tx + (1-t)\frac{a+b}{2}\right) \right. \\ &\left. + f\left(t(a+b-x) + (1-t)\frac{a+b}{2}\right) \right] g(x) dx. \end{aligned}$$

If we choose  $A = (1-t)a + tb$ ,  $C = tx + (1-t)\frac{a+b}{2}$ ,  $D = t(a+b-x) + (1-t)\frac{a+b}{2}$  and  $B = ta + (1-t)b$  in Lemma 2, then the inequality

$$(2.36) \quad \begin{aligned} f\left(tx + (1-t)\frac{a+b}{2}\right) + f\left(t(a+b-x) + (1-t)\frac{a+b}{2}\right) \\ \leq f((1-t)a + tb) + f(ta + (1-t)b) \end{aligned}$$

holds for all  $t \in \left[0, \frac{1}{3}\right]$  and  $x \in \left[a, \frac{a+b}{2}\right]$ . Multiplying the inequality (2.36) by  $g(x)$ , integrating both sides over  $x$  on  $\left[a, \frac{a+b}{2}\right]$  and using identity (2.35), we derive the first inequality of (2.32). From Lemma 4, we have

$$\sup_{t \in \left[0, \frac{1}{3}\right]} Q(t) = \frac{f(a) + f(b)}{2}.$$

Then the second inequality of (2.32) can be obtained. This proves (2.32).

**Case 2.**  $t \in \left[\frac{1}{3}, 1\right]$ .

If we choose  $A = ta + (1-t)x$ ,  $C = ta + (1-t)b$ ,  $D = (1-t)a + tb$  and  $B = tb + (1-t)(a+b-x)$  in Lemma 3, then the inequality

$$(2.37) \quad f(ta + (1-t)b) + f(tb + (1-t)a) \\ \leq f(ta + (1-t)x) + f(tb + (1-t)(a+b-x))$$

holds for all  $t \in [\frac{1}{3}, 1]$  and  $x \in [a, \frac{a+b}{2}]$ . Multiplying the inequality (2.37) by  $g(x)$ , integrating both sides over  $x$  on  $[a, \frac{a+b}{2}]$  and using identity (2.20), we obtain the second inequality of (2.33). From Lemma 4, we have

$$\inf_{t \in [\frac{1}{3}, 1]} Q(t) = f\left(\frac{a+b}{2}\right).$$

Then the first inequality of (2.33) can be obtained. This proves (2.33).

(2) Using substitution rules for integration and the hypothesis of  $g$ , we have the following identity

$$(2.38) \quad 2S_g(t) = \int_a^{\frac{a+b}{2}} [f(ta + (1-t)x) + f(tb + (1-t)x)] g(2x-a) dx \\ + \int_{\frac{a+b}{2}}^b [f(ta + (1-t)x) + f(tb + (1-t)x)] g(2x-b) dx \\ = \int_a^{\frac{a+b}{2}} [f(ta + (1-t)x) + f(tb + (1-t)x) \\ + f(ta + (1-t)(a+b-x)) + f(tb + (1-t)(a+b-x))] \\ \times g(2x-a) dx \\ = \int_a^{\frac{3a+b}{4}} \left[ f(ta + (1-t)x) + f\left(ta + (1-t)\left(\frac{3a+b}{2} - x\right)\right) \right. \\ \left. + f\left(ta + (1-t)\left(\frac{b-a}{2} + x\right)\right) + f(ta + (1-t)(a+b-x)) \right. \\ \left. + f(tb + (1-t)x) + f\left(tb + (1-t)\left(\frac{3a+b}{2} - x\right)\right) \right. \\ \left. + f\left(tb + (1-t)\left(\frac{b-a}{2} + x\right)\right) + f(tb + (1-t)(a+b-x)) \right] \\ \times g(2x-a) dx$$

for all  $t \in [0, 1]$ .

By Lemma 2, the following inequalities hold for all  $t \in [0, 1]$  and  $x \in [a, \frac{3a+b}{4}]$ .

$$(2.39) \quad f(ta + (1-t)x) + f\left(ta + (1-t)\left(\frac{3a+b}{2} - x\right)\right) \\ \leq f(a) + f\left(ta + (1-t)\frac{a+b}{2}\right)$$

holds when  $A = a$ ,  $C = ta + (1-t)x$ ,  $D = ta + (1-t)\left(\frac{3a+b}{2} - x\right)$  and  $B = ta + (1-t)\frac{a+b}{2}$  in Lemma 2.

$$(2.40) \quad f\left(ta + (1-t)\left(\frac{b-a}{2} + x\right)\right) + f(ta + (1-t)(a+b-x)) \\ \leq f\left(ta + (1-t)\frac{a+b}{2}\right) + f(ta + (1-t)b)$$

holds when  $A = ta + (1-t)\frac{a+b}{2}$ ,  $C = ta + (1-t)\left(\frac{b-a}{2} + x\right)$ ,  $D = ta + (1-t)(a+b-x)$  and  $B = ta + (1-t)b$  in Lemma 2.

$$(2.41) \quad f(tb + (1-t)x) + f\left(tb + (1-t)\left(\frac{3a+b}{2} - x\right)\right) \\ \leq f(tb + (1-t)a) + f\left(tb + (1-t)\frac{a+b}{2}\right)$$

holds when  $A = tb + (1-t)a$ ,  $C = tb + (1-t)x$ ,  $D = tb + (1-t)\left(\frac{3a+b}{2} - x\right)$  and  $B = tb + (1-t)\frac{a+b}{2}$  in Lemma 2.

$$(2.42) \quad f\left(tb + (1-t)\left(\frac{b-a}{2} + x\right)\right) + f(tb + (1-t)(a+b-x)) \\ \leq f\left(tb + (1-t)\frac{a+b}{2}\right) + f(b)$$

holds when  $A = tb + (1-t)\frac{a+b}{2}$ ,  $C = tb + (1-t)\left(\frac{b-a}{2} + x\right)$ ,  $D = tb + (1-t)(a+b-x)$  and  $B = b$  in Lemma 2. Multiplying the inequalities (2.39) – (2.42) by  $g(2x-a)$ , integrating them over  $x$  on  $[a, \frac{3a+b}{4}]$  and using identity (2.38), we have

$$(2.43) \quad 2S_g(t) \leq G(t) \int_a^b g(x) dx + \frac{1}{2} \left[ \frac{f(a) + f(b)}{2} + Q(t) \right] \int_a^b g(x) dx$$

for all  $t \in [0, 1]$ . Using (1.10) and (2.43), we derive (2.34). This completes the proof.  $\square$

Let  $g(x) = \frac{1}{b-a}(x \in [a, b])$ . Then the following Hermite-Hadamard-type inequalities, which are given in [14], are natural consequences of Theorem 9.

**Corollary 10.** *Let  $f, G, H, L, P$  be defined as above. Then:*

(1) *The inequalities*

$$H(t) \leq Q(t) \leq \frac{f(a) + f(b)}{2} \quad \left(t \in \left[0, \frac{1}{3}\right]\right)$$

and

$$f\left(\frac{a+b}{2}\right) \leq Q(t) \leq P(t) \quad \left(t \in \left[\frac{1}{3}, 1\right]\right)$$

hold for all  $t \in [0, 1]$ .

(2) *The inequality*

$$0 \leq L(t) - G(t) \leq \frac{1}{2} \left[ \frac{f(a) + f(b)}{2} + Q(t) \right] - L(t)$$

holds for all  $t \in [0, 1]$ .

The following Fejér-type inequalities are natural consequences of Theorems A – B, E – I, 5, 9 and Lemma 4 and we shall omit their proofs.

**Theorem 11.** *Let  $f, g, G, H_g, P_g, I, L_g, S_g$  be defined as above.*

$$\begin{aligned} f\left(\frac{a+b}{2}\right) \int_a^b g(x) dx &\leq H_g(t) \leq G(t) \int_a^b g(x) dx \leq S_g(t) \\ &\leq (1-t) \int_a^b \frac{1}{2} \left[ f\left(\frac{x+a}{2}\right) + f\left(\frac{x+b}{2}\right) \right] g(x) dx \\ &\quad + t \cdot \frac{f(a)+f(b)}{2} \int_a^b g(x) dx \\ &\leq \frac{f(a)+f(b)}{2} \int_a^b g(x) dx \end{aligned}$$

and

$$\begin{aligned} f\left(\frac{a+b}{2}\right) \int_a^b g(x) dx &\leq I(t) \leq G(t) \int_a^b g(x) dx \\ &\leq L_g(t) \leq P_g(t) \\ &\leq (1-t) \int_a^b f(x) g(x) dx + t \cdot \frac{f(a)+f(b)}{2} \int_a^b g(x) dx \\ &\leq \frac{f(a)+f(b)}{2} \int_a^b g(x) dx. \end{aligned}$$

**Theorem 12.** *Let  $f, g, G, Q, H_g, I$  be defined as above. Then, for all  $t \in [0, \frac{1}{4}]$ , we have*

$$\begin{aligned} f\left(\frac{a+b}{2}\right) \int_a^b g(x) dx &\leq H_g(t) \leq H_g(2t) \leq G(2t) \int_a^b g(x) dx \\ &\leq Q(t) \int_a^b g(x) dx \leq \frac{f(a)+f(b)}{2} \int_a^b g(x) dx \end{aligned}$$

and

$$\begin{aligned} f\left(\frac{a+b}{2}\right) \int_a^b g(x) dx &\leq I(t) \leq I(2t) \leq G(2t) \int_a^b g(x) dx \\ &\leq Q(t) \int_a^b g(x) dx \leq \frac{f(a)+f(b)}{2} \int_a^b g(x) dx. \end{aligned}$$

**Theorem 13.** *Let  $f, g, G, Q, H_g, P_g, L_g, S_g$  be defined as above. Then, for all  $t \in [\frac{1}{4}, \frac{1}{3}]$ , we have*

$$\begin{aligned} f\left(\frac{a+b}{2}\right) \int_a^b g(x) dx &\leq H_g(t) \leq Q(t) \int_a^b g(x) dx \leq G(2t) \int_a^b g(x) dx \\ &\leq L_g(2t) \leq P_g(2t) \\ &\leq (1-2t) \int_a^b f(x) g(x) dx + 2t \cdot \frac{f(a)+f(b)}{2} \int_a^b g(x) dx \\ &\leq \frac{f(a)+f(b)}{2} \int_a^b g(x) dx \end{aligned}$$



and

$$\begin{aligned}
f\left(\frac{a+b}{2}\right) \int_a^b g(x) dx &\leq H_g(t) \leq Q(t) \int_a^b g(x) dx \\
&\leq G(2t) \int_a^b g(x) dx \leq S_g(2t) \\
&\leq (1-2t) \int_a^b \frac{1}{2} \left[ f\left(\frac{x+a}{2}\right) + f\left(\frac{x+b}{2}\right) \right] g(x) dx \\
&\quad + 2t \cdot \frac{f(a)+f(b)}{2} \int_a^b g(x) dx \\
&\leq \frac{f(a)+f(b)}{2} \int_a^b g(x) dx.
\end{aligned}$$

**Theorem 14.** Let  $f, g, G, Q, P_g, L_g, S_g$  be defined as above. Then, for all  $t \in [\frac{1}{3}, \frac{1}{2}]$ , we have

$$\begin{aligned}
f\left(\frac{a+b}{2}\right) \int_a^b g(x) d &\leq Q(t) \int_a^b g(x) dx \\
&\leq G(2t) \int_a^b g(x) dx \leq L_g(2t) \leq P_g(2t) \\
&\leq (1-2t) \int_a^b f(x) g(x) dx + 2t \cdot \frac{f(a)+f(b)}{2} \int_a^b g(x) dx \\
&\leq \frac{f(a)+f(b)}{2} \int_a^b g(x) dx;
\end{aligned}$$

$$\begin{aligned}
f\left(\frac{a+b}{2}\right) \int_a^b g(x) d &\leq Q(t) \int_a^b g(x) dx \\
&\leq G(2t) \int_a^b g(x) dx \leq S_g(2t) \\
&\leq (1-2t) \int_a^b \frac{1}{2} \left[ f\left(\frac{x+a}{2}\right) + f\left(\frac{x+b}{2}\right) \right] g(x) dx \\
&\quad + 2t \cdot \frac{f(a)+f(b)}{2} \int_a^b g(x) dx \\
&\leq \frac{f(a)+f(b)}{2} \int_a^b g(x) dx
\end{aligned}$$

and

$$\begin{aligned}
f\left(\frac{a+b}{2}\right) \int_a^b g(x) dx &\leq Q(t) \int_a^b g(x) dx \leq P_g(t) \leq P_g(2t) \\
&\leq (1-2t) \int_a^b f(x) g(x) dx + 2t \cdot \frac{f(a)+f(b)}{2} \int_a^b g(x) dx \\
&\leq \frac{f(a)+f(b)}{2} \int_a^b g(x) dx.
\end{aligned}$$

**Theorem 15.** *Let  $f, g, G, Q, P_g, L_g, S_g$  be defined as above. Then, for all  $t \in [\frac{1}{2}, \frac{2}{3}]$ , we have*

$$\begin{aligned} f\left(\frac{a+b}{2}\right) \int_a^b g(x) dx &\leq Q(t) \int_a^b g(x) dx \leq G(2(1-t)) \int_a^b g(x) dx \\ &\leq L_g(2(1-t)) \leq P_g(2(1-t)) \\ &\leq (2t-1) \int_a^b f(x)g(x) dx + 2(1-t) \cdot \frac{f(a)+f(b)}{2} \int_a^b g(x) dx \\ &\leq \frac{f(a)+f(b)}{2} \int_a^b g(x) dx \end{aligned}$$

and

$$\begin{aligned} f\left(\frac{a+b}{2}\right) \int_a^b g(x) dx &\leq Q(t) \int_a^b g(x) dx \\ &\leq G(2(1-t)) \int_a^b g(x) dx \leq S_g(2(1-t)) \\ &\leq (2t-1) \int_a^b \frac{1}{2} \left[ f\left(\frac{x+a}{2}\right) + f\left(\frac{x+b}{2}\right) \right] g(x) dx \\ &\quad + 2(1-t) \cdot \frac{f(a)+f(b)}{2} \int_a^b g(x) dx \\ &\leq \frac{f(a)+f(b)}{2} \int_a^b g(x) dx. \end{aligned}$$

**Theorem 16.** *Let  $f, g, G, Q, H_g, P_g, L_g, S_g$  be defined as above. Then, for all  $t \in [\frac{2}{3}, \frac{3}{4}]$ , we have*

$$\begin{aligned} f\left(\frac{a+b}{2}\right) \int_a^b g(x) dx &\leq Q(t) \int_a^b g(x) dx \\ &\leq G(2(1-t)) \int_a^b g(x) dx \\ &\leq G(t) \int_a^b g(x) dx \leq L_g(t) \leq P_g(t) \\ &\leq (1-t) \int_a^b f(x)g(x) dx + t \cdot \frac{f(a)+f(b)}{2} \int_a^b g(x) dx \\ &\leq \frac{f(a)+f(b)}{2} \int_a^b g(x) dx \end{aligned}$$

and

$$\begin{aligned} f\left(\frac{a+b}{2}\right) \int_a^b g(x) dx &\leq Q(t) \int_a^b g(x) dx \leq G(2(1-t)) \int_a^b g(x) dx \\ &\leq G(t) \int_a^b g(x) dx \leq S_g(t) \end{aligned}$$

$$\begin{aligned}
&\leq (1-t) \int_a^b \frac{1}{2} \left[ f\left(\frac{x+a}{2}\right) + f\left(\frac{x+b}{2}\right) \right] g(x) dx \\
&\quad + t \cdot \frac{f(a)+f(b)}{2} \int_a^b g(x) dx \\
&\leq \frac{f(a)+f(b)}{2} \int_a^b g(x) dx.
\end{aligned}$$

**Theorem 17.** *Let  $f, g, G, Q, H_g, P_g, I, S_g$  be defined as above. Then, for all  $t \in [\frac{3}{4}, 1]$ , we have*

$$\begin{aligned}
f\left(\frac{a+b}{2}\right) \int_a^b g(x) dx &\leq H_g(2(1-t)) \leq G(2(1-t)) \int_a^b g(x) dx \\
&\leq Q(t) \int_a^b g(x) dx \leq P_g(t) \\
&\leq \frac{1-t}{b-a} \int_a^b f(x)g(x) dx + t \cdot \frac{f(a)+f(b)}{2} \int_a^b g(x) dx \\
&\leq \frac{f(a)+f(b)}{2} \int_a^b g(x) dx
\end{aligned}$$

and

$$\begin{aligned}
f\left(\frac{a+b}{2}\right) \int_a^b g(x) dx &\leq I(2(1-t)) \leq G(2(1-t)) \int_a^b g(x) dx \\
&\leq Q(t) \int_a^b g(x) dx \leq P_g(t) \\
&\leq \frac{1-t}{b-a} \int_a^b f(x)g(x) dx + t \cdot \frac{f(a)+f(b)}{2} \int_a^b g(x) dx \\
&\leq \frac{f(a)+f(b)}{2} \int_a^b g(x) dx.
\end{aligned}$$

Let  $g(x) = \frac{1}{b-a}$  ( $x \in [a, b]$ ). Then the following Hermite-Hadamard-type inequalities are natural consequences of Theorems 11 – 17, which are given in [14].

**Corollary 18.** *Let  $f, Q, G, H, P, L$  be defined as above. Then we have:*

(1) *For all  $t \in [0, \frac{1}{4}]$  one has the inequality*

$$f\left(\frac{a+b}{2}\right) \leq H(t) \leq H(2t) \leq G(2t) \leq Q(t) \leq \frac{f(a)+f(b)}{2}.$$

(2) *For all  $t \in [\frac{1}{4}, \frac{1}{3}]$  one has the inequality*

$$\begin{aligned}
f\left(\frac{a+b}{2}\right) &\leq H(t) \leq Q(t) \leq G(2t) \leq L(2t) \leq P(2t) \\
&\leq \frac{1-2t}{b-a} \int_a^b f(x) dx + 2t \cdot \frac{f(a)+f(b)}{2} \\
&\leq \frac{f(a)+f(b)}{2}.
\end{aligned}$$

(3) For all  $t \in [\frac{1}{3}, \frac{1}{2}]$  one has the inequalities

$$\begin{aligned} f\left(\frac{a+b}{2}\right) &\leq Q(t) \leq G(2t) \leq L(2t) \leq P(2t) \\ &\leq \frac{1-2t}{b-a} \int_a^b f(x) dx + 2t \cdot \frac{f(a)+f(b)}{2} \\ &\leq \frac{f(a)+f(b)}{2} \end{aligned}$$

and

$$\begin{aligned} f\left(\frac{a+b}{2}\right) &\leq Q(t) \leq P(t) \leq P(2t) \\ &\leq \frac{1-2t}{b-a} \int_a^b f(x) dx + 2t \cdot \frac{f(a)+f(b)}{2} \\ &\leq \frac{f(a)+f(b)}{2}. \end{aligned}$$

(4) For all  $t \in [\frac{1}{2}, \frac{2}{3}]$  one has the inequality

$$\begin{aligned} f\left(\frac{a+b}{2}\right) &\leq Q(t) \leq G(2(1-t)) \leq L(2(1-t)) \leq P(2(1-t)) \\ &\leq \frac{2t-1}{b-a} \int_a^b f(x) dx + 2(1-t) \cdot \frac{f(a)+f(b)}{2} \\ &\leq \frac{f(a)+f(b)}{2}. \end{aligned}$$

(5) For all  $t \in [\frac{2}{3}, \frac{3}{4}]$  one has the inequality

$$\begin{aligned} f\left(\frac{a+b}{2}\right) &\leq Q(t) \leq G(2(1-t)) \leq G(t) \leq L(t) \leq P(t) \\ &\leq \frac{1-t}{b-a} \int_a^b f(x) dx + t \cdot \frac{f(a)+f(b)}{2} \leq \frac{f(a)+f(b)}{2}. \end{aligned}$$

(6) For all  $t \in [\frac{3}{4}, 1]$  one has the inequality

$$\begin{aligned} f\left(\frac{a+b}{2}\right) &\leq H(2(1-t)) \leq G(2(1-t)) \leq Q(t) \leq P(t) \\ &\leq \frac{1-t}{b-a} \int_a^b f(x) dx + t \cdot \frac{f(a)+f(b)}{2} \leq \frac{f(a)+f(b)}{2}. \end{aligned}$$

#### REFERENCES

- [1] J. Hadamard, Étude sur les propriétés des fonctions entières en particulier d'une fonction considérée par Riemann, *J. Math. Pures Appl.* 58 (1893), 171-215.
- [2] S. S. Dragomir, Two mappings in Connection to Hadamard's Inequalities, *J. Math. Anal. Appl.* 167 (1992), 49-56.
- [3] S. S. Dragomir, A Refinement of Hadamard's Inequality for Isotonic Linear Functionals, *Tamkang. J. Math.*
- [4] S. S. Dragomir, On the Hadamard's Inequality for Convex on the Co-ordinates in a Rectangle from the Plane, *Taiwanese J. Math.*, 5 (4) (2001), 775-788.
- [5] S. S. Dragomir, Further Properties of Some Mappings Associated with Hermite-Hadamard Inequalities, *Tamkang. J. Math.* 34 (1) (2003), 45-57.

- [6] S. S. Dragomir, Y.-J. Cho and S.-S. Kim, Inequalities of Hadamard's type for Lipschitzian Mappings and their Applications, *J. Math. Anal. Appl.* 245 (2000),489-501.
- [7] S. S. Dragomir, D. S. Milošević and József Sándor, On Some Refinements of Hadamard's Inequalities and Applications, *Univ. Belgrad. Publ. Elek. Fak. Sci. Math.* 4 (1993), 3-10.
- [8] L. Fejér, Über die Fourierreihen, II, *Math. Naturwiss. Anz Ungar. Akad. Wiss.* 24 (1906),369-390.(In Hungarian).
- [9] D.-Y. Hwang, K.-L. Tseng and G.-S. Yang, Some Hadamard's Inequalities for Co-ordinated Convex Functions in a Rectangle from the Plane, *Taiwanese J. Math.*, 11 (1) (2007), 63-73.
- [10] K.-L. Tseng, S.-R. Hwang and S. S. Dragomir, On Some New Inequalities of Hermite-Hadamard-Fejér Type Involving Convex Functions, *Demonstratio Math.* XL (1) (2007), 51-64.
- [11] K.-L. Tseng, S.-R. Hwang and S. S. Dragomir, Fejér-type Inequalities (I). (Submitted), Preprint *RGMA Res. Rep. Coll.* 12(2009), No.4, Article 5. [Online <http://www.staff.vu.edu.au/RGMA/v12n4.asp>].
- [12] K.-L. Tseng, S.-R. Hwang and S. S. Dragomir, Fejér-type Inequalities (II). (Submitted) Preprint *RGMA Res. Rep. Coll.* 12(2009), Supplement, Article 15.[Online [http://www.staff.vu.edu.au/RGMA/v12\(E\).asp](http://www.staff.vu.edu.au/RGMA/v12(E).asp)].
- [13] K.-L. Tseng, S.-R. Hwang and S. S. Dragomir, Some Companions of Fejer's Inequality for Convex Functions. (Submitted)*RGMA Res. Rep. Coll.* 12(2009), Supplement, Article 19.[Online [http://www.staff.vu.edu.au/RGMA/v12\(E\).asp](http://www.staff.vu.edu.au/RGMA/v12(E).asp)].
- [14] K.-L. Tseng, G.-S. Yang and K.-C. Hsu, On Some Inequalities of Hadamard's Type and Applications, *Taiwanese J. Math.*, 13.
- [15] G.-S. Yang and M.-C. Hong, A Note on Hadamard's Inequality, *Tamkang. J. Math.* 28 (1) (1997), 33-37.
- [16] G.-S. Yang and K.-L. Tseng, On Certain Integral Inequalities Related to Hermite-Hadamard Inequalities, *J. Math. Anal. Appl.* 239 (1999), 180-187.
- [17] G.-S. Yang and K.-L. Tseng, Inequalities of Hadamard's Type for Lipschitzian Mappings, *J. Math. Anal. Appl.* 260 (2001),230-238.
- [18] G.-S. Yang and K.-L. Tseng, On Certain Multiple Integral Inequalities Related to Hermite-Hadamard Inequalities, *Utilitas Math.*, 62 (2002), 131-142.
- [19] G.-S. Yang and K.-L. Tseng, Inequalities of Hermite-Hadamard-Fejér Type for Convex Functions and Lipschitzian Functions, *Taiwanese J. Math.*, 7 (3) (2003), 433-440.

DEPARTMENT OF MATHEMATICS, ALETHEIA UNIVERSITY, TAMSUI, TAIWAN 25103.  
*E-mail address:* [kltseng@email.au.edu.tw](mailto:kltseng@email.au.edu.tw)

CHINA UNIVERSITY OF SCIENCE AND TECHNOLOGY, NANKANG, TAIPEI, TAIWAN 11522  
*E-mail address:* [hsru@cc.cust.edu.tw](mailto:hsru@cc.cust.edu.tw)

SCHOOL OF ENGINEERING AND SCIENCE, VICTORIA UNIVERSITY, PO BOX 14428, MELBOURNE CITY MC, VICTORIA 8001, AUSTRALIA.

*E-mail address:* [sever.dragomir@vu.edu.au](mailto:sever.dragomir@vu.edu.au)  
*URL:* <http://www.staff.vu.edu.au/RGMA/dragomir/>