



VICTORIA UNIVERSITY
MELBOURNE AUSTRALIA

A Subadditive Property of the Digamma Function

This is the Published version of the following publication

Gao, Peng (2005) A Subadditive Property of the Digamma Function. Research report collection, 8 (3).

The publisher's official version can be found at

Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/18103/>

A SUBADDITIVE PROPERTY OF THE DIGAMMA FUNCTION

PENG GAO

ABSTRACT. We prove that the function $x \mapsto \psi(a + e^x)$ ($a > 0$) is subadditive on $(-\infty, +\infty)$ if and only if $a \geq c_0$, where c_0 is the only positive zero of $\psi(x)$.

1. INTRODUCTION

Euler's gamma function for $x > 0$ is defined as

$$\Gamma(x) = \int_0^{\infty} t^x e^{-t} \frac{dt}{t}.$$

The digamma (or psi) function $\psi(x)$ is defined as the logarithmic derivative of $\Gamma(x)$. It has the following series representation(see, for example, [7, (1.8)]):

$$(1.1) \quad \psi(x) = -\gamma + \sum_{n=0}^{\infty} \left(\frac{1}{n+1} - \frac{1}{x+n} \right).$$

Here $\gamma = 0.57721\dots$ denotes Euler's constant. We further note the following asymptotic expression(see, for example, [2, (1.4)]):

$$(1.2) \quad \psi(x) = \log x + O\left(\frac{1}{x}\right).$$

There exist many inequalities for the gamma and digamma functions. For the recent developments in this area, we refer the reader to the articles [1]-[3], [7], [8] and the references therein.

A function f defined on a set D of real numbers is said to be subadditive on D if $f(x+y) \leq f(x) + f(y)$ for all $x, y \in D$ such that $x+y \in D$. If instead $f(xy) \leq f(x)f(y)$ for all $x, y \in D$ such that $xy \in D$, then $f(x)$ is said to be submultiplicative on D .

Inspired by (1.2) and a result of Gustavsson et al. [6], which asserts that if $a \geq 1$, then the function $f(x) = \log(a+x)$ is submultiplicative on $[0, +\infty)$ if and only if $a \geq e$, Alzer and Ruehr[3] proved the following submultiplicative property of the digamma function:

Theorem 1.1. *Let $a > 0$ be fixed. Then $\psi(a+x)$ as a function of x is submultiplicative on $[0, +\infty)$ if and only if $a \geq a_0 = 3.203171\dots$ (Here, a_0 denotes the only positive real number which satisfies $\psi(a_0) = 1$.)*

The above result asserts that for $a \geq a_0, x, y \geq 0$,

$$(1.3) \quad \psi(a+xy) \leq \psi(a+x)\psi(a+y).$$

We note here[5] that for $a > 0, x, y \geq 0$,

$$\psi(a) + \psi(a+x+y) \leq \psi(a+x) + \psi(a+y).$$

As $\psi(x)$ is an increasing function(see [5]), we obtain from the above that

$$(1.4) \quad \psi(a) + \psi(a+xy) \leq \psi(a+x) + \psi(a+y),$$

Date: August 18, 2005.

1991 Mathematics Subject Classification. Primary 33D05.

Key words and phrases. Digamma function, subadditive function.

for $a > 0, x, y \geq 0$, provided that $xy \leq x + y$ or $1 \leq 1/x + 1/y$. In particular, when $x \leq 1$ or $y \leq 1$, inequality (1.4) is valid and this would then give a refinement of (1.3) when $a \geq a_0$ since we now have

$$1 + \psi(a + xy) \leq \psi(a) + \psi(a + xy) \leq \psi(a + x) + \psi(a + y),$$

and that

$$\psi(a + x) + \psi(a + y) - 1 \leq \psi(a + x)\psi(a + y).$$

We point out here (1.4) does not hold in general. In fact, it follows from (1.2) and (1.4) that when $x \rightarrow +\infty$,

$$\psi(a) + \log y \leq \psi(a + y),$$

and the above inequality does not hold for $y > 1$ if we take $a \rightarrow +\infty$, in view of (1.2) again. Similarly, it is not possible to compare $\psi(a)\psi(a + xy)$ and $\psi(a + x)\psi(a + y)$ for $a \geq a_0, x, y \geq 0$ in general. Certainly when we fix a and set $x = y \rightarrow +\infty$, we will have by (1.2),

$$\psi(a)\psi(a + xy) < \psi(a + x)\psi(a + y),$$

and the above inequality is reversed when we set $x = y = a \rightarrow +\infty$.

It is now interesting to ask whether the following inequality holds for all $x, y \geq 0$:

$$(1.5) \quad \psi(a + xy) \leq \psi(a + x) + \psi(a + y),$$

for a larger than some constant. In view of (1.2), this can be thought as an analogue of the property $\log(xy) = \log x + \log y$, $x, y > 0$ and the easily checked fact: $\log(a + xy) \leq \log(a + x) + \log(a + y)$, $a \geq 1, x, y \geq 0$. We may also interpret (1.5) as asserting that the function $x \mapsto \psi(a + e^x)$ is subadditive on $(-\infty, +\infty)$ for a larger than some constant.

By taking $x = y = 0$ in (1.5), we see that it is necessary to have $a \geq c_0$, where c_0 is the only positive zero of $\psi(x)$ and it is our goal in this paper to prove that this condition is also sufficient for (1.5) to hold.

2. A LEMMA

Lemma 2.1. *The function $f(x) = x\psi'(a + x)$ is strictly increasing on $[0, +\infty)$ for $a \geq 1$.*

Proof. We have

$$f'(x) = \psi'(a + x) + x\psi''(a + x).$$

We now use a method in [4] to show that $f'(x) > 0$ for $x \geq 0$. First note the following two asymptotic expressions([4, p. 2668]):

$$\psi'(x) = \frac{1}{x} + O\left(\frac{1}{x^2}\right), \psi''(x) = -\frac{1}{x^2} + O\left(\frac{1}{x^3}\right).$$

It follows that

$$\lim_{x \rightarrow +\infty} f'(x) = 0.$$

Hence it suffices to show that $f(x) - f(x + 1) > 0$. Using (1.1) we get

$$\begin{aligned} f(x) - f(x + 1) &= -\frac{1}{(x + a)^2} + \frac{2a}{(x + a)^3} + \sum_{n=1}^{\infty} \frac{2}{(n + x + a)^3} \\ &\geq -\frac{1}{(x + a)^2} + \sum_{n=0}^{\infty} \frac{2}{(n + x + a)^3} \\ &> -\frac{1}{(x + a)^2} + \sum_{n=0}^{\infty} \left(\frac{1}{(n + x + a)^2} - \frac{1}{(n + x + a + 1)^2} \right) = 0, \end{aligned}$$

where the last inequality follows from the inequality

$$\frac{2}{u^3} > \frac{1}{u^2} - \frac{1}{(1+u)^2}, \quad u > 0.$$

The proof is now complete. \square

3. MAIN RESULT

Theorem 3.1. *The function $x \mapsto \psi(a + e^x)$ ($a > 0$) is subadditive on $(-\infty, +\infty)$ if and only if $a \geq c_0$, where c_0 is the only positive zero of $\psi(x)$.*

Proof. From our discussions above, it suffices to show (1.5) holds for all $x, y \geq 0$ when $a \geq c_0$. If $x \leq 1$ or $y \leq 1$, then (1.5) follows from (1.4). Hence from now on we may assume that $x, y \geq 1$. In this case we define

$$f(x, y) = \psi(a + x) + \psi(a + y) - \psi(a + xy),$$

and note that $f(x, 1) = \psi(a + 1) > 0$. If $y > 1$, then as $c_0 > 1$ (since $\psi(1) = -\gamma$ by (1.1)), Lemma 2.1 implies that $\partial f / \partial x < 0$ so that by (1.2),

$$f(x, y) > \lim_{u \rightarrow +\infty} f(u, y) = \psi(a + y) - \log y.$$

Now that we have [1, (2.2)] for $x > 0$,

$$\frac{1}{2x} < \log x - \psi(x) < \frac{1}{x}.$$

It follows from this and the mean value theorem that

$$\psi(a + y) - \log y > \log(a + y) - \log y - \frac{1}{a + y} > \frac{a}{a + y} - \frac{1}{a + y} \geq 0.$$

From this we conclude that $f(x, y) \geq 0$ for $x, y \geq 0$ and this completes the proof. \square

REFERENCES

- [1] H. Alzer, On some inequalities for the gamma and psi functions, *Math. Comp.*, **66** (1997), 373–389.
- [2] H. Alzer, Sharp inequalities for the digamma and polygamma functions, *Forum Math.*, **16** (2004), 181–221.
- [3] H. Alzer and O. G. Ruehr, A submultiplicative property of the psi function, *J. Comput. Appl. Math.*, **101** (1999), 53–60.
- [4] Á. Elbert and A. Laforgia, On some properties of the gamma function, *Proc. Amer. Math. Soc.*, **128** (2000), 2667–2673.
- [5] P. Gao, Some monotonicity properties of the q -gamma Function, *RGMA Research Report Collection* **8**(3), Article 4, 2005.
- [6] J. Gustavsson, L. Maligranda and J. Peetre, A submultiplicative function, *Nederl. Akad. Wetensch. Indag. Math.*, **51** (1989), 435–442.
- [7] M.E.H. Ismail and M.E. Muldoon, Inequalities and monotonicity properties for gamma and q -gamma functions. In: *Approximation and computation, Internat. Ser. Numer. Math.*, **119**, Birkhäuser, Boston, 1994, 309–323.
- [8] S.-L. Qiu and M. Vuorinen, Some properties of the gamma and psi functions, with applications, *Math. Comp.*, **74** (2005), 723–742.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109
E-mail address: penggao@umich.edu