
VICTORIA UNIVERSITY
MELBOURNE AUSTRALIA

Approximation of pn by Hn

This is the Published version of the following publication

Alikhani, Akram and Hassani, Mehdi (2005) Approximation of pn by Hn. Research report collection, 8 (4).

The publisher's official version can be found at

Note that access to this version may require subscription.

Downloaded from VU Research Repository https://vuir.vu.edu.au/18117/

APPROXIMATION OF p_n BY H_n

AKRAM ALIKHANI
MEHDI HASSANI

Abstract. In this paper we introduce some bounds including $H_n = \sum_{k=1}^{n} \frac{1}{k}$, for p_n, n^{th} prime number. Then we observe that the Prime Number Theorem is equivalent with $p_n \sim nH_n$, when n tends to infinity.

1. Introduction

As usual, let p_n be the n^{th} prime. According to the Prime Number Theorem (PNT) [3], we know that:

$$p_n = n\log n + o(n\log n) \qquad (n \to \infty). \tag{1.1}$$

Also, we know that [1], if $H_n = \sum_{k=1}^{n} \frac{1}{k}$, then:

$$H_n = \log n + O(1) \qquad (n \to \infty). \tag{1.2}$$

So, considering (1.1) and (1.2), we obtain:

$$p_n = n(H_n + O(1)) + o(n\log n) = nH_n + o(n\log n) \qquad (n \to \infty).$$

Therefore, comparing p_n and nH_n seems to be a nice problem. To consider this problem, we need some bounds concerning p_n and H_n, which we recall them from literatures. About p_n, we have the following bounds [4]:

(1.3)

$$n\log n + n\log_2 n - n + n\frac{\log_2 n - 2.25}{\log n} \leq p_n \leq n\log n + n\log_2 n - n + n\frac{\log_2 n - 1.8}{\log n},$$

which left hand side of it holds true for every $n \geq 2$ and the right hand side of it holds true for every $n \geq 27076$, $\log_2 n$ means $\log\log n$ and base of all logarithms is e. Also, for H_n we have the following bounds [2]:

$$\gamma + \log(n + 0.5) < H_n \leq \gamma + \log(n - 1 + e^{1-\gamma}) \qquad (n \geq 1), \tag{1.4}$$

where γ is Euler constant. In this note, we search some bounds of the form $nH_n + e(n)$, which we will find $e(n)$ in both cases lower and upper bounds.

2. Inequalities of the form nH_n

Consider the following inequality:

$$nH_n + a(n) \leq p_n \leq nH_n + b(n). \tag{2.1}$$

Here, we try to find some suitable functions $a(n)$ and $b(n)$, such that (2.1) holds true.

1991 *Mathematics Subject Classification.* 11A41, 26D15.
Key words and phrases. Primes, Harmonic series.

Upper Bound. To find above mentioned upper bound, considering (1.4), we have:

$$n\left(\gamma+\log(n+0.5)\right)<nH_n\leq n\left(\gamma+\log(n-1+e^{1-\gamma})\right). \tag{2.2}$$

Using left hand side of above inequality, we have:

$$n\left(\gamma+\log(n+0.5)\right)+b(n)\leq nH_n+b(n),$$

and considering this inequality, with the right hand side of (1.3), we must have:

$$n\log n+n\log_2 n-n+n\frac{\log_2 n-1.8}{\log n}\leq n\left(\gamma+\log(n+0.5)\right)+b(n),$$

or equivalently,

$$n\log n+n\log_2 n-n+n\frac{\log_2 n-1.8}{\log n}-n\left(\gamma+\log(n+0.5)\right)\leq b(n),$$

Since, $b(n)$ is going to appears in upper bound for p_n , the best possible case is:

$$b(n)=n\log_2 n-n(1+\gamma)+n\left(\frac{\log_2 n-1.8}{\log n}-\log\left(1+\frac{0.5}{n}\right)\right). \tag{2.3}$$

Thus, we have:

$$p_n\leq nH_n+n\log_2 n-n(1+\gamma)+n\left(\frac{\log_2 n-1.8}{\log n}-\log\left(1+\frac{0.5}{n}\right)\right), \tag{2.4}$$

which holds for $n\geq 27076$.

Lower Bound. To find above mentioned lower bound, considering (2.2), we have:

$$nH_n+a(n)\leq n\gamma+n\log(n-1+e^{1-\gamma})+a(n).$$

Considering this inequality with the left hand side of (2.1) and the left hand side of (1.3), we must have:

$$n\gamma+n\log(n-1+e^{1-\gamma})+a(n)\leq n\log n+n\log_2 n-n+n\frac{\log_2 n-2.25}{\log n}.$$

Since, we want to find the maximum lower bound in the left hand side of (2.1), the best possible choice for $a(n)$, is:

$$a(n)=n\log_2 n-n(1+\gamma)+n\left(\frac{\log_2 n-2.25}{\log n}-\log\left(1+\frac{e^{1-\gamma}-1}{n}\right)\right).$$

So, we have:

$$nH_n+n\log_2 n-n(1+\gamma)+n\left(\frac{\log_2 n-2.25}{\log n}-\log\left(1+\frac{e^{1-\gamma}-1}{n}\right)\right)\leq p_n, \tag{2.5}$$

which holds for $n\geq 2$. Therefore, considering (2.4) and (2.5), for every $n\geq 27076$, we obtain:

$$|p_n-(nH_n+n\log_2 n-n(1+\gamma))|\leq n\left(\frac{\log_2 n-1.8}{\log n}-\log\left(1+\frac{0.5}{n}\right)\right). \tag{2.6}$$

3. An Equivalent for the PNT

Considering (2.6), we obtain:

$$p_n = nH_n + n\log_2 n - n(1+\gamma) + O\left(\frac{n\log_2 n}{\log n}\right),$$

which is a very strong form of an equivalent for the PNT. In fact we observe that the PNT holds if and only if $p_n \sim nH_n$, when n tends to infinity. To see this, first suppose that the PNT holds true. So, when n tends to infinity, we have:

$$p_n = n\log n + o(n\log n).$$

Considering this with $H_n \sim \log n$, we obtain:

$$\begin{aligned} p_n &= n(H_n + O(1)) + o(n\log n) \\ &= nH_n + O(n) + o(n\log n) \\ &= nH_n + o(n\log n) \\ &= nH_n + o(n(H_n + O(1))) \\ &= nH_n + o(nH_n) + o(O(n)) \\ &= nH_n + o(nH_n). \end{aligned}$$

Inversely, suppose $p_n \sim nH_n$, then:

$$p_n = nH_n + o(nH_n),$$

which considering this with $H_n \sim \log n$, we obtain:

$$\begin{aligned} p_n &= n(\log n + O(1)) + o(n(\log n + O(1))) \\ &= n\log n + O(n) + o(n\log n + O(n)) \\ &= n\log n + O(n) + o(n\log n) \\ &= n\log n + o(n\log n), \end{aligned}$$

and this is PNT.

References

[1] M. Abramowitz and I.A. Stegun, *HANDBOOK OF MATHEMATICAL FUNCTIONS: with Formulas, Graphs, and Mthematical Tables*, Dover Publications, 1972.

[2] N. Batir, "Some new inequalities for Gamma and polygamma functions", *RGMIA Research Report Collection*, **7**(3), Article 1, 2004.

[3] H. Davenport, Multiplicative Number Theory (Second Edition), Springer-Verlag, 1980.

[4] P. Dusart, Inégalités explicites pour $\psi(X)$, $\theta(X)$, $\pi(X)$ et les nombres premiers, *C. R. Math. Acad. Sci. Soc. R. Can.* **21** (1999), no. 2, 53–59.

Institute for Advanced, Studies in Basic Sciences, P.O. Box 45195-1159, Zanjan, Iran.
E-mail address: alikhani@iasbs.ac.ir, mmhassany@srttu.edu