



VICTORIA UNIVERSITY
MELBOURNE AUSTRALIA

A Function Involving Gamma Function and Having Logarithmically Absolute Convexity

This is the Published version of the following publication

Qi, Feng, Guo, Bai-Ni and Guo, Senlin (2007) A Function Involving Gamma Function and Having Logarithmically Absolute Convexity. Research report collection, 10 (1).

The publisher's official version can be found at

Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/18270/>

A FUNCTION INVOLVING GAMMA FUNCTION AND HAVING LOGARITHMICALLY ABSOLUTE CONVEXITY

FENG QI, BAI-NI GUO, AND SENLIN GUO

ABSTRACT. In the paper, the logarithmically complete monotonicity, logarithmically absolute monotonicity and logarithmically absolute convexity of the function $\frac{[\Gamma(1+tx)]^s}{[\Gamma(1+sx)]^t}$ for $x, s, t \in \mathbb{R}$ such that $1+sx > 0$ and $1+tx > 0$ with $s \neq t$ are verified, some known results are generalized.

1. INTRODUCTION

By using a geometrical method in [1], the following double inequality was proved:

$$\frac{1}{n!} \leq \frac{[\Gamma(1+x)]^n}{\Gamma(1+nx)} \leq 1 \quad (1)$$

for $x \in [0, 1]$ and $n \in \mathbb{N}$, where $\Gamma(x)$ stands for the classical Euler's gamma function defined for $x > 0$ by $\Gamma(x) = \int_0^\infty e^{-t} t^{x-1} dt$.

By analytical arguments in [29], it was presented that the function

$$f(x, y) = \frac{[\Gamma(1+x)]^y}{\Gamma(1+xy)} \quad (2)$$

for all $y \geq 1$ is decreasing in $x \geq 0$. From this, it is deduced that

$$\frac{1}{\Gamma(1+y)} \leq \frac{[\Gamma(1+x)]^y}{\Gamma(1+xy)} \leq 1 \quad (3)$$

for all $y \geq 1$ and $x \in [0, 1]$, which is a generalization of inequality (1).

Recall [12, 30, 31] that a function f is called completely monotonic on an interval I if f has derivatives of all orders on I and $(-1)^k f^{(k)}(x) \geq 0$ for all $k \geq 0$ on I . The set of completely monotonic functions on I is denoted by $\mathcal{C}[I]$. Recall also [20, 21, 22] that a positive function f is said to be logarithmically completely monotonic on an interval I if f has derivatives of all orders on I and its logarithm $\ln f$ satisfies $(-1)^k [\ln f(x)]^{(k)} \geq 0$ for all $k \in \mathbb{N}$ on I . The set of logarithmically completely monotonic functions on I is denoted by $\mathcal{L}[I]$. Recall [3] that if $f^{(k)}(x)$ for some nonnegative integer k is completely monotonic on an interval I , but $f^{(k-1)}(x)$

Date: This paper was completed on 4 February 2007.

2000 Mathematics Subject Classification. Primary 33B15; Secondary 26A48.

Key words and phrases. logarithmically absolutely convex function, logarithmically completely monotonic function, logarithmically absolutely monotonic function, gamma function, generalization.

This paper was typeset using $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{L}\mathcal{A}\mathcal{T}\mathcal{E}\mathcal{X}$.

is not completely monotonic on I , then $f(x)$ is called a completely monotonic function of k -th order on an interval I . Among other things, it was proved implicitly or explicitly in [4, 14, 18, 20, 22, 30] that a logarithmically completely monotonic function is always completely monotonic, but not conversely. Motivated or stimulated by the papers [20, 22], among other things, it is further revealed in [4] that $\mathcal{S} \setminus \{0\} \subset \mathcal{L}[(0, \infty)] \subset \mathcal{C}[(0, \infty)]$, where \mathcal{S} denotes the set of Stieltjes transforms. In [4, Theorem 1.1] and [9, 21] it is pointed out that the logarithmically completely monotonic functions on $(0, \infty)$ can be characterized as the infinitely divisible completely monotonic functions studied by Horn in [10, Theorem 4.4]. For more information on the logarithmically completely monotonic functions, please refer to [3, 4, 6, 7, 8, 9, 15, 18, 19, 20, 21, 22, 23, 25, 28, 30] and the references therein.

Definition 1. Let f be a positive function which has derivatives of all orders on an interval I . If $[\ln f(x)]^{(k)}$ for some nonnegative integer k is completely monotonic on I , but $[\ln f(x)]^{(k-1)}$ is not completely monotonic on I , then f is said to be a logarithmically completely monotonic function of k -th order on I .

In [25, 28], the following logarithmically complete monotonicities, as generalizations of the decreasingly monotonic property in [29], are presented:

- (1) For given $y > 1$, the function $f(x, y)$ defined by (2) is decreasing and logarithmically concave with respect to $x \in (0, \infty)$, and $\frac{1}{f(x, y)}$ is a logarithmically completely monotonic function of second order in $x \in (0, \infty)$.
- (2) For given $0 < y < 1$, the function $f(x, y)$ defined by (2) is increasing and logarithmically convex with respect to $x \in (0, \infty)$, and $f(x, y)$ is a logarithmically completely monotonic function of second order in $x \in (0, \infty)$.
- (3) For given $x \in (0, \infty)$, the function $f(x, y)$ defined by (2) is logarithmically concave with respect to $y \in (0, \infty)$, and $\frac{1}{f(x, y)}$ is a logarithmically completely monotonic function of first order in $y \in (0, \infty)$.
- (4) For given $x \in (0, \infty)$, let

$$F_x(y) = \frac{\Gamma(1+y)[\Gamma(1+x)]^y}{\Gamma(1+xy)} \quad (4)$$

in $(0, \infty)$. If $0 < x < 1$, then $F_x(y)$ is a logarithmically completely monotonic function of second order in $(0, \infty)$; if $x > 1$, then $\frac{1}{F_x(y)}$ is a logarithmically completely monotonic function of second order in $(0, \infty)$.

In [13], It was shown that if f is a differentiable and logarithmically convex function in $[0, \infty)$, then the function $\frac{[f(x)]^a}{f(ax)}$ for $a \geq 1$ (or $0 < a \leq 1$ respectively) is decreasing (or increasing respectively) in $[0, \infty)$. As one of applications to inequalities involving gamma function, Riemann's zeta function and the complete elliptic integrals of the first kind, inequalities (1) and (3) were deduced.

In [11], an inequality involving a positive linear operator acting on the composition of two continuous functions is presented and, as applications of this inequality,

some new inequalities involving the beta, gamma and Riemann's zeta functions and a large family of functions which are Mellin transforms are produced. In particular, for $\beta > \delta > 0$, $\alpha\beta > -1$ and $\alpha\delta > -1$, if either $\alpha < 0$ or $\alpha > 1$, then inequality

$$\frac{[\Gamma(1+\delta)]^\alpha}{\Gamma(1+\alpha\delta)} > \frac{[\Gamma(1+\beta)]^\alpha}{\Gamma(1+\alpha\beta)} \quad (5)$$

holds true; if $0 < \alpha \leq 1$, inequality (5) is reversed. It is not difficult to see that the left hand sides in inequalities (1) and (3) are special cases of inequality (5).

In [5, Theorem 2.1], it was proved that the function

$$G_{s,t}(x) = \frac{[\Gamma(1+tx)]^s}{[\Gamma(1+sx)]^t} \quad (6)$$

is decreasing (or increasing respectively) in $x \in [0, \infty)$ if either $s \geq t > 0$ or $0 > s \geq t$ (or both $s > 0$ and $t < 0$ respectively) such that $1+sx > 0$ and $1+tx > 0$. This result generalized and extended the corresponding conclusions in [1, 11, 29].

Recall [12, 15, 16, 27, 30, 31] that a function f is said to be absolutely monotonic on an interval I if it has derivatives of all orders and $f^{(k-1)}(t) \geq 0$ for $t \in I$ and $k \in \mathbb{N}$. Recall also [14, 26] that a positive function f is said to be logarithmically absolutely monotonic on an interval I if it has derivatives of all orders and $[\ln f(t)]^{(k)} \geq 0$ for $t \in I$ and $k \in \mathbb{N}$. In [14, 26] it was proved that a logarithmically absolutely monotonic function on an interval I is also absolutely monotonic on I , but not conversely.

Definition 2. Let f be a positive function which has derivatives of all orders on an interval I . If $[\ln f(x)]^{(k)}$ for some nonnegative integer k is absolutely monotonic on I , but $[\ln f(x)]^{(k-1)}$ is not absolutely monotonic on I , then f is said to be a logarithmically absolutely monotonic function of k -th order on I .

Recall [12, p. 375, Definition 3] and [16, 17, 30, 31] that a function f which has derivatives of all orders on an interval I is said to be absolutely convex on I if $f^{(2k)}(x) \geq 0$ on I for any nonnegative integer k .

Definition 3. A positive function f which has derivatives of all orders on an interval I is said to be logarithmically absolutely convex on I if $[\ln f(x)]^{(2k)} \geq 0$ on I for $k \in \mathbb{N}$.

The main aim of this paper is to generalize and extend some results obtained in [1, 5, 11, 13, 25, 28, 29].

Our main results can be stated as the following theorem.

Theorem 1. *The function $G_{s,t}(x)$ defined by (6) for $x, s, t \in \mathbb{R}$ such that $1+sx > 0$ and $1+tx > 0$ with $s \neq t$ has the following properties:*

- (1) $G_{s,t}(x) = \frac{1}{G_{t,s}(x)}$;
- (2) For $t > s > 0$ and $x \in (0, \infty)$, $G_{s,t}(x)$ is an increasing function and a logarithmically completely monotonic function of second order in x ;

- (3) For $t > s > 0$ and $x \in (-\frac{1}{t}, 0)$, $G_{s,t}(x)$ is a logarithmically completely monotonic function in x ;
- (4) For $s < t < 0$ and $x \in (-\infty, 0)$, $G_{s,t}(x)$ is a decreasing function and a logarithmically absolutely monotonic function of second order in x ;
- (5) For $s < t < 0$ and $x \in (0, -\frac{1}{s})$, $G_{s,t}(x)$ is a logarithmically completely monotonic function in x ;
- (6) For $s < 0 < t$ and $x \in (-\frac{1}{t}, 0)$, $G_{t,s}(x)$ is an increasing function and a logarithmically absolutely convex function in x ;
- (7) For $s < 0 < t$ and $x \in (0, -\frac{1}{s})$, $G_{t,s}(x)$ is a decreasing function and a logarithmically absolutely convex function in x .

2. PROOF OF THEOREM 1

It is clear that $G_{s,t}(x) = \frac{1}{G_{t,s}(x)}$. Therefore, it is sufficient to show Theorem 1 only for $s < t$.

Taking logarithm of $G_{s,t}(x)$ yields

$$\ln G_{s,t}(x) = s \ln \Gamma(1 + tx) - t \ln \Gamma(1 + sx) \quad (7)$$

and

$$\begin{aligned} \frac{\partial^k [\ln G_{s,t}(x)]}{\partial x^k} &= st^k \psi^{(k-1)}(1 + tx) - ts^k \psi^{(k-1)}(1 + sx) \\ &= \frac{st}{x^{k-1}} [(tx)^{k-1} \psi^{(k-1)}(1 + tx) - (sx)^{k-1} \psi^{(k-1)}(1 + sx)] \end{aligned} \quad (8)$$

for $k \in \mathbb{N}$.

In [2, Lemma 2.2], it was obtained that the function $x^\alpha |\psi^{(i)}(1+x)|$ is strictly increasing in $(0, \infty)$ if and only if $\alpha \geq i$, where $i \in \mathbb{N}$ and $\alpha \in \mathbb{R}$. This was generalized in [24, Theorem 1] as follows: For $\beta \geq \frac{1}{2}$, the function $x^\alpha |\psi^{(i)}(x+\beta)|$ is strictly increasing in $[0, \infty)$ if and only if $\alpha \geq i \in \mathbb{N}$. In particular, the functions $x^{2i} \psi^{(2i)}(1+x)$ are decreasing and the functions $x^{2i-1} \psi^{(2i-1)}(1+x)$ are increasing in $[0, \infty)$ for $i \in \mathbb{N}$. From this, it is readily obtained for $i \in \mathbb{N}$, $t > s > 0$ and $x > 0$ that

$$\frac{\partial^{2i} [\ln G_{s,t}(x)]}{\partial x^{2i}} > 0 \quad \text{and} \quad \frac{\partial^{2i+1} [\ln G_{s,t}(x)]}{\partial x^{2i+1}} < 0.$$

Since

$$\frac{\partial [\ln G_{s,t}(x)]}{\partial x} = st[\psi(1+tx) - \psi(1+sx)] \quad (9)$$

and the psi function ψ is increasing in $(0, \infty)$, it is easy to see

$$\frac{\partial [\ln G_{s,t}(x)]}{\partial x} > 0.$$

Consequently, for $t > s > 0$ and $x > 0$, $G_{s,t}(x)$ is an increasing function in $x \in (0, \infty)$ and a logarithmically completely monotonic function of second order in $x \in (0, \infty)$.

If $t > s > 0$ and $0 > x > -\frac{1}{t}$, then

$$(-1)^{k-1} \frac{\partial^k [\ln G_{s,t}(x)]}{\partial x^k} = \frac{st}{(-x)^{k-1}} [(tx)^{k-1} \psi^{(k-1)}(1+tx) - (sx)^{k-1} \psi^{(k-1)}(1+sx)]$$

for $k \in \mathbb{N}$. Let $h_k(u) = u^k \psi^{(k)}(1+u)$ for $u \in (-1, 0)$ and $k \in \mathbb{N}$. Straightforward computation gives

$$\begin{aligned} h'_k(u) &= u^{k-1} [k\psi^{(k)}(1+u) + u\psi^{(k+1)}(1+u)] \\ &= (-u)^{k-1} \{k[(-1)^{k+1}\psi^{(k)}(1+u)] + (-u)[(-1)^{k+2}\psi^{(k+1)}(1+u)]\} > 0. \end{aligned}$$

This means that the function $h_k(u)$ for $u \in (-1, 0)$ and $k \in \mathbb{N}$ is increasing. Then it is concluded that $(-1)^{k-1} \frac{\partial^k [\ln G_{s,t}(x)]}{\partial x^k} < 0$ for $k \in \mathbb{N}$. As a result, the function $G_{s,t}(x)$ is logarithmically completely monotonic in $x \in (-\frac{1}{t}, 0)$ for $t > s > 0$.

If $s < t < 0$ and $x < 0$, then

$$(-1)^k \frac{\partial^k [\ln G_{s,t}(x)]}{\partial x^k} = \frac{st}{(-x)^{k-1}} [(sx)^{k-1} \psi^{(k-1)}(1+sx) - (tx)^{k-1} \psi^{(k-1)}(1+tx)]$$

for $k \in \mathbb{N}$, accordingly

$$(-1)^{2i} \frac{\partial^{2i} [\ln G_{s,t}(x)]}{\partial x^{2i}} > 0 \quad \text{and} \quad (-1)^{2i+1} \frac{\partial^{2i+1} [\ln G_{s,t}(x)]}{\partial x^{2i+1}} < 0$$

for $i \in \mathbb{N}$. From (9), it follows that $\frac{\partial [\ln G_{s,t}(x)]}{\partial x} < 0$. Hence, $G_{s,t}(x)$ is a logarithmically absolutely monotonic function of second order in $x \in (-\infty, 0)$ for $s < t < 0$.

If $s < t < 0$ and $0 < x < -\frac{1}{s}$, then formula (8) is valid. In virtue of the increasingly monotonic property of the function $h_k(u)$ for $u \in (-1, 0)$ and $k \in \mathbb{N}$, it is deduced that $\frac{\partial^k [\ln G_{s,t}(x)]}{\partial x^k} > 0$ for $k \in \mathbb{N}$. This means that the function $G_{s,t}(x)$ for $s < t < 0$ is logarithmically completely monotonic in $x \in (0, -\frac{1}{s})$.

If $s < 0 < t$ and $-\frac{1}{t} < x < 0$, then

$$\begin{aligned} \frac{\partial^k [\ln G_{s,t}(x)]}{\partial x^k} &= \frac{-st}{(-x)^{k-1}} \{(-1)^{k-1} (-tx)^{k-1} [(-1)^k \psi^{(k-1)}(1+tx)] \\ &\quad - (sx)^{k-1} [(-1)^k \psi^{(k-1)}(1+sx)]\} \end{aligned}$$

for $k \in \mathbb{N}$. Accordingly, it is obtained readily that $\frac{\partial^{2i} [\ln G_{s,t}(x)]}{\partial x^{2i}} < 0$ for $i \in \mathbb{N}$. This implies that the function $\frac{1}{G_{s,t}(x)}$ is logarithmically absolutely convex in $x \in (-\frac{1}{t}, 0)$ for $s < 0 < t$. Formula (9) implies that the function $G_{s,t}(x)$ is increasing in $x \in (-\frac{1}{t}, 0)$ for $s < 0 < t$.

If $s < 0 < t$ and $0 < x < -\frac{1}{s}$, then

$$\begin{aligned} \frac{\partial^k [\ln G_{s,t}(x)]}{\partial x^k} &= \frac{-st}{x^{k-1}} (-1)^k \{(sx)^{k-1} [(-1)^k \psi^{(k-1)}(1+sx)] \\ &\quad - (tx)^{k-1} [(-1)^k \psi^{(k-1)}(1+tx)]\} \end{aligned}$$

for $k \in \mathbb{N}$. Hence, it is deduced apparently that $\frac{\partial^{2i} [\ln G_{s,t}(x)]}{\partial x^{2i}} < 0$ for $i \in \mathbb{N}$. This implies that the function $\frac{1}{G_{s,t}(x)}$ is logarithmically absolutely convex in $x \in (0, -\frac{1}{s})$

for $s < 0 < t$. Formula (9) implies that the function $G_{s,t}(x)$ is decreasing in $x \in (0, -\frac{1}{s})$ for $s < 0 < t$. The proof of Theorem 1 is finished.

REFERENCES

- [1] C. Alsina and M. S. Tomás, *A geometrical proof of a new inequality for the gamma function*, J. Inequal. Pure Appl. Math. **6** (2005), no. 2, Art. 48. Available online at <http://jipam.vu.edu.au/article.php?sid=517>.
- [2] H. Alzer, *Sharp inequalities for the digamma and polygamma functions*, Forum Math. **16** (2004), 1851–221.
- [3] R. D. Atanassov and U. V. Tsoukrovski, *Some properties of a class of logarithmically completely monotonic functions*, C. R. Acad. Bulgare Sci. **41** (1988), no. 2, 21–23.
- [4] C. Berg, *Integral representation of some functions related to the gamma function*, Mediterr. J. Math. **1** (2004), no. 4, 433–439.
- [5] L. Bougoffa, *Some inequalities involving the gamma function*, J. Inequal. Pure Appl. Math. **7** (2006), no. 5, Art. 179. Available online at <http://jipam.vu.edu.au/article.php?sid=796>.
- [6] Ch.-P. Chen and F. Qi, *Logarithmically completely monotonic functions relating to the gamma function*, J. Math. Anal. Appl. **321** (2006), no. 1, 405–411.
- [7] Ch.-P. Chen and F. Qi, *Logarithmically completely monotonic ratios of mean values and an application*, Glob. J. Math. Math. Sci. **1** (2005), no. 1, 71–76. RGMIA Res. Rep. Coll. **8** (2005), no. 1, Art. 18. Available online at <http://rgmia.vu.edu.au/v8n1.html>.
- [8] Ch.-P. Chen and F. Qi, *Logarithmically complete monotonicity properties for the gamma functions*, Austral. J. Math. Anal. Appl. **2** (2005), no. 2, Art. 8. Available online at <http://ajmaa.org/cgi-bin/paper.pl?string=v2n2/V2I2P8.tex>.
- [9] A. Z. Grinshpan and M. E. H. Ismail, *Completely monotonic functions involving the Gamma and q-gamma functions*, Proc. Amer. Math. Soc. **134** (2006), 1153–1160.
- [10] R. A. Horn, *On infinitely divisible matrices, kernels and functions*, Z. Wahrscheinlichkeitstheorie und Verw. Geb **8** (1967), 219–230.
- [11] A. McD. Mercer, *Some new inequalities for the gamma, beta and zeta functions*, J. Inequal. Pure Appl. Math. **7** (2006), no. 1, Art. 29. Available online at <http://jipam.vu.edu.au/article.php?sid=636>.
- [12] D. S. Mitrinović, J. E. Pečarić and A. M. Fink, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, Dordrecht/Boston/London, 1993.
- [13] E. Neuman, *Inequalities involving a logarithmically convex function and their applications to special functions*, J. Inequal. Pure Appl. Math. **7** (2006), no. 1, Art. 16. Available online at <http://jipam.vu.edu.au/article.php?sid=646>.
- [14] F. Qi, *A property of logarithmically absolutely monotonic functions and logarithmically complete monotonicities of $(1 + \alpha/x)^{x+\beta}$* , Integral Transforms Spec. Funct. **18** (2007), accepted.
- [15] F. Qi, *Certain logarithmically N-alternating monotonic functions involving gamma and q-gamma functions*, RGMIA Res. Rep. Coll. **8** (2005), no. 3, Art. 5. Available online at <http://rgmia.vu.edu.au/v8n3.html>.
- [16] F. Qi, *Generalized weighted mean values with two parameters*, R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci. **454** (1998), no. 1978, 2723–2732.
- [17] F. Qi, *Logarithmic convexity of extended mean values*, Proc. Amer. Math. Soc. **130** (2002), no. 6, 1787–1796. RGMIA Res. Rep. Coll. **2** (1999), no. 5, Art. 5, 643–652. Available online at <http://rgmia.vu.edu.au/v2n5.html>.
- [18] F. Qi and Ch.-P. Chen, *A complete monotonicity property of the gamma function*, J. Math. Anal. Appl. **296** (2004), no. 2, 603–607.

- [19] F. Qi and W.-S. Cheung, *Logarithmically completely monotonic functions concerning gamma and digamma functions*, Integral Transforms Spec. Funct. **18** (2007), in press.
- [20] F. Qi and B.-N. Guo, *Complete monotonicities of functions involving the gamma and digamma functions*, RGMIA Res. Rep. Coll. **7** (2004), no. 1, Art. 8, 63–72. Available online at <http://rgmia.vu.edu.au/v7n1.html>.
- [21] F. Qi, B.-N. Guo, and Ch.-P. Chen, *Some completely monotonic functions involving the gamma and polygamma functions*, J. Austral. Math. Soc. **80** (2006), no. 1, 81–88.
- [22] F. Qi, B.-N. Guo, and Ch.-P. Chen, *Some completely monotonic functions involving the gamma and polygamma functions*, RGMIA Res. Rep. Coll. **7** (2004), no. 1, Art. 5, 31–36. Available online at <http://rgmia.vu.edu.au/v7n1.html>.
- [23] F. Qi, B.-N. Guo, and Ch.-P. Chen, *The best bounds in Gautschi-Kershaw inequalities*, Math. Inequal. Appl. **9** (2006), no. 3, 427–436. RGMIA Res. Rep. Coll. **8** (2005), no. 2, Art. 17. Available online at <http://rgmia.vu.edu.au/v8n2.html>.
- [24] F. Qi, S. Guo and B.-N. Guo, *Note on a class of completely monotonic functions involving the polygamma functions*, RGMIA Res. Rep. Coll. **10** (2006), no. 1, Art. 5. Available online at <http://rgmia.vu.edu.au/v10n1.html>.
- [25] F. Qi and W. Li, *Two logarithmically completely monotonic functions connected with the gamma function*, RGMIA Res. Rep. Coll. **8** (2005), no. 3, Art. 13, 497–493. Available online at <http://rgmia.vu.edu.au/v8n3.html>.
- [26] F. Qi, W. Li and B.-N. Guo, *Generalizations of a theorem of I. Schur*, RGMIA Res. Rep. Coll. **9** (2006), no. 3, Art. 15. Available online at <http://rgmia.vu.edu.au/v9n3.html>. Bùdēngshì Yānjiū Tōngxùn (Communications in Studies on Inequalities) **13** (2006), no. 4, 355–364.
- [27] F. Qi and S.-L. Xu, *The function $(b^x - a^x)/x$: Inequalities and properties*, Proc. Amer. Math. Soc. **126** (1998), no. 11, 3355–3359.
- [28] F. Qi, Q. Yang and W. Li, *Two logarithmically completely monotonic functions connected with gamma function*, Integral Transforms Spec. Funct. **17** (2006), no. 7, 539–542.
- [29] J. Sándor, *A note on certain inequalities for the gamma function*, J. Inequal. Pure Appl. Math. **6** (2005), no. 3, Art. 61. Available online at <http://jipam.vu.edu.au/article.php?sid=534>.
- [30] H. van Haeringen, *Completely Monotonic and Related Functions*, Report 93-108, Faculty of Technical Mathematics and Informatics, Delft University of Technology, Delft, The Netherlands, 1993.
- [31] D. V. Widder, *The Laplace Transform*, Princeton University Press, Princeton, 1941.

(F. Qi) RESEARCH INSTITUTE OF MATHEMATICAL INEQUALITY THEORY, HENAN POLYTECHNIC UNIVERSITY, JIAOZUO CITY, HENAN PROVINCE, 454010, CHINA

E-mail address: qifeng@hpu.edu.cn, qifeng618@hotmail.com, qifeng618@msn.com, qifeng618@qq.com, qifeng618@gmail.com, fengqi618@member.ams.org

URL: <http://rgmia.vu.edu.au/qi.html>

(B.-N. Guo) SCHOOL OF MATHEMATICS AND INFORMATICS, HENAN POLYTECHNIC UNIVERSITY, JIAOZUO CITY, HENAN PROVINCE, 454010, CHINA

E-mail address: guobaini@hpu.edu.cn

(S. Guo) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MANITOBA, WINNIPEG, MB, CANADA R3T 2N2

E-mail address: sguo@hotmail.com