



VICTORIA UNIVERSITY
MELBOURNE AUSTRALIA

On Weighted Mean Matrices Whose l_p Norms are Determined on Decreasing Sequences

This is the Published version of the following publication

Gao, Peng (2008) On Weighted Mean Matrices Whose l_p Norms are Determined on Decreasing Sequences. Research report collection, 11 (4).

The publisher's official version can be found at

Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/18397/>

ON WEIGHTED MEAN MATRICES WHOSE l^p NORMS ARE DETERMINED ON DECREASING SEQUENCES

PENG GAO

ABSTRACT. We give a condition on weighted mean matrices so that their l^p norms are determined on decreasing sequences when the condition is satisfied. We apply our result to give a proof of a conjecture of Bennett and discuss some related results.

1. INTRODUCTION

Suppose throughout that $p \neq 0$, $\frac{1}{p} + \frac{1}{q} = 1$. For $p \geq 1$, let l^p be the Banach space of all complex sequences $\mathbf{a} = (a_n)_{n \geq 1}$ with norm

$$\|\mathbf{a}\|_p := \left(\sum_{n=1}^{\infty} |a_n|^p \right)^{1/p} < \infty.$$

The celebrated Hardy's inequality ([17, Theorem 326]) asserts that for $p > 1$,

$$(1.1) \quad \sum_{n=1}^{\infty} \left| \frac{1}{n} \sum_{k=1}^n a_k \right|^p \leq \left(\frac{p}{p-1} \right)^p \sum_{n=1}^{\infty} |a_n|^p.$$

Hardy's inequality can be regarded as a special case of the following inequality:

$$(1.2) \quad \left\| C \cdot \mathbf{a} \right\|_p^p = \sum_{n=1}^{\infty} \left| \sum_{k=1}^{\infty} c_{n,k} a_k \right|^p \leq U_p \sum_{n=1}^{\infty} |a_n|^p,$$

in which $C = (c_{n,k})$ and the parameter $p > 1$ are assumed fixed, and the estimate is to hold for all complex sequences $\mathbf{a} \in l^p$. The l^p operator norm of C is then defined as

$$\|C\|_{p,p} = \sup_{\|\mathbf{a}\|_p=1} \left\| C \cdot \mathbf{a} \right\|_p.$$

It follows that inequality (1.2) holds for any $\mathbf{a} \in l^p$ when $U_p^{1/p} \geq \|C\|_{p,p}$ and fails to hold for some $\mathbf{a} \in l^p$ when $U_p^{1/p} < \|C\|_{p,p}$. Hardy's inequality thus asserts that the Cesàro matrix operator C , given by $c_{n,k} = 1/n$, $k \leq n$ and 0 otherwise, is bounded on l^p and has norm $\leq p/(p-1)$. (The norm is in fact $p/(p-1)$.)

We say a matrix $A = (a_{n,k})$ is a lower triangular matrix if $a_{n,k} = 0$ for $n < k$ and a lower triangular matrix A is a summability matrix if $a_{n,k} \geq 0$ and $\sum_{k=1}^n a_{n,k} = 1$. We say a summability matrix A is a weighted mean matrix if its entries satisfy:

$$(1.3) \quad a_{n,k} = \lambda_k / \Lambda_n, \quad 1 \leq k \leq n; \quad \Lambda_n = \sum_{i=1}^n \lambda_i, \quad \lambda_i \geq 0, \quad \lambda_1 > 0.$$

We shall also say that a weighted mean matrix A is generated by $\{\lambda_n\}_{n=1}^{\infty}$ (resp. $\{\lambda_n\}_{n=1}^N$) when A is an infinite weighted mean matrix (resp. finite $N \times N$ weighted mean matrix) whose entries are given by (1.3).

Date: October 6, 2008.

2000 Mathematics Subject Classification. Primary 47A30.

Key words and phrases. Hardy's inequality, Schur's test, weighted mean matrices.

Hardy's inequality (1.1) motivates one to determine the l^p operator norm of an arbitrary summability or weighted mean matrix A . In the weighted mean matrix case, as the diagonal entries $\{\lambda_n/\Lambda_n\}$ uniquely determine one such a matrix, one certainly expects to obtain a bound for its norm using only the diagonal terms. In [16], the author proved the following result:

Theorem 1.1. *Let $1 < p < \infty$ be fixed. Let A be a weighted mean matrix generated by $\{\lambda_n\}_{n=1}^\infty$. If for any integer $n \geq 1$, there exists a positive constant $0 < L < p$ such that*

$$(1.4) \quad \frac{\Lambda_{n+1}}{\lambda_{n+1}} \leq \frac{\Lambda_n}{\lambda_n} \left(1 - \frac{L\lambda_n}{p\Lambda_n}\right)^{1-p} + \frac{L}{p},$$

then $\|A\|_{p,p} \leq p/(p-L)$.

It is easy to see that the above result implies the following well-known result of Cartlidge [9] (see also [2, p. 416, Theorem C]):

Theorem 1.2. *Let $1 < p < \infty$ be fixed. Let A be a weighted mean matrix generated by $\{\lambda_n\}_{n=1}^\infty$. If*

$$(1.5) \quad L = \sup_n \left(\frac{\Lambda_{n+1}}{\lambda_{n+1}} - \frac{\Lambda_n}{\lambda_n} \right) < p,$$

then $\|A\|_{p,p} \leq p/(p-L)$.

The above result of Cartlidge is often very handy to apply for determining l^p norms of certain weighted mean matrices, when combined with a result of Cass and Kratz [10], which says that for a weighted mean matrix A generated by $\{\lambda_n\}_{n=1}^\infty$, with the λ_n 's generated by a positive logarithmico-exponential function (for details, see [14]) and satisfying $\lim_{n \rightarrow \infty} \Lambda_n/(n\lambda_n) = L < p$, then $\|A\|_{p,p} \geq p/(p-L)$. As an example, we note the following two inequalities were claimed to hold (with no proofs supplied) by Bennett ([4, p. 40-41]; see also [5, p. 407]):

$$(1.6) \quad \sum_{n=1}^{\infty} \left| \frac{1}{n^\alpha} \sum_{i=1}^n (i^\alpha - (i-1)^\alpha) a_i \right|^p \leq \left(\frac{\alpha p}{\alpha p - 1} \right)^p \sum_{n=1}^{\infty} |a_n|^p,$$

$$(1.7) \quad \sum_{n=1}^{\infty} \left| \frac{1}{\sum_{i=1}^n i^{\alpha-1}} \sum_{i=1}^n i^{\alpha-1} a_i \right|^p \leq \left(\frac{\alpha p}{\alpha p - 1} \right)^p \sum_{n=1}^{\infty} |a_n|^p,$$

whenever $p > 1, \alpha p > 1$. We note here the constant $(\alpha p/(\alpha p - 1))^p$ is best possible by the result of Cass and Kratz (or see [6]).

Straightforward applications of Theorem 1.2 allow the author [14] to prove inequalities (1.6) for $p > 1, \alpha \geq 1$ and (1.7) for $p > 1, \alpha \geq 2$ or $0 < \alpha \leq 1, \alpha p > 1$. The same result was obtained for (1.7) by Bennett himself [6] independently and his proof also relies on Cartlidge's result. Using a different approach, Bennett was able to prove (1.6) for the full range of α (see [6, Theorem 1] with $\beta = 1$ there). Using the result of Theorem 1.1, the author [16] has shown that inequality (1.7) holds for $p \geq 2, 1 < \alpha < 2$ (in fact, as pointed out in [16], for fixed $1 < p < 2$, one can also prove (1.7) for some cases of $1 < \alpha < 2$).

We note here that by a change of variables $a_k \rightarrow a_k^{1/p}$ in (1.1) and on letting $p \rightarrow +\infty$, one obtains the following well-known Carleman's inequality [8], which asserts that for convergent infinite series $\sum a_n$ with non-negative terms, one has

$$\sum_{n=1}^{\infty} \left(\prod_{k=1}^n a_k \right)^{\frac{1}{n}} \leq e \sum_{n=1}^{\infty} a_n,$$

with the constant e being best possible.

It is then natural to study the following weighted version of Carleman's inequality:

$$(1.8) \quad \sum_{n=1}^N \left(\prod_{k=1}^n a_k^{\lambda_k/\Lambda_n} \right) \leq E_N \sum_{n=1}^N a_n,$$

where the notations are as in (1.3) and $N \geq 1$ is an integer or $N = \infty$. The task here is to determine the best constant E_N so that inequality (1.8) holds for any (convergent when $N = \infty$) series $\sum a_n$ with non-negative terms. Note that (1.8) can be regarded as the $p \rightarrow +\infty$ case of the following inequality (once again by a change of variables):

$$(1.9) \quad \sum_{n=1}^N \left(\sum_{k=1}^n \frac{\lambda_k}{\Lambda_n} a_k \right)^p \leq U_{p,N} \sum_{n=1}^N a_n^p,$$

where $U_{p,N}$ is a positive constant, $a_n \geq 0$ and λ_n, Λ_n 's are given as in (1.3).

Note that Cartlidge's result (Theorem 1.2) implies that when (1.5) is satisfied, then for any $\mathbf{a} \in l^p$, inequality (1.9) holds for any N with $U_{p,N} = (p/(p-L))^p$. Similar to our discussions above, by a change of variables $a_k \rightarrow a_k^{1/p}$ in (1.9) and on letting $p \rightarrow +\infty$, one obtains inequality (1.8) with $E_N = e^L$ as long as (1.5) is satisfied with p replaced by $+\infty$ there.

In connection to (1.7), Bennett [6, p. 829] further conjectured that inequality (1.8) holds for $\lambda_k = k^\alpha$ for $\alpha > -1$ with $E_\infty = e^{1/(\alpha+1)}$. As the cases $-1 < \alpha \leq 0$ or $\alpha \geq 1$ follow directly from the known cases of inequalities (1.7) upon changes of variables $\alpha \rightarrow \alpha + 1, a_k \rightarrow a_k^{1/p}$ and on letting $p \rightarrow +\infty$, the only nontrivial cases are when $0 < \alpha < 1$. As these cases are the limits of the corresponding l^p cases and the author [16] has shown (1.7) hold for $p \geq 2, 1 < \alpha < 2$ using Theorem 1.1, it follows that Bennett's conjecture is true.

Motivated by the study of inequalities (1.6)-(1.7), we seek for extra inputs that may lead to a resolution of the remaining case of (1.7) for $1 < p < 2, 1 < \alpha < 2$. For this, we note the following natural question related to the l^p norms of any matrix asked by Bennett [5, Problem 7.23]: When is the norm of a matrix determined by its action on decreasing sequences? In other words, when do we have

$$(1.10) \quad \|C\|_{p,p} = \sup \left\{ \|C \cdot \mathbf{a}\|_p : \|\mathbf{a}\|_p = 1 \text{ and } \mathbf{a} \text{ decreasing} \right\}?$$

For weighted mean matrices, it is known that [2, p. 422] that sequences \mathbf{a} , with $a_n/\lambda_n^{1/(p-1)}$ decreasing in n , are sufficient to determine the norm. Note that this certainly implies (1.10) when the λ_n 's are decreasing. A slightly generalization of this later case is given in the following lemma:

Lemma 1.1. [11, Lemma 2.4] *Let $p > 1$ and $C = (c_{n,k})_{n,k \geq 1}$ be an arbitrary lower triangular matrix. If $c_{n,k} \geq c_{n,k+1} \geq 0$ for all $n \geq 1, 1 \leq k \leq n-1$, then (1.10) holds.*

We refer the reader to the articles [12] and [13] for more recent developments in this area. It is our goal in this paper to give a condition on weighted mean matrices in Section 2 so that (1.10) will hold. As an application, we will give another proof of the above mentioned Bennett's conjecture.

We note that Cartlidge's result (Theorem 1.2) only allows one to prove (1.6) with some restrictions on the α 's, as in [14], leaving alone the cases $1/p < \alpha \leq 1$. However, for these cases, Lemma 1.1 implies that (1.10) holds for the corresponding matrices. This extra information can be used to give a proof of these cases and in fact we shall prove a more general result in Section 3.

In [16], the author has shown that several approaches in the literature concerning the l^p norms of weighted mean matrices are equivalent. In Section 4, we will consider another approach to the l^p norms of weighted mean matrices, namely the Schur's test. We will show that Schur's test is equivalent to the other approaches mentioned in [16] and we shall point out how Bennett's proof of (1.6) can be rewritten using Schur's test. We shall also apply Schur's test to give extensions of (1.6) which in turn allows us to view both inequalities (1.6) and (1.7) as special cases of a family of inequalities.

2. ON THE VALIDITY OF (1.10) FOR WEIGHTED MEAN MATRICES

In this section, we want to first present a result regarding the validity of (1.10) for weighted mean matrices. Since one can often reduce the questions of finding the norms of infinite weighted mean matrices to that of finite ones, we consider only finite weighted mean matrices here. Thus instead of (1.2), we consider (1.9) instead and we have

Theorem 2.1. *Let $p > 1$ be fixed and let $N \geq 1$ be a fixed integer and A a weighted mean matrix generated by $\{\lambda_n\}_{n=1}^N$. Suppose that (1.9) is satisfied for some positive constant $U_{p,N}$. If for any $1 \leq k \leq N - 1$, the following condition*

$$(2.1) \quad \frac{1}{\Lambda_k} \geq U_{p,N} \left(\frac{1}{\lambda_k} - \frac{1}{\lambda_{k+1}} \right)$$

is satisfied, then (1.10) holds for $C = A$ in this case.

Proof. Since our matrix A is of finite dimension, it is easy to see that in this case we have

$$\mu_{p,N}^{1/p} := \|A\|_{p,p} = \max_{\|\mathbf{a}\|_p=1} \|A \cdot \mathbf{a}\|_p.$$

Thus without loss of generality, we may assume that the maximum is reached at some \mathbf{a} with $\|\mathbf{a}\|_p = 1$. It is shown in [16] that in this case we have $a_n > 0$ for all $1 \leq n \leq N$ and on setting

$$A_n = \sum_{k=1}^n \frac{\lambda_k a_k}{\Lambda_n},$$

we also have

$$(2.2) \quad \mu_{p,N} \left(\frac{a_k^{p-1}}{\lambda_k} - \frac{a_{k+1}^{p-1}}{\lambda_{k+1}} \right) = \frac{A_k^{p-1}}{\Lambda_k}, \quad 1 \leq k \leq N - 1; \quad \mu_{p,N} \frac{a_N^{p-1}}{\lambda_N} = \frac{A_N^{p-1}}{\Lambda_N}; \quad \sum_{n=1}^N a_n^p = 1.$$

We now show by induction on k that if (2.1) is satisfied, then the sequence \mathbf{a} satisfying (2.2) must be decreasing. First, it is easy to see that $a_1 \geq a_2$ using the relation $k = 1$ in (2.2) and noting that $A_1 = a_1$ and $0 < \mu_{p,N} \leq U_{p,N}$ by assumption. It now follows by induction that $A_k \geq a_k$ for $k \geq 1$ and that $a_k \geq a_{k+1}$ now follows from the k -th relation in (2.2) and this establishes our assertion. \square

We note here that one sees from (2.2) that that sequence \mathbf{a} with $a_n/\lambda_n^{1/(p-1)}$ decreasing in n , are sufficient to determine the norm, this is mentioned in Section 1.

Now to apply Theorem 2.1, one needs to find some constant $U_{p,N}$ so that (1.9) holds. This is not a problem in many cases, as one can apply Theorem 1.1 or Theorem 1.2. For example, if we use Theorem 1.2, then we can deduce the following result from Theorem 2.1:

Corollary 2.1. *Let $p > 1$ be fixed and let $N \geq 1$ be a fixed integer and A a weighted mean matrix generated by $\{\lambda_n\}_{n=1}^N$. Suppose that (1.5) is satisfied and for any $1 \leq k \leq N - 1$, we have*

$$(2.3) \quad \left(1 - \frac{L}{p} \right)^p \geq \Lambda_k \left(\frac{1}{\lambda_k} - \frac{1}{\lambda_{k+1}} \right),$$

then (1.10) holds for $C = A$ in this case.

We note that the left-hand side expression of (2.3) is an increasing function of p for fixed L . Thus if $L < 1$, then upon taking $p = 1$, we see that (1.10) holds for any $p > 1$ as long as

$$(2.4) \quad \inf_n \left(\frac{\Lambda_{n+1}}{\lambda_{n+1}} - \frac{\Lambda_n}{\lambda_n} \right) \geq L.$$

One should compare the above with (1.5). Interestingly enough, (2.4) tells us that if the condition (1.5) fails in the worst possible way (so that (2.4) holds), then Cartledge's result (Theorem 1.2)

does not help in determining the norm but then we can have an extra input by knowing that in this case (1.10) holds, provided that we know the norm is bounded by $p/(p-L)$. In particular, we point out here that if inequalities (1.7) were true for $p > 1, 1 < \alpha < 2$ (note that it is shown in [16] that this is the case when $p \geq 2$), then Theorem 2.1 implies that one may focus on decreasing sequences when trying to prove (1.7), since in this case (2.4) holds with $\lambda_k = k^{\alpha-1}$ and $L = 1/\alpha$ (see [6, Theorem 6]). Of course one is not able to apply (2.1) using the constant $U_{p,N} = (\alpha p/(\alpha p - 1))^p$ for the unknown cases of (1.7). However, for the case of p being large, one may hope to find a coarse bound $U_{p,N}$ so that (1.7) hold with the constant $(\alpha p/(\alpha p - 1))^p$ replaced by $U_{p,N}$ and (2.1) is also satisfied and hopefully the extra information (that one may focus on decreasing sequences) will allow one to give a proof of (1.7) for the cases $1 < \alpha < 2$ and p large. We shall not worry about finding such a coarse bound here but we will show later in this section that the $p \rightarrow +\infty$ case (corresponding to the conjecture of Bennett mentioned in Section 1) follows from this approach.

By looking at the case $k = 1$ of (2.2), we see that the case $k = 1$ of (2.1) with $U_{p,N}$ replaced by $\mu_{p,N}$ is a necessary condition for $a_2 \geq a_1$. When $A = (a_{i,j})$ is an infinite weighted mean matrix, then we denote $A_N = (a_{i,j})_{1 \leq i,j \leq N}$ and let $\mu_{p,N} = \|A_N\|_{p,p}^p$ and note that we have $\mu_{p,N-1} \leq \mu_{p,N}$ for $N \geq 2$ (one sets $a_N = 0$ in (1.9) to see this), thus the sequence $\{\mu_{p,N}\}_{N=1}^\infty$ is increasing and thus we have $\mu_{p,N} \rightarrow \|A\|_{p,p}^p$ as $N \rightarrow +\infty$, which allows us to deduce immediately the following

Corollary 2.2. *Let $p > 1$ be fixed and A a weighted mean matrix generated by $\{\lambda_n\}_{n=1}^N$. A necessary condition for (1.10) to hold for $C = A$ is*

$$\frac{1}{\lambda_1} \geq \|A\|_{p,p}^p \left(\frac{1}{\lambda_1} - \frac{1}{\lambda_2} \right).$$

If moreover, the sequence $\{\Lambda_n/\lambda_n\}_{n=1}^\infty$ is convex, then the above condition is also sufficient.

We note here by a result of Bennett [6, Theorem 2], we know that the sequence $\{\Lambda_n/\lambda_n\}_{n=1}^\infty$ is convex when $\lambda_n = n^\alpha$ for $\alpha \geq 1$ or $\alpha \leq 0$ and is concave for $0 \leq \alpha \leq 1$.

We now consider two analogues of Theorem 2.1 here. First we note that we have a similar result concerning inequality (1.8), namely,

Theorem 2.2. *Let $N \geq 1$ be a fixed integer and suppose that E_N is the best possible constant to make (1.8) hold. If for any $1 \leq k \leq N-1$, inequality (2.1) is satisfied with $U_{p,N}$ replaced by E'_N for some constant $E'_N \geq E_N$ there, then to prove (1.8), it suffices to establish it for decreasing sequences.*

Next, we note that one can also study inequality (1.9) when $p < 0$ and one often expects to get result analogue to the case $p > 0$. To be precise, we consider the following inequality for $a_n \geq 0$ and $p < 0$,

$$(2.5) \quad \sum_{n=1}^N \left(\sum_{k=1}^n \frac{\lambda_k}{\Lambda_n} a_k^{1/p} \right)^p \leq U_{p,N} \sum_{n=1}^N a_n.$$

Here we define the value of the left-hand side expression above to be 0 when one or more of the a_n 's is zero. This makes the left-hand side expression above a continuous function on the compact set $\{\sum_{n=1}^N a_n = 1 | a_n \geq 0\}$ and therefore we have $U_{p,N} < \infty$. From now on, for a weighted mean matrix A generated by $\{\lambda_n\}_{n=1}^N$ (N finite or infinite) and a fixed $p < 0$, we shall denote $\|A\|_{p,p}^p$ for the supreme of the left-hand side expression of (2.5), over the set $\{\sum_{n=1}^N a_n = 1 | a_n \geq 0\}$. We now have the following analogue of Cartlidge's result for $p < 0$, which can be easily established by following the proof for the case $p > 1$ given in [15] by noting that the case $n = 1$ of (1.5) implies $L \geq 0$.

Theorem 2.3. *Let $p < 0$ be fixed and A a weighted mean matrix generated by $\{\lambda_n\}_{n=1}^N$. Then*

$$\sum_{n=1}^{\infty} A_n^p \geq \frac{p}{p-L} \sum_{n=1}^{\infty} a_n A_n^{p-1},$$

where L is given as in (1.5). In particular, inequality (2.5) holds with $\|A\|_{p,p}^p \leq (p/(p-L))^p$.

Now, analogue to Theorem 2.1, we have

Theorem 2.4. *Let $p < 0$ be fixed and $N \geq 1$ a fixed integer and A a weighted mean matrix generated by $\{\lambda_n\}_{n=1}^N$ and suppose that (2.5) holds for some constant $U_{p,N}$. If for any $1 \leq k \leq N-1$, inequality (2.1) is satisfied with $U_{p,N}$, then $\|A\|_{p,p}^p$ is determined on an increasing sequence.*

Now, we want to see what can be said about the l^p norm of a given matrix, taking into the account that (1.10) holds for such a matrix. One strategy is to find a matrix whose l^p norm (or an upper bound of it) is known, say by Cartlidge's result. Then one can make a comparison of the two matrices, thanks to the following result:

Lemma 2.1. [5, Lemma 2.1] *Let \mathbf{u}, \mathbf{v} be n -tuples with non-negative entries with $n \geq 1$ and*

$$\sum_{i=1}^k u_i \leq \sum_{i=1}^k v_i, \quad 1 \leq k \leq n-1; \quad \sum_{i=1}^n u_i = \sum_{i=1}^n v_i.$$

then

$$\sum_{i=1}^n u_i a_i \leq \sum_{i=1}^n v_i a_i,$$

for any decreasing n -tuple \mathbf{a} and the above inequality reverses when \mathbf{a} is increasing.

We note that the above lemma is given in [5, Lemma 2.1] for a slightly general statement, but only for the case when \mathbf{a} is decreasing and the case of \mathbf{a} being increasing follows by applying the previous case to $-\mathbf{a}$.

The above lemma allows us to deduce the following result:

Theorem 2.5. *Let A, A' be two weighted mean matrix generated by $\{\lambda_n\}_{n=1}^N$ and $\{\lambda'_n\}_{n=1}^N$ respectively. Suppose that $\Lambda_n/\lambda_n \leq \Lambda'_n/\lambda'_n$ for all n . Then for fixed $p > 1$, if (1.10) holds for $C = A$, we have $\|A\|_{p,p} \leq \|A'\|_{p,p}$. Similarly, for fixed $p < 0$, if $\|A\|_{p,p}^p$ is determined on an increasing sequence, we have $\|A\|_{p,p}^p \leq \|A'\|_{p,p}^p$.*

Proof. Since the proofs are similar, we will only prove the $p > 1$ case here. In this case as (1.10) holds for $C = A$, it follows from Lemma 2.1 that $\|A\|_{p,p} \leq \|A'\|_{p,p}$ as long as one can show that for any $k \leq n$,

$$\frac{\Lambda_k}{\Lambda_n} \leq \frac{\Lambda'_k}{\Lambda'_n}.$$

By induction, it suffices to establish the above inequality for $k = n-1$ and one sees easily in this case the above inequality is equivalent to $\Lambda_n/\lambda_n \leq \Lambda'_n/\lambda'_n$ and this completes the proof. \square

We note here the above theorem can be regarded as in the spirit of Bennett's "right is tight principle" (see page 409 of [5]) concerning the l^p norms of summability matrices. According to the above theorem, we can interpret this principle for the weighted mean matrices as saying that for two given weighted mean matrices, the one with termwise larger diagonal entries has smaller norm, provided its norm is determined on decreasing sequences.

As a concrete example of an application of the above theorem, we consider (1.7) for the cases $p > 1, 1 < \alpha < 2$. As we mentioned earlier, if we assume (1.7) hold for those cases, then (1.10) holds for the corresponding matrix and in fact this is the case at least for $p \geq 2, 1 < \alpha < 2$ as (1.7) are known to hold for these cases. Now assume (1.10) does hold for the corresponding matrix for

the cases $p > 1, 1 < \alpha < 2$ of (1.7), then in order to apply Theorem 2.5 to establish (1.7), we need to find a weighted mean matrix A' (we may again focus on the finite matrices) whose l^p norm is bounded by $\alpha p/(\alpha p - 1)$. Now for the cases $1 < \alpha < 2$ of (1.7), we consider the following choice of the matrix A' generated by $\{\lambda'_n\}_{n=1}^N$, satisfying

$$\lambda'_1 = 1, \quad \frac{\Lambda'_n}{\lambda'_n} = \frac{n + \alpha/2}{\alpha}, \quad n \geq 2.$$

Note that this defines the λ'_n 's uniquely and $\lambda'_n > 0$ for all n . For a fixed $1 < \alpha < 2$, we now apply Theorem 1.1 to conclude $\|A'\|_{p,p} \leq \alpha p/(\alpha p - 1)$ for $p > 1/(\alpha - 1)^2$ by noting that it suffices to prove the case $n = 1$ of (1.4) with $L = 1/\alpha$ and this case follows when we bound $(1 - 1/(p\alpha))^{1-p}$ from below by $1 - (1 - p)/(p\alpha) + (1 - 1/p)/(2\alpha^2)$. It is also easy to check that for $n \geq 2$,

$$\frac{\sum_{k=1}^n k^{\alpha-1}}{n^{\alpha-1}} \leq \frac{n + \alpha/2}{\alpha}.$$

One can similarly discuss the case $p < 0, 1 < \alpha < 2$ using the following analogue of Theorem 1.1:

Theorem 2.6. *Let $p < 0$ be fixed. Let A be a weighted mean matrix generated by $\{\lambda_n\}_{n=1}^N$. If for any integer $n \geq 1$, there exists a positive constant $L > 0$ such that*

$$\frac{\Lambda_{n+1}}{\lambda_{n+1}} \leq \frac{\Lambda_n}{\lambda_n} \left(1 - \frac{L\lambda_n}{p\Lambda_n}\right)^{1-p} + \frac{L}{p},$$

then $\|A\|_{p,p}^p \leq (p/(p - L))^p$.

Apply the above theorem to A' defined above, we see that $\|A'\|_{p,p}^p \leq (\alpha p/(\alpha p - 1))^p$ and we then deduce immediately from Theorem 2.5 the following

Corollary 2.3. *Inequalities (1.7) hold for $p < 0, 1 < \alpha < 2$ for any increasing sequence \mathbf{a} .*

Now, Corollary 2.3 allows us to give another proof of the nontrivial cases $0 < \alpha < 1$ of Bennett's conjecture and in fact we shall prove a slightly general version by first establishing

Theorem 2.7. *Let $p < 0$ be fixed and N an integer and A a weighted mean matrix generated by $\{\lambda_n\}_{n=1}^N$. Suppose that the sequence $\{\Lambda_n/\lambda_n\}_{n=1}^\infty$ is concave and that $\lim_{n \rightarrow +\infty} \Lambda_n/(n\lambda_n) = L$. If we have*

$$(2.6) \quad e^{\lambda_1/\lambda_2}(1 - L) < 1,$$

then $\|A\|_{p,p}^p$ is determined on an increasing sequence.

Proof. As $\{\Lambda_n/\lambda_n\}_{n=1}^\infty$ is concave and that $\lim_{n \rightarrow +\infty} \Lambda_n/(n\lambda_n) = L$, a result of Bennett [6, Lemma 2] implies that $L \leq \Lambda_{n+1}/\lambda_{n+1} - \Lambda_n/\lambda_n \leq \Lambda_2/\lambda_2 - \Lambda_1/\lambda_1 = \lambda_1/\lambda_2$. It follows from Theorem 2.3 that $\|A\|_{p,p}^p \leq (p/(p - \lambda_1/\lambda_2))^p$ for $p < 0$. Thus inequality (2.5) holds with $U_{p,N} = (p/(p - \lambda_1/\lambda_2))^p$. As $\lim_{p \rightarrow -\infty} (p/(p - \lambda_1/\lambda_2))^p = e^{\lambda_1/\lambda_2}$ and $(p/(p - \lambda_1/\lambda_2))^p$ is a decreasing function of $p < 0$, we see that inequality (2.1) holds with $U_{p,N} = (p/(p - \lambda_1/\lambda_2))^p$ by (2.6). Now our assertion follows from Theorem 2.4. \square

We now apply the above theorem to $\lambda_n = n^\alpha$ for $0 < \alpha < 1$, in which case (2.6) is equivalent to

$$1 + \frac{1}{\alpha} > e^{1/2^\alpha}.$$

As $e^{1/2^\alpha} < e$ when $0 < \alpha < 1$, it follows that the above inequality holds for $\alpha < 1/(e - 1) \approx 0.58$. Thus we may assume that $1/2 \leq \alpha < 1$ and in this case $e^{1/2^\alpha} < e^{1/\sqrt{2}}$ and by repeating the above argument, we see that we may further assume that $0.8 \leq \alpha < 1$ but then the above inequality holds since $e^{1/2^{0.8}} < 2$. Therefore, combined with Corollary 2.3, we see that inequalities (1.7) hold for $p < 0, 1 < \alpha < 2$ and for the other positive α 's, we can apply Theorem 2.3 to conclude that inequalities (1.7) hold as well and we summarize our result in the following

Corollary 2.4. *Inequalities (1.7) hold for $p < 0, \alpha > 0$.*

We note here that the above corollary implies the nontrivial cases $0 < \alpha < 1$ of Bennett's conjecture, which one obtains by taking $p \rightarrow -\infty$ of the corresponding cases of (1.7).

3. A GENERALIZATION OF A RESULT OF BENNETT

As we mentioned in the introduction, the validity of (1.10) will allow us to deduce the cases $1/p < \alpha \leq 1$ of inequalities (1.6). In this section, we shall generalize a result of Bennett which in turn implies these cases. We shall assume all the infinite sums converge and we start by noting the following result of Bliss [7]:

Theorem 3.1. *Let $r > p > 1$ and let α be a real number satisfying $(\alpha + 1)p > 1$. Let $f(x)$ be a non-negative measurable function on $[0, +\infty)$ such that $f \in L^p(0, +\infty)$. Then the integral $\int_0^x f(t)t^\alpha dt$ is finite for every x and*

$$\int_0^\infty \left(\int_0^x f(t)t^\alpha dt \right)^r \frac{dx}{x^{(\alpha+1)r-s}} \leq K_{r,s,\alpha} \left(\int_0^\infty f(x)^p dx \right)^{r/p},$$

where

$$s = r/p - 1, \quad K_{r,s,\alpha} = \frac{1}{(r-s-1)(1+\alpha q)^{r-s}} \left(\frac{s\Gamma(r/s)}{\Gamma(1/s)\Gamma((r-1)/s)} \right)^s.$$

We note here Bliss only proved the case $\alpha = 0$ in [7] but the general case can be obtained by some changes of variables. Based on the above result, we now prove the following

Theorem 3.2. *Let $r > s > 1$ and $s/r < \alpha \leq 1$. Let $\mathbf{u}, \mathbf{v}, \mathbf{a}$ be sequences with positive entries. Let $V_n = \sum_{k=1}^n v_k$ for $n \geq 1$ and $V_0 = 0$. If for $m \geq 1$,*

$$\sum_{n=1}^m u_n V_n^{\alpha r} \leq V_m^s.$$

Then

$$\sum_{n=1}^\infty u_n \left(\sum_{k=1}^n (V_k^\alpha - V_{k-1}^\alpha) a_k \right)^r \leq s\alpha^r K_{r,s-1,\alpha-1} \left(\sum_{n=1}^\infty v_n a_n^{r/s} \right)^s.$$

Proof. The proof is almost identical to the proof of Theorem 2 in [3], taking account into Theorem 3.1, as long as one can show (see also the proof of Theorem 1 in [2]) that for $1 \leq i < j, a_i < a_j$,

$$\frac{(V_i^\alpha - V_{i-1}^\alpha)a_i + (V_j^\alpha - V_{j-1}^\alpha)a_j}{(V_i^\alpha - V_{i-1}^\alpha) + (V_j^\alpha - V_{j-1}^\alpha)} \leq \frac{v_i a_i + v_j a_j}{v_i + v_j}.$$

The above inequality follows from Lemma 2.1 (note that $a_j > a_i$ here) provided that

$$\frac{V_j^\alpha - V_{j-1}^\alpha}{V_j - V_{j-1}} \leq \frac{V_i^\alpha - V_{i-1}^\alpha}{V_i - V_{i-1}}.$$

The above inequality holds by the mean value theorem, since the right-hand side is no less than $\alpha V_i^{\alpha-1}$ and the left-hand side is no greater than $\alpha V_{j-1}^{\alpha-1}$ and this completes the proof. \square

We now take $u_n = (V_n^s - V_{n-1}^s)/V_n^{\alpha r}$ in the above theorem and make a change of variables $a_n \rightarrow a_n^{s/r}$ and let $r \rightarrow +\infty$ to deduce that

Corollary 3.1. *Let $s > 1$ and $0 < \alpha \leq 1$. Let \mathbf{v}, \mathbf{a} be sequences with positive entries. Let $V_n = \sum_{k=1}^n v_k$ for $n \geq 1$ and $V_0 = 0$. Then*

$$\sum_{n=1}^\infty (V_n^s - V_{n-1}^s) \left(\prod_{k=1}^n a_k^{V_k^\alpha - V_{k-1}^\alpha} \right)^{s/V_n^\alpha} \leq \frac{e^{-(\alpha-1)s/\alpha}}{\alpha^{1-s}} \frac{s}{s-1} \left(\frac{s-1}{\Gamma(1/(s-1))} \right)^{s-1} \left(\sum_{n=1}^\infty v_n a_n \right)^s.$$

Note that we will get back Carleman-type inequalities on letting $s \rightarrow 1^+$ in the above corollary. We can also take $v_n = 1$ and $u_n = (n^s - (n-1)^s)/n^{\alpha r}$ in Theorem 3.1 to deduce that

Corollary 3.2. *Let $r > s > 1$ and $s/r < \alpha \leq 1$. Let \mathbf{a} be sequences with positive entries. Then*

$$\sum_{n=1}^{\infty} (n^s - (n-1)^s) \left(\frac{1}{n^\alpha} \sum_{k=1}^n (k^\alpha - (k-1)^\alpha) a_k \right)^r \leq s\alpha^r K_{r,s-1,\alpha-1} \left(\sum_{n=1}^{\infty} v_n a_n^{r/s} \right)^s.$$

Note that we get back the cases $1/p < \alpha \leq 1$ of (1.6) on setting $r = p$ and letting $s \rightarrow 1^+$ in the above corollary.

4. SCHUR'S TEST AND SOME GENERALIZATIONS OF INEQUALITIES (1.6) AND (1.7)

In this section we first state a discrete version of Schur's test concerning the norms of linear operators:

Lemma 4.1. *Let $p > 1$ be fixed and let $A = (\alpha_{j,i})_{1 \leq i, j \leq N}$ be a matrix with non-negative entries. If there exist positive numbers U_1, U_2 and two positive sequences $\mathbf{c} = (c_i), 1 \leq i \leq N; \mathbf{d} = (d_i), 1 \leq i \leq N$, such that*

$$(4.1) \quad \sum_{i=1}^N \alpha_{j,i} c_i^{1/p} \leq U_1 d_j^{1/p}, \quad 1 \leq j \leq N;$$

$$(4.2) \quad \sum_{j=1}^N \alpha_{j,i} d_j^{1/q} \leq U_2 c_i^{1/q}, \quad 1 \leq i \leq N.$$

Then

$$\|A\|_{p,p} \leq U_1^{1/q} U_2^{1/p}.$$

We now point out that Schur's test is equivalent to the approaches mentioned in [16] in determining the operator norms of weighted mean matrices. It suffices to show that it is equivalent to the approach of Kaluza and Szegö. To see this, note that our goal in general is to find some (smallest possible) constant $U_{p,N}$ so that for a weighted mean matrix generated by $\{\lambda_n\}_{n=1}^N$ (we may assume $\lambda_n > 0$ for all n), inequality (1.9) holds for any integer $N \geq 1$ and any $\mathbf{a} \in l^p$. We now apply Lemma 4.1 with $\alpha_{j,i} = \lambda_i/\Lambda_j$ for $i \leq j$ and $\alpha_{j,i} = 0$ for $i > j$ with

$$c_i = \left(\frac{w_i}{\lambda_i} \right)^p, \quad d_j = \left(\frac{\sum_{k=1}^j w_k}{\Lambda_j} \right)^p,$$

where the auxiliary sequence $\{w_n\}_{n=1}^{\infty}$ is of positive terms and to be determined later. The choice of the c_i 's and d_j 's is to make inequality (4.1) satisfied with $U_1 = 1$ (it becomes an identity) and inequality (4.2) becomes

$$(4.3) \quad \sum_{j=i}^N \frac{\lambda_i}{\Lambda_j^p} \left(\sum_{k=1}^j w_k \right)^{p-1} \leq U_2 \left(\frac{w_i}{\lambda_i} \right)^{p-1}.$$

Suppose now one can find for each $p > 1$ a positive constant U_2 , a sequence \mathbf{w} of positive terms with w_n^{p-1}/λ_n^p decreasing to 0, such that for any integer $n \geq 1$,

$$(w_1 + \cdots + w_n)^{p-1} < U_2 \Lambda_n^p \left(\frac{w_n^{p-1}}{\lambda_n^p} - \frac{w_{n+1}^{p-1}}{\lambda_{n+1}^p} \right),$$

then inequality (4.3) will follow from this and this is exactly the starting point of Kaluza and Szegö's approach.

In what follows, we will give an account of Bennett's proof of (1.6) in the form of Schur's test. First we consider the case $\alpha > 1/p$ of (1.6) and we can replace the infinite sums by finite sums from 1 to N with $N \geq 1$ here and we note the following estimation ([6, (99)]):

$$(4.4) \quad \sum_{j=i}^N \frac{\int_{i-1}^j x^{\alpha-1/p} dx}{j^{\alpha+1/q}} \leq \frac{1}{\alpha - 1/p}.$$

We now apply Lemma 4.1 with $\alpha_{j,i} = \alpha \left(\int_{i-1}^j x^{\alpha-1/p} dx \right)^{1/p} \left(\int_{i-1}^j x^{\alpha-1/p-1} dx \right)^{1/q} / j^\alpha$ for $i \leq j$ and $a_{j,i} = 0$ otherwise and $c_i = \left(\int_{i-1}^i x^{\alpha-1/p-1} dx / \int_{i-1}^i x^{\alpha-1/p} dx \right)$, $d_j = 1/j$, $U_1 = U_2 = (\alpha p)/(\alpha p - 1)$ to see that in this case inequality (4.1) becomes an identity and inequality (4.2) becomes exactly (4.4). From this we deduce the following inequality for $p > 1$, $\alpha > 1/p$ and any $\mathbf{a} \in l^p$,

$$\sum_{n=1}^{\infty} \left| \frac{1}{n^\alpha} \sum_{i=1}^n \alpha \left(\int_{i-1}^i x^{\alpha-1/p} dx \right)^{1/p} \left(\int_{i-1}^i x^{\alpha-1-1/p} dx \right)^{1/q} a_i \right|^p \leq \left(\frac{\alpha p}{\alpha p - 1} \right)^p \sum_{n=1}^{\infty} |a_n|^p.$$

from which one deduces the corresponding cases of (1.6) easily.

We note here in Bennett's proof of (1.6) given above, a key ingredient is inequality (4.4). We point out here that when $1 \leq \alpha \leq 1 + 1/p$, a better estimation exists, namely,

$$(4.5) \quad \sum_{j=i}^N \frac{\alpha \left(i - \frac{1}{2} \right)^{\alpha-1+1/q}}{j^{\alpha+1/q}} \leq \frac{\alpha p}{\alpha p - 1}.$$

Inequality (4.5) can be easily deduced from the following inequality for all integers $i \geq 1$ and $1 \leq \alpha \leq 1 + 1/p$,

$$i^{-\alpha-1/q} \leq \frac{1}{\alpha - 1/p} \left(\left(i - \frac{1}{2} \right)^{1-\alpha-1/q} - \left(i + \frac{1}{2} \right)^{1-\alpha-1/q} \right) = \int_{i-1/2}^{i+1/2} x^{-\alpha-1/q} dx.$$

The above inequality follows from the well-known Hadamard's inequality (with $h(x) = x^{-\alpha-1/q}$, $a = i - 1/2$, $b = i + 1/2$ below), which asserts that for a continuous convex function $h(x)$ on $[a, b]$,

$$h\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b h(x) dx \leq \frac{h(a) + h(b)}{2}.$$

The above inequality also allows us to see easily that inequality (4.5) improves upon (4.4) for $1 \leq \alpha \leq 1 + 1/p$.

Now, inequality (4.5) allows us to establish the following

Theorem 4.1. *Let $p > 1$ be fixed, then the following inequality holds for $1 \leq \alpha \leq 1 + 1/p$ and any $\mathbf{a} \in l^p$,*

$$\sum_{n=1}^{\infty} \left| \frac{1}{n^\alpha} \sum_{i=1}^n \alpha \left(i - \frac{1}{2} \right)^{\frac{1}{p}(\alpha-\frac{1}{p})} \left(\int_{i-1}^i x^{\alpha-1-1/p} dx \right)^{1/q} a_i \right|^p \leq \left(\frac{\alpha p}{\alpha p - 1} \right)^p \sum_{n=1}^{\infty} |a_n|^p.$$

Proof. We can replace the infinite sums by finite sums from 1 to N with $N \geq 1$ here and we apply Lemma 4.1 here with $\alpha_{j,i} = \alpha \left(i - \frac{1}{2} \right)^{\frac{1}{p}(\alpha-\frac{1}{p})} \left(\int_{i-1}^i x^{\alpha-1-1/p} dx \right)^{1/q} / j^\alpha$ for $i \leq j$ and 0 otherwise and $c_i = \left(i - \frac{1}{2} \right)^{-(\alpha-\frac{1}{p})} \left(\int_{i-1}^i x^{\alpha-1-1/p} dx \right)$, $d_j = j^{-1}$ to see that estimations (4.1)-(4.2) hold by (4.5) with $U_1 = U_2 = \alpha p/(\alpha p - 1)$ and this completes the proof. \square

To deduce interesting corollaries from Theorem 4.1, we note the following lemma:

Lemma 4.2 ([1, Lemma 2.1]). *Let $a > 0, b > 0$ and r be real numbers with $a \neq b$, and let*

$$\begin{aligned} L_r(a, b) &= \left(\frac{a^r - b^r}{r(a - b)} \right)^{1/(r-1)} \quad (r \neq 0, 1), \\ L_0(a, b) &= \frac{a - b}{\log a - \log b}, \\ L_1(a, b) &= \frac{1}{e} \left(\frac{a^a}{b^b} \right)^{1/(a-b)}. \end{aligned}$$

The function $r \mapsto L_r(a, b)$ is strictly increasing on \mathbb{R} .

It readily follows from the above lemma that for $1 \leq \alpha \leq 1 + 1/p$, we have

$$\begin{aligned} i^\alpha - (i-1)^\alpha &= \alpha L_\alpha^{\alpha-1}(i, i-1) \leq \alpha L_2^{\alpha-1}(i, i-1) = \alpha \left(i - 1/2 \right)^{\alpha-1} \\ &\leq \alpha \left(i - 1/2 \right)^{\frac{1}{p}(\alpha - \frac{1}{p})} \left(\int_{i-1}^i x^{\alpha-1-1/p} dx \right)^{1/q} = \alpha L_2^{\frac{1}{p}(\alpha - \frac{1}{p})}(i, i-1) \cdot L_{\alpha - \frac{1}{p}}^{\frac{1}{q}(\alpha - 1 - \frac{1}{p})}(i, i-1). \end{aligned}$$

It follows from this that Theorem 4.1 not only implies the corresponding cases of (1.6) but also the following stronger result:

Corollary 4.1. *Let $p > 1$ be fixed, then the following inequality holds for $1 \leq \alpha \leq 1 + 1/p$ and any $\mathbf{a} \in l^p$,*

$$\sum_{n=1}^{\infty} \left| \frac{1}{n^\alpha} \sum_{i=1}^n \alpha \left(i - \frac{1}{2} \right)^{\alpha-1} a_i \right|^p \leq \left(\frac{\alpha p}{\alpha p - 1} \right)^p \sum_{n=1}^{\infty} |a_n|^p.$$

As an interesting consequence of Corollary 4.1, we note for the case $p = 2$ we have $2\alpha - 1 \leq 2$ for $\alpha \leq 3/2$ so that Corollary 4.1 implies the following inequality for $\mathbf{a} \in l^2$ and $1 \leq \alpha \leq 3/2$:

$$(4.6) \quad \sum_{n=1}^N \left| \sum_{i=1}^n \frac{\alpha L_{2\alpha-1}^{\alpha-1}(i, i-1)}{n^\alpha} a_i \right|^2 \leq \frac{\alpha^2}{(\alpha - 1/2)^2} \sum_{i=1}^N |a_i|^2.$$

We now apply the duality principle [18, Lemma 2] to deduce from (4.6) the following inequality for $\mathbf{a} \in l^2, a_i \geq 0$ and $1 \leq \alpha \leq 3/2$:

$$\sum_{i,j=1}^N \frac{\alpha^2 \min(i^{2\alpha-1}, j^{2\alpha-1})}{(2\alpha - 1) i^\alpha j^\alpha} a_i a_j = \sum_{n=1}^N \left(\sum_{i=n}^N \frac{\alpha L_{2\alpha-1}^{\alpha-1}(n, n-1)}{i^\alpha} a_i \right)^2 \leq \frac{\alpha^2}{(\alpha - 1/2)^2} \sum_{i=1}^N a_i^2.$$

We note here the case $\alpha = 1$ above gives back a result of Schur in [19], who showed that for $\mathbf{x}, \mathbf{y} \in l^2$,

$$\sum_{i,j=1}^{\infty} \frac{x_i y_j}{\max(i, j)} \leq 4 \|\mathbf{x}\|_2 \|\mathbf{y}\|_2.$$

By the duality principle, the above inequality is equivalent to Hardy's inequality (1.1) for the case $p = 2$, even though this was not mentioned in [19] (this is actually prior to Hardy's discovery of (1.1)).

Our discussions above allow us to regard the cases of $\alpha \geq 1$ of inequalities (1.6) and (1.7) as special cases of a family of inequalities. Namely, it is interesting to determine the best constant $U = U(\alpha, \beta, p)$ so that the following inequality holds for all $\mathbf{a} \in l^p$ ($p > 1, \beta \geq \alpha \geq 1$):

$$(4.7) \quad \sum_{n=1}^{\infty} \left| \frac{1}{\sum_{k=1}^n L_\beta^{\alpha-1}(k, k-1)} \sum_{i=1}^n L_\beta^{\alpha-1}(i, i-1) a_i \right|^p \leq U \sum_{n=1}^{\infty} |a_n|^p.$$

Note that the case of $\beta = \alpha$ above corresponds to inequality (1.6) and the case of $\beta \rightarrow +\infty$ above corresponds to inequality (1.7) by Lemma 4.2. In both cases, we expect $U = (\alpha p / (\alpha p - 1))^p$ (of

course this is known except for some cases of (1.7) when $1 < p < 2, 1 < \alpha < 2$. Thanks to Corollary 4.1 and Lemma 4.2, we also know that inequality (4.7) holds with $U = (\alpha p / (\alpha p - 1))^p$ for $p > 1, 1 \leq \alpha \leq 1 + 1/p, \alpha \leq \beta \leq 2$.

ACKNOWLEDGEMENT

The author is supported by a research fellowship from an Academic Research Fund Tier 1 grant at Nanyang Technological University for this work.

REFERENCES

- [1] H. Alzer, Sharp bounds for the ratio of q -gamma functions, *Math. Nachr.*, **222** (2001), 5–14.
- [2] G. Bennett, Some elementary inequalities, *Quart. J. Math. Oxford Ser. (2)* **38** (1987), 401–425.
- [3] G. Bennett, Some elementary inequalities. III, *Quart. J. Math. Oxford Ser. (2)* **42** (1991), 149–174.
- [4] G. Bennett, Factorizing the classical inequalities, *Mem. Amer. Math. Soc.*, **120** (1996), 1–130.
- [5] G. Bennett, Inequalities complimentary to Hardy, *Quart. J. Math. Oxford Ser. (2)*, **49** (1998), 395–432.
- [6] G. Bennett, Sums of powers and the meaning of l^p , *Houston J. Math.*, **32** (2006), 801–831.
- [7] G. A. Bliss, An integral inequality, *J. London. Math. Soc.* **5** (1930), 40–46.
- [8] T. Carleman, Sur les fonctions quasi-analytiques, in *Proc. 5th Scand. Math. Congress*, Helsingfors, Finland, 1923, 181–196.
- [9] J. M. Cartlidge, Weighted mean matrices as operators on l^p , Ph.D. thesis, Indiana University, 1978.
- [10] F. P. Cass and W. Kratz, Nörlund and weighted mean matrices as operators on l_p , *Rocky Mountain J. Math.* **20** (1990), 59–74.
- [11] C.-P. Chen, D.-C. Luor and Z.-Y. Ou, Extensions of Hardy inequality, *J. Math. Anal. Appl.*, **273** (2002), 160–171.
- [12] C.-P. Chen, H.-W. Huang and C.-Y. Shen, Matrices whose norms are determined by their actions on decreasing sequences, *Canad. J. Math.*, **60** (2008), 520–531.
- [13] C.-P. Chen, C.-Y. Shen and K.-Z. Wang, Characterization of the matrix whose norm is determined by its action on decreasing sequences: The exceptional cases, arXiv:0710.0038.
- [14] P. Gao, A note on Hardy-type inequalities, *Proc. Amer. Math. Soc.*, **133** (2005), 1977–1984.
- [15] P. Gao, On a result of Cartlidge, *J. Math. Anal. Appl.*, **332** (2007), 1477–1481.
- [16] P. Gao, On l^p norms of weighted mean matrices, arXiv:0707.1473.
- [17] G. H. Hardy, J. E. Littlewood and G. Pólya, *Inequalities*, Cambridge Univ. Press, 1952.
- [18] H. L. Montgomery, The analytic principle of the large sieve, *Bull. Amer. Math. Soc.* **84** (1978), 547–567.
- [19] I. Schur, Bemerkungen zur Theorie der beschränkten Bilinearformen mit unendlich vielen Veränderlichen, *J. Reine Angew. Math.*, **140** (1911), 1–28.

DIVISION OF MATHEMATICAL SCIENCES, SCHOOL OF PHYSICAL AND MATHEMATICAL SCIENCES, NANYANG TECHNOLOGICAL UNIVERSITY, 637371 SINGAPORE

E-mail address: penggao@ntu.edu.sg