



VICTORIA UNIVERSITY
MELBOURNE AUSTRALIA

Bounding the Čebyšev function for a differentiable function whose derivative is h or λ -convex in absolute value and applications

This is the Published version of the following publication

Dragomir, Sever S (2016) Bounding the Čebyšev function for a differentiable function whose derivative is h or λ -convex in absolute value and applications. *Le Matematiche*, 71 (1). 173 - 202. ISSN 2037-5298

The publisher's official version can be found at
<https://www.dmi.unict.it/ojs/index.php/lematematiche/article/view/1243>
Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/33236/>

BOUNDING THE ČEBYŠEV FUNCTIONAL FOR A DIFFERENTIABLE FUNCTION WHOSE DERIVATIVE IS h OR λ -CONVEX IN ABSOLUTE VALUE AND APPLICATIONS

SILVESTRU SEVER DRAGOMIR

Some bounds for the Čebyšev functional of a differentiable function whose derivative is h or λ -convex in absolute value and applications for functions of selfadjoint operators in Hilbert spaces via the spectral representation theorem are given.

1. Introduction

For two Lebesgue integrable functions $f, g : [a, b] \rightarrow \mathbb{C}$, in order to compare the integral mean of the product with the product of the integral means, we consider the Čebyšev functional defined by

$$C(f, g) := \frac{1}{b-a} \int_a^b f(t)g(t) dt - \frac{1}{b-a} \int_a^b f(t) dt \cdot \frac{1}{b-a} \int_a^b g(t) dt.$$

In 1934, G. Grüss [55] showed that

$$|C(f, g)| \leq \frac{1}{4} (M - m)(N - n), \quad (1)$$

Entrato in redazione: 12 maggio 2015

AMS 2010 Subject Classification: 26D15, 47A63, 47A99.

Keywords: Čebyšev Functional, Grüss' inequality, Čebyšev's inequality, Ostrowski's inequality, Lupaş's inequality, Selfadjoint operators, Spectral Representation.

provided m, M, n, N are real numbers with the property that

$$-\infty < m \leq f \leq M < \infty, \quad -\infty < n \leq g \leq N < \infty \quad \text{a.e. on } [a, b]. \quad (2)$$

The constant $\frac{1}{4}$ is best possible in (1) in the sense that it cannot be replaced by a smaller one.

Another lesser known inequality for $C(f, g)$ was derived in [14] under the assumption that f', g' exist and are continuous on $[a, b]$, and is given by

$$|C(f, g)| \leq \frac{1}{12} \|f'\|_\infty \|g'\|_\infty (b - a)^2, \quad (3)$$

where $\|f'\|_\infty := \sup_{t \in [a, b]} |f'(t)| < \infty$.

The constant $\frac{1}{12}$ cannot be improved in general in (3).

Čebyšev's inequality (3) also holds if $f, g : [a, b] \rightarrow \mathbb{R}$ are assumed to be absolutely continuous and $f', g' \in L_\infty [a, b]$.

In 1970, A.M. Ostrowski [69] proved, amongst others, the following result that is in a sense a combination of the Čebyšev and Grüss results:

$$|C(f, g)| \leq \frac{1}{8} (b - a) (M - m) \|g'\|_\infty, \quad (4)$$

provided f is Lebesgue integrable on $[a, b]$ and satisfying (2) while $g : [a, b] \rightarrow \mathbb{R}$ is absolutely continuous and $g' \in L_\infty [a, b]$. Here the constant $\frac{1}{8}$ is also sharp.

In 1973, A. Lupuş [61] (see also [66, p. 210]) obtained the following result as well:

$$|C(f, g)| \leq \frac{1}{\pi^2} \|f'\|_2 \|g'\|_2 (b - a), \quad (5)$$

provided f, g are absolutely continuous and $f', g' \in L_2 [a, b]$.

Here the constant $\frac{1}{\pi^2}$ is the best possible as well.

In [11], P. Cerone and S.S. Dragomir proved the following inequalities:

$$|C(f, g)| \leq \begin{cases} \inf_{\gamma \in \mathbb{R}} \|g - \gamma\|_\infty \cdot \frac{1}{b-a} \int_a^b \left| f(t) - \frac{1}{b-a} \int_a^b f(s) ds \right| dt \\ \inf_{\gamma \in \mathbb{R}} \|g - \gamma\|_q \cdot \frac{1}{b-a} \left(\int_a^b \left| f(t) - \frac{1}{b-a} \int_a^b f(s) ds \right|^p dt \right)^{\frac{1}{p}} \end{cases} \quad (6)$$

where $p > 1, 1/p + 1/q = 1$.

For $\gamma = 0$, we get from the first inequality in (6)

$$|C(f, g)| \leq \|g\|_\infty \cdot \frac{1}{b-a} \int_a^b \left| f(t) - \frac{1}{b-a} \int_a^b f(s) ds \right| dt \quad (7)$$

for which the constant 1 cannot be replaced by a smaller constant.

If $m \leq g \leq M$ for a.e. $x \in [a, b]$, then $\|g - \frac{m+M}{2}\|_\infty \leq \frac{1}{2}(M - m)$ and by the first inequality in (6) we can deduce the following result obtained by Cheng and Sun [15]

$$|C(f, g)| \leq \frac{1}{2}(M - m) \cdot \frac{1}{b - a} \int_a^b \left| f(t) - \frac{1}{b - a} \int_a^b f(s) ds \right| dt. \tag{8}$$

The constant $\frac{1}{2}$ is best in (8) as shown by Cerone and Dragomir in [12].

The following result holds [33].

Theorem 1.1. *Let $f : [a, b] \rightarrow \mathbb{C}$ be of bounded variation on $[a, b]$ and $g : [a, b] \rightarrow \mathbb{C}$ a Lebesgue integrable function on $[a, b]$. Then*

$$|C(f, g)| \leq \frac{1}{2} \bigvee_a^b(f) \cdot \frac{1}{b - a} \int_a^b \left| g(t) - \frac{1}{b - a} \int_a^b g(s) ds \right| dt \tag{9}$$

where $\bigvee_a^b(f)$ denotes the total variation of f on the interval $[a, b]$.

The constant $\frac{1}{2}$ is best possible in (9).

We denote the variance of the function $f : [a, b] \rightarrow \mathbb{C}$ by $D(f)$ and defined as

$$D(f) = [C(f, \bar{f})]^{1/2} = \left[\frac{1}{b - a} \int_a^b |f(t)|^2 dt - \left| \frac{1}{b - a} \int_a^b f(t) dt \right|^2 \right]^{1/2}, \tag{10}$$

where \bar{f} denotes the complex conjugate function of f .

We have [33]:

Corollary 1.2. *If the function $f : [a, b] \rightarrow \mathbb{C}$ is of bounded variation on $[a, b]$, then*

$$D(f) \leq \frac{1}{2} \bigvee_a^b(f). \tag{11}$$

The constant $\frac{1}{2}$ is best possible in (11).

Now we can state the following result when both functions are of bounded variation [33]:

Corollary 1.3. *If $f, g : [a, b] \rightarrow \mathbb{C}$ are of bounded variation on $[a, b]$, then*

$$|C(f, g)| \leq \frac{1}{4} \bigvee_a^b(f) \bigvee_a^b(g). \tag{12}$$

The constant $\frac{1}{4}$ is best possible in (12).

Remark 1.4. We can consider the following quantity associated with a complex valued function $f : [a, b] \rightarrow \mathbb{C}$,

$$E(f) := |C(f, f)|^{1/2} = \left| \frac{1}{b-a} \int_a^b f^2(t) dt - \left(\frac{1}{b-a} \int_a^b f(t) dt \right)^2 \right|^{1/2}.$$

Utilising the above results we can state that

$$\begin{aligned} E^2(f) &\leq \frac{1}{2} \bigvee_a^b(f) \cdot \frac{1}{b-a} \int_a^b \left| f(t) - \frac{1}{b-a} \int_a^b f(s) ds \right| dt \quad (13) \\ &\leq \frac{1}{2} \bigvee_a^b(f) D(f) \leq \frac{1}{4} \left[\bigvee_a^b(f) \right]^2. \end{aligned}$$

If we consider

$$\begin{aligned} G(f) &:= |C(f, |f|)|^{1/2} \\ &= \left| \frac{1}{b-a} \int_a^b f(t) |f(t)| dt - \frac{1}{b-a} \int_a^b f(t) dt \cdot \frac{1}{b-a} \int_a^b |f(t)| dt \right|^{1/2}, \end{aligned}$$

then we also have

$$\begin{aligned} G^2(f) &\leq \frac{1}{2} \bigvee_a^b(f) \cdot \frac{1}{b-a} \int_a^b \left| |f(t)| - \frac{1}{b-a} \int_a^b |f(s)| ds \right| dt \quad (14) \\ &\leq \frac{1}{2} \bigvee_a^b(f) D(|f|) \leq \frac{1}{4} \bigvee_a^b(f) \bigvee_a^b(|f|) \leq \frac{1}{4} \left[\bigvee_a^b(f) \right]^2 \end{aligned}$$

and

$$\begin{aligned} G^2(f) &\leq \frac{1}{2} \bigvee_a^b(|f|) \cdot \frac{1}{b-a} \int_a^b \left| f(t) - \frac{1}{b-a} \int_a^b f(s) ds \right| dt \quad (15) \\ &\leq \frac{1}{2} \bigvee_a^b(|f|) D(f) \leq \frac{1}{4} \bigvee_a^b(f) \bigvee_a^b(|f|) \leq \frac{1}{4} \left[\bigvee_a^b(f) \right]^2. \end{aligned}$$

Motivated by the results presented above, we establish in this paper some new bounds for the magnitude of $C(f, g)$ in the case when one of the complex valued function, say f , is differentiable and the derivative is h -convex or λ -convex in absolute value while the other is Lebesgue integrable on $[a, b]$. Applications for functions of selfadjoint operators in Hilbert spaces via the spectral representation theorem are also given.

Before we are able to state our new results, we need the following preliminary facts about h -convex and λ -convex functions.

2. h -Convex and λ -Convex Functions

2.1. h -Convex Functions

We recall here some concepts of convexity that are well known in the literature.

Let I be an interval in \mathbb{R} .

Definition 2.1 ([54]). We say that $f : I \rightarrow \mathbb{R}$ is a Godunova-Levin function or that f belongs to the class $Q(I)$ if f is non-negative and for all $x, y \in I$ and $t \in (0, 1)$ we have

$$f(tx + (1 - t)y) \leq \frac{1}{t}f(x) + \frac{1}{1-t}f(y). \tag{16}$$

Some further properties of this class of functions can be found in [43], [44], [46], [63], [73] and [74]. Among others, it has been noted that non-negative monotone and non-negative convex functions belong to this class of functions.

Definition 2.2 ([46]). We say that a function $f : I \rightarrow \mathbb{R}$ belongs to the class $P(I)$ if it is nonnegative and for all $x, y \in I$ and $t \in [0, 1]$ we have

$$f(tx + (1 - t)y) \leq f(x) + f(y). \tag{17}$$

Obviously $Q(I)$ contains $P(I)$ and for applications it is important to note that also $P(I)$ contain all nonnegative monotone, convex and *quasi convex functions*, i. e. nonnegative functions satisfying

$$f(tx + (1 - t)y) \leq \max \{f(x), f(y)\} \tag{18}$$

for all $x, y \in I$ and $t \in [0, 1]$.

For some results on P -functions see [46] and [71] while for quasi convex functions, the reader can consult [45].

Definition 2.3 ([7]). Let s be a real number, $s \in (0, 1]$. A function $f : [0, \infty) \rightarrow [0, \infty)$ is said to be s -convex (in the second sense) or Breckner s -convex if

$$f(tx + (1 - t)y) \leq t^s f(x) + (1 - t)^s f(y)$$

for all $x, y \in [0, \infty)$ and $t \in [0, 1]$.

For some properties of this class of functions see [1], [2], [7], [8], [41], [42], [57], [59] and [76].

In order to unify the above concepts for functions of real variable, Varošanec introduced the concept of h -convex functions as follows.

Assume that I and J are intervals in \mathbb{R} , $(0, 1) \subseteq J$ and functions h and f are real non-negative functions defined in J and I , respectively.

Definition 2.4 ([80]). Let $h : J \rightarrow [0, \infty)$ with h not identical to 0. We say that $f : I \rightarrow [0, \infty)$ is an h -convex function if for all $x, y \in I$ we have

$$f(tx + (1-t)y) \leq h(t)f(x) + h(1-t)f(y) \quad (19)$$

for all $t \in (0, 1)$.

For some results concerning this class of functions see [80], [6], [61], [77], [75] and [79].

This concept can be extended for functions defined on convex subsets of linear spaces in the same way as above replacing the interval I by the corresponding convex subset C of the linear space X .

We can introduce now another class of functions.

Definition 2.5. We say that the function $f : C \subseteq X \rightarrow [0, \infty)$ is of s -Godunova-Levin type, with $s \in [0, 1]$, if

$$f(tx + (1-t)y) \leq \frac{1}{t^s}f(x) + \frac{1}{(1-t)^s}f(y), \quad (20)$$

for all $t \in (0, 1)$ and $x, y \in C$.

We observe that for $s = 0$ we obtain the class of P -functions while for $s = 1$ we obtain the class of Godunova-Levin. If we denote by $Q_s(C)$ the class of s -Godunova-Levin functions defined on C , then we obviously have

$$P(C) = Q_0(C) \subseteq Q_{s_1}(C) \subseteq Q_{s_2}(C) \subseteq Q_1(C) = Q(C)$$

for $0 \leq s_1 \leq s_2 \leq 1$.

For different inequalities related to these classes of functions, see [1]-[4], [6], [9]-[52], [58]-[60] and [71]-[79].

A function $h : J \rightarrow \mathbb{R}$ is said to be *supermultiplicative* if

$$h(ts) \geq h(t)h(s) \text{ for any } t, s \in J. \quad (21)$$

If the inequality (21) is reversed, then h is said to be *submultiplicative*. If the equality holds in (21) then h is said to be a multiplicative function on J .

In [80] it has been noted that if $h : [0, \infty) \rightarrow [0, \infty)$ with $h(t) = (x+c)^{p-1}$, then for $c = 0$ the function h is multiplicative. If $c \geq 1$, then for $p \in (0, 1)$ the function h is supermultiplicative and for $p > 1$ the function is submultiplicative.

We observe that, if h, g are nonnegative and supermultiplicative, the same is their product. In particular, if h is supermultiplicative then its product with a power function $\ell_r(t) = t^r$ is also supermultiplicative.

The following generalization of the Hermite-Hadamard inequality for h -convex functions defined on convex subsets of linear spaces holds [36].

Theorem 2.6. Assume that the function $f : C \subseteq X \rightarrow [0, \infty)$ is an h -convex function with $h \in L[0, 1]$. Let $y, x \in C$ with $y \neq x$ and assume that the mapping $[0, 1] \ni t \mapsto f[(1 - t)x + ty]$ is Lebesgue integrable on $[0, 1]$. Then

$$\frac{1}{2h\left(\frac{1}{2}\right)}f\left(\frac{x+y}{2}\right) \leq \int_0^1 f[(1-t)x+ty] dt \leq [f(x) + f(y)] \int_0^1 h(t) dt. \quad (22)$$

Remark 2.7. If $f : I \rightarrow [0, \infty)$ is an h -convex function on an interval I of real numbers with $h \in L[0, 1]$ and $f \in L[a, b]$ with $a, b \in I, a < b$, then from (22) we get the Hermite-Hadamard type inequality obtained by Sarikaya et al. in [75]

$$\frac{1}{2h\left(\frac{1}{2}\right)}f\left(\frac{a+b}{2}\right) \leq \int_a^b f(u) du \leq [f(a) + f(b)] \int_0^1 h(t) dt. \quad (23)$$

If we write (22) for $h(t) = t$, then we get the classical Hermite-Hadamard inequality for convex functions.

If we write (22) for the case of P -type functions $f : C \rightarrow [0, \infty)$, i.e., $h(t) = 1, t \in [0, 1]$, then we get the inequality

$$\frac{1}{2}f\left(\frac{x+y}{2}\right) \leq \int_0^1 f[(1-t)x+ty] dt \leq f(x) + f(y), \quad (24)$$

that has been obtained for functions of real variable in [46].

If f is Breckner s -convex on C , for $s \in (0, 1)$, then by taking $h(t) = t^s$ in (22) we get

$$2^{s-1}f\left(\frac{x+y}{2}\right) \leq \int_0^1 f[(1-t)x+ty] dt \leq \frac{f(x) + f(y)}{s+1}, \quad (25)$$

that was obtained for functions of a real variable in [41].

If $f : C \rightarrow [0, \infty)$ is of s -Godunova-Levin type, with $s \in [0, 1)$, then

$$\frac{1}{2^{s+1}}f\left(\frac{x+y}{2}\right) \leq \int_0^1 f[(1-t)x+ty] dt \leq \frac{f(x) + f(y)}{1-s}. \quad (26)$$

We notice that for $s = 1$ the first inequality in (26) still holds, i.e.

$$\frac{1}{4}f\left(\frac{x+y}{2}\right) \leq \int_0^1 f[(1-t)x+ty] dt. \quad (27)$$

The case for functions of real variables was obtained for the first time in [46].

2.2. λ -Convex Functions

We start with the following definition (see [37]):

Definition 2.8. Let $\lambda : [0, \infty) \rightarrow [0, \infty)$ be a function with the property that $\lambda(t) > 0$ for all $t > 0$. A mapping $f : C \rightarrow \mathbb{R}$ defined on convex subset C of a linear space X is called λ -convex on C if

$$f\left(\frac{\alpha x + \beta y}{\alpha + \beta}\right) \leq \frac{\lambda(\alpha)f(x) + \lambda(\beta)f(y)}{\lambda(\alpha + \beta)} \quad (28)$$

for all $\alpha, \beta \geq 0$ with $\alpha + \beta > 0$ and $x, y \in C$.

We observe that if $f : C \rightarrow \mathbb{R}$ is λ -convex on C , then f is h -convex on C with $h(t) = \frac{\lambda(t)}{\lambda(1)}$, $t \in [0, 1]$.

If $f : C \rightarrow [0, \infty)$ is h -convex function with h supermultiplicative on $[0, \infty)$, then f is λ -convex with $\lambda = h$.

Indeed, if $\alpha, \beta \geq 0$ with $\alpha + \beta > 0$ and $x, y \in C$ then

$$\begin{aligned} f\left(\frac{\alpha x + \beta y}{\alpha + \beta}\right) &\leq h\left(\frac{\alpha}{\alpha + \beta}\right)f(x) + h\left(\frac{\beta}{\alpha + \beta}\right)f(y) \\ &\leq \frac{h(\alpha)f(x) + h(\beta)f(y)}{h(\alpha + \beta)}. \end{aligned}$$

The following proposition contain some properties of λ -convex functions [37].

Proposition 2.9. Let $f : C \rightarrow \mathbb{R}$ be a λ -convex function on C .

(i) If $\lambda(0) > 0$, then we have $f(x) \geq 0$ for all $x \in C$;

(ii) If there exists $x_0 \in C$ so that $f(x_0) > 0$, then

$$\lambda(\alpha + \beta) \leq \lambda(\alpha) + \lambda(\beta)$$

for all $\alpha, \beta > 0$, i.e. the mapping λ is subadditive on $(0, \infty)$.

(iii) If there exists $x_0, y_0 \in C$ with $f(x_0) > 0$ and $f(y_0) < 0$, then

$$\lambda(\alpha + \beta) = \lambda(\alpha) + \lambda(\beta)$$

for all $\alpha, \beta > 0$, i.e. the mapping λ is additive on $(0, \infty)$.

We have the following result providing many examples of subadditive functions $\lambda : [0, \infty) \rightarrow [0, \infty)$.

Theorem 2.10 ([37]). *Let $h(z) = \sum_{n=0}^{\infty} a_n z^n$ a power series with nonnegative coefficients $a_n \geq 0$ for all $n \in \mathbb{N}$ and convergent on the open disk $D(0, R)$ with $R > 0$ or $R = \infty$. If $r \in (0, R)$ then the function $\lambda_r : [0, \infty) \rightarrow [0, \infty)$ given by*

$$\lambda_r(t) := \ln \left[\frac{h(r)}{h(r \exp(-t))} \right] \tag{29}$$

is nonnegative, increasing and subadditive on $[0, \infty)$.

Remark 2.11. Now, if we take $h(z) = \frac{1}{1-z}$, $z \in D(0, 1)$, then

$$\lambda_r(t) = \ln \left[\frac{1 - r \exp(-t)}{1 - r} \right] \tag{30}$$

is nonnegative, increasing and subadditive on $[0, \infty)$ for any $r \in (0, 1)$.

If we take $h(z) = \exp(z)$, $z \in \mathbb{C}$ then

$$\lambda_r(t) = r [1 - \exp(-t)] \tag{31}$$

is nonnegative, increasing and subadditive on $[0, \infty)$ for any $r > 0$.

Corollary 2.12 ([37]). *Let $h(z) = \sum_{n=0}^{\infty} a_n z^n$ a power series with nonnegative coefficients $a_n \geq 0$ for all $n \in \mathbb{N}$ and convergent on the open disk $D(0, R)$ with $R > 0$ or $R = \infty$ and $r \in (0, R)$. For a mapping $f : C \rightarrow \mathbb{R}$ defined on convex subset C of a linear space X , the following statements are equivalent:*

(i) *The function f is λ_r -convex with $\lambda_r : [0, \infty) \rightarrow [0, \infty)$,*

$$\lambda_r(t) := \ln \left[\frac{h(r)}{h(r \exp(-t))} \right];$$

(ii) *We have the inequality*

$$\left[\frac{h(r)}{h(r \exp(-\alpha - \beta))} \right]^{f\left(\frac{\alpha x + \beta y}{\alpha + \beta}\right)} \leq \left[\frac{h(r)}{h(r \exp(-\alpha))} \right]^{f(x)} \left[\frac{h(r)}{h(r \exp(-\beta))} \right]^{f(y)} \tag{32}$$

for any $\alpha, \beta \geq 0$ with $\alpha + \beta > 0$ and $x, y \in C$.

(iii) *We have the inequality*

$$\frac{[h(r \exp(-\alpha))]^{f(x)} [h(r \exp(-\beta))]^{f(y)}}{[h(r \exp(-\alpha - \beta))]^{f\left(\frac{\alpha x + \beta y}{\alpha + \beta}\right)}} \leq [h(r)]^{f(x) + f(y) - f\left(\frac{\alpha x + \beta y}{\alpha + \beta}\right)} \tag{33}$$

for any $\alpha, \beta \geq 0$ with $\alpha + \beta > 0$ and $x, y \in C$.

Remark 2.13. We observe that, in the case when

$$\lambda_r(t) = r[1 - \exp(-t)], t \geq 0$$

then the function f is λ_r -convex on convex subset C of a linear space X iff

$$f\left(\frac{\alpha x + \beta y}{\alpha + \beta}\right) \leq \frac{[1 - \exp(-\alpha)]f(x) + [1 - \exp(-\beta)]f(y)}{1 - \exp(-\alpha - \beta)} \tag{34}$$

for any $\alpha, \beta \geq 0$ with $\alpha + \beta > 0$ and $x, y \in C$.

We observe that this definition is independent of $r > 0$.

The inequality (34) is equivalent with

$$f\left(\frac{\alpha x + \beta y}{\alpha + \beta}\right) \leq \frac{\exp(\beta)[\exp(\alpha) - 1]f(x) + \exp(\alpha)[\exp(\beta) - 1]f(y)}{\exp(\alpha + \beta) - 1} \tag{35}$$

for any $\alpha, \beta \geq 0$ with $\alpha + \beta > 0$ and $x, y \in C$.

We also can introduce the following mapping $k_{x,y} : [0, 1] \rightarrow \mathbb{R}$

$$k_{x,y}(t) := \frac{1}{2} [f(tx + (1-t)y) + f((1-t)x + ty)]$$

for $x, y \in C, x \neq y$.

The following result holds [37]:

Theorem 2.14. Let $f : C \rightarrow [0, \infty)$ be a λ -convex function on C . Assume that $x, y \in C$ with $x \neq y$.

(i) We have the equality

$$k_{x,y}(1-t) = k_{x,y}(t) \text{ for all } t \in [0, 1];$$

(ii) The mapping $k_{x,y}$ is λ -convex on $[0, 1]$;

(iii) One has the inequalities

$$k_{x,y}(t) \leq \frac{\lambda(t) + \lambda(1-t)}{\lambda(1)} \cdot \frac{f(x) + f(y)}{2} \tag{36}$$

and

$$\frac{\lambda(2\alpha)}{2\lambda(\alpha)} f\left(\frac{x+y}{2}\right) \leq k_{x,y}(t) \tag{37}$$

for all $t \in [0, 1]$ and $\alpha > 0$.

(iv) Let $y, x \in C$ with $y \neq x$ and assume that the mappings $[0, 1] \ni t \mapsto f[(1-t)x + ty]$ and λ are Lebesgue integrable on $[0, 1]$, then we have the Hermite-Hadamard type inequalities

$$\frac{\lambda(2\alpha)}{2\lambda(\alpha)} f\left(\frac{x+y}{2}\right) \leq \int_0^1 f((1-t)x + ty) dt \leq \frac{f(x) + f(y)}{\lambda(1)} \int_0^1 \lambda(t) dt \tag{38}$$

for any $\alpha > 0$.

Corollary 2.15. *If $f : I \rightarrow [0, \infty)$ is an λ -convex function on an interval I of real numbers with $\lambda \in L[0, 1]$ and $f \in L[a, b]$ with $a, b \in I, a < b$, then*

$$\frac{\lambda(2\alpha)}{2\lambda(\alpha)} f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(u) du \leq \frac{f(a)+f(b)}{\lambda(1)} \int_0^1 \lambda(t) dt \quad (39)$$

for any $\alpha > 0$.

3. New Results for Čebyšev Functional

We have the following result:

Theorem 3.1. *Let $f : I \rightarrow \mathbb{C}$ be a differentiable function on $\overset{\circ}{I}$, the interior of the interval I , $[a, b] \subset \overset{\circ}{I}$ and $g : [a, b] \rightarrow \mathbb{C}$ is an integrable function on $[a, b]$.*

(i) *If $|f'|$ is λ -convex integrable on $[a, b]$ and λ is integrable on $[0, 1]$, then*

$$|C(f, g)| \leq \frac{|f'(a)| + |f'(b)|}{2\lambda(1)} \int_0^1 \lambda(t) dt \int_a^b \left| g(t) - \frac{1}{b-a} \int_a^b g(s) ds \right| dt. \quad (40)$$

(ii) *If $|f'|$ is h -convex integrable on $[a, b]$ and h is integrable on $(0, 1)$, then*

$$|C(f, g)| \leq \frac{|f'(a)| + |f'(b)|}{2} \int_0^1 h(t) dt \int_a^b \left| g(t) - \frac{1}{b-a} \int_a^b g(s) ds \right| dt. \quad (41)$$

Proof. We use Sonin's identity

$$C(f, g) = \frac{1}{b-a} \int_a^b (f(t) - \mu) \left[g(t) - \frac{1}{b-a} \int_a^b g(s) ds \right] dt, \quad (42)$$

for $\mu = \frac{f(a)+f(b)}{2}$ to get

$$C(f, g) = \frac{1}{b-a} \int_a^b \left[f(t) - \frac{f(a)+f(b)}{2} \right] \left[g(t) - \frac{1}{b-a} \int_a^b g(s) ds \right] dt. \quad (43)$$

Since f is differentiable, then we have

$$\begin{aligned} f(t) - \frac{f(a)+f(b)}{2} &= \frac{1}{2} [f(t) - f(a) + f(t) - f(b)] \\ &= \frac{1}{2} \left[\int_a^t f'(s) ds - \int_t^b f'(s) ds \right] \\ &= \frac{1}{2} \int_a^b \operatorname{sgn}(t-s) f'(s) ds \end{aligned}$$

for any $t \in [a, b]$.

Therefore we have the representation:

$$C(f, g) = \frac{1}{2(b-a)} \int_a^b \left(\int_a^b \operatorname{sgn}(t-s) f'(s) ds \right) \times \left(g(t) - \frac{1}{b-a} \int_a^b g(s) ds \right) dt. \quad (44)$$

Taking the modulus we have

$$\begin{aligned} |C(f, g)| &\leq \frac{1}{2(b-a)} \int_a^b \left| \int_a^b \operatorname{sgn}(t-s) f'(s) ds \right| \\ &\times \left| g(t) - \frac{1}{b-a} \int_a^b g(s) ds \right| dt \\ &\leq \frac{1}{2(b-a)} \int_a^b \left(\int_a^b |\operatorname{sgn}(t-s) f'(s)| ds \right) \\ &\times \left| g(t) - \frac{1}{b-a} \int_a^b g(s) ds \right| dt \\ &= \frac{1}{2(b-a)} \int_a^b \left(\int_a^b |f'(s)| ds \right) \left| g(t) - \frac{1}{b-a} \int_a^b g(s) ds \right| dt \\ &= \frac{1}{2(b-a)} \int_a^b |f'(s)| ds \int_a^b \left| g(t) - \frac{1}{b-a} \int_a^b g(s) ds \right| dt. \end{aligned} \quad (45)$$

(i) Since $|f'|$ is λ -convex integrable on $[a, b]$, then by Corollary 2.15 we have

$$\frac{1}{b-a} \int_a^b |f'(s)| ds \leq \frac{|f'(a)| + |f'(b)|}{2\lambda(1)} \int_0^1 \lambda(t) dt$$

and by (45) we get (40).

(ii) Follows by (23) and the details are omitted. \square

With the notations from the introduction we have:

Corollary 3.2. *Let $f : I \rightarrow \mathbb{C}$ be a differentiable function on \dot{I} and $[a, b] \subset \dot{I}$. If $|f'|$ is λ -convex integrable on $[a, b]$ and λ is integrable on $[0, 1]$, then*

$$\begin{aligned} D^2(f), E^2(f) &\leq \frac{|f'(a)| + |f'(b)|}{2\lambda(1)} \int_0^1 \lambda(t) dt \\ &\times \int_a^b \left| f(t) - \frac{1}{b-a} \int_a^b f(s) ds \right| dt, \end{aligned} \quad (46)$$

and

$$G^2(f) \leq \frac{|f'(a)| + |f'(b)|}{2\lambda(1)} \int_0^1 \lambda(t) dt \int_a^b \left| |f(t)| - \frac{1}{b-a} \int_a^b |f(s)| ds \right| dt. \quad (47)$$

A similar result holds for h -convex functions.

Remark 3.3. Let $f : I \rightarrow \mathbb{C}$ be a differentiable function on \dot{I} , $[a, b] \subset \dot{I}$ and $g : [a, b] \rightarrow \mathbb{C}$ is an integrable function on $[a, b]$.

If $|f'|$ is convex on $[a, b]$, then

$$|C(f, g)| \leq \frac{|f'(a)| + |f'(b)|}{4} \int_a^b \left| g(t) - \frac{1}{b-a} \int_a^b g(s) ds \right| dt. \tag{48}$$

If $|f'|$ is of P -type on $[a, b]$, then

$$|C(f, g)| \leq \frac{|f'(a)| + |f'(b)|}{2} \int_a^b \left| g(t) - \frac{1}{b-a} \int_a^b g(s) ds \right| dt. \tag{49}$$

If $|f'|$ is Breckner s -convex on $[a, b]$, for $s \in (0, 1)$, then

$$|C(f, g)| \leq \frac{|f'(a)| + |f'(b)|}{2(s+1)} \int_a^b \left| g(t) - \frac{1}{b-a} \int_a^b g(s) ds \right| dt. \tag{50}$$

If $|f'|$ is of s -Godunova-Levin type, with $s \in [0, 1)$, then

$$|C(f, g)| \leq \frac{|f'(a)| + |f'(b)|}{2(1-s)} \int_a^b \left| g(t) - \frac{1}{b-a} \int_a^b g(s) ds \right| dt. \tag{51}$$

Remark 3.4. We notice, from the proof of Theorem 3.1, if $|f'|$ satisfies the second Hemite-Hadamard inequality with a certain term $R(|f'(a)|, |f'(b)|)$, i.e.

$$\frac{1}{b-a} \int_a^b |f'(u)| du \leq R(|f'(a)|, |f'(b)|),$$

then we have the inequality

$$|C(f, g)| \leq R(|f'(a)|, |f'(b)|) \int_a^b \left| g(t) - \frac{1}{b-a} \int_a^b g(s) ds \right| dt. \tag{52}$$

The case of p -norm of the deviation

$$\left| f - \frac{1}{b-a} \int_a^b f(s) ds \right|$$

is as follows:

Theorem 3.5. Let $f : I \rightarrow \mathbb{C}$ be a differentiable function on \dot{I} , $[a, b] \subset \dot{I}$ and $g : [a, b] \rightarrow \mathbb{C}$ is an integrable function on $[a, b]$. Assume that $p, q > 1$ with $\frac{1}{p} + \frac{1}{q} = 1$.

(i) If $|f'|^q$ is λ -convex integrable on $[a, b]$ and λ is integrable on $[0, 1]$, then

$$|C(f, g)| \leq (b-a) \left[\frac{|f'(a)|^q + |f'(b)|^q}{2\lambda(1)} \int_0^1 \lambda(t) dt \right]^{1/q} \quad (53)$$

$$\times \left[\frac{1}{b-a} \int_a^b \left| g(t) - \frac{1}{b-a} \int_a^b g(s) ds \right|^p dt \right]^{1/p}.$$

(ii) If $|f'|^q$ is h -convex integrable on $[a, b]$ and h is integrable on $(0, 1)$, then

$$|C(f, g)| \leq (b-a) \left[\frac{|f'(a)|^q + |f'(b)|^q}{2} \int_0^1 h(t) dt \right]^{1/q} \quad (54)$$

$$\times \left[\frac{1}{b-a} \int_a^b \left| g(t) - \frac{1}{b-a} \int_a^b g(s) ds \right|^p dt \right]^{1/p}.$$

Proof. Making use of Hölder's integral inequality, we have

$$|C(f, g)| \leq \frac{1}{2(b-a)} \int_a^b \left| \int_a^b \operatorname{sgn}(t-s) f'(s) ds \right| \quad (55)$$

$$\times \left| g(t) - \frac{1}{b-a} \int_a^b g(s) ds \right| dt$$

$$\leq \frac{1}{2(b-a)} \left(\int_a^b \left| \int_a^b \operatorname{sgn}(t-s) f'(s) ds \right|^q dt \right)^{1/q}$$

$$\times \left(\int_a^b \left| g(t) - \frac{1}{b-a} \int_a^b g(s) ds \right|^p dt \right)^{1/p}.$$

Observe that, by Jensen's integral inequality for $q > 1$, we have

$$\left| \frac{\int_a^b \operatorname{sgn}(t-s) f'(s) ds}{b-a} \right|^q \leq \frac{\int_a^b |\operatorname{sgn}(t-s) f'(s)|^q ds}{b-a}$$

$$= \frac{\int_a^b |f'(s)|^q ds}{b-a},$$

which shows that

$$\left| \int_a^b \operatorname{sgn}(t-s) f'(s) ds \right|^q \leq (b-a)^{q-1} \int_a^b |f'(s)|^q ds$$

for any $t \in [a, b]$.

Therefore,

$$\left(\int_a^b \left| \int_a^b \operatorname{sgn}(t-s) f'(s) ds \right|^q dt \right)^{1/q} \leq \left[(b-a)^q \int_a^b |f'(s)|^q ds \right]^{1/q}$$

$$= (b-a) \left[\int_a^b |f'(s)|^q ds \right]^{1/q}$$

and by (55) we get

$$|C(f, g)| \leq \frac{1}{2} \left(\int_a^b |f'(s)|^q ds \right)^{1/q} \left(\int_a^b \left| g(t) - \frac{1}{b-a} \int_a^b g(s) ds \right|^p dt \right)^{1/p}.$$

(i) Since $|f'|^q$ is λ -convex integrable on $[a, b]$, then by Corollary 2.15 we have

$$\frac{1}{b-a} \int_a^b |f'(s)|^q ds \leq \frac{|f'(a)|^q + |f'(b)|^q}{2\lambda(1)} \int_0^1 \lambda(t) dt$$

and by (55) we get (53).

(ii) Follows by (23) and the details are omitted. □

The case $p = q = 2$ is of interest.

Corollary 3.6. *Let $f : I \rightarrow \mathbb{C}$ be a differentiable function on \mathring{I} , $[a, b] \subset \mathring{I}$ and $g : [a, b] \rightarrow \mathbb{C}$ is an integrable function on $[a, b]$.*

If $|f'|^2$ is λ -convex integrable on $[a, b]$ and λ is integrable on $[0, 1]$, then

$$|C(f, g)| \leq (b-a) \left[\frac{|f'(a)|^2 + |f'(b)|^2}{2\lambda(1)} \int_0^1 \lambda(t) dt \right]^{1/2} D(g), \quad (56)$$

where

$$D(g) = [C(g, \bar{g})]^{1/2} = \left[\frac{1}{b-a} \int_a^b |g(t)|^2 dt - \left| \frac{1}{b-a} \int_a^b g(t) dt \right|^2 \right]^{1/2}.$$

If $|f'|^2$ is h -convex integrable on $[a, b]$ and h is integrable on $(0, 1)$, then a similar inequality is valid.

The following particular cases are of interest as well:

Corollary 3.7. *Let $f : I \rightarrow \mathbb{C}$ be a differentiable function on \mathring{I} , $[a, b] \subset \mathring{I}$ and $g : [a, b] \rightarrow \mathbb{C}$ is an integrable function on $[a, b]$. Assume that $p, q > 1$ and $\frac{1}{p} + \frac{1}{q} = 1$. If $|f'|^q$ is λ -convex integrable on $[a, b]$ and λ is integrable on $[0, 1]$, then*

$$D^2(f), E^2(f) \leq (b-a) \left[\frac{|f'(a)|^q + |f'(b)|^q}{2\lambda(1)} \int_0^1 \lambda(t) dt \right]^{1/q} \times \left[\frac{1}{b-a} \int_a^b \left| f(t) - \frac{1}{b-a} \int_a^b f(s) ds \right|^p dt \right]^{1/p}, \quad (57)$$

and

$$G^2(f) \leq (b-a) \left[\frac{|f'(a)|^q + |f'(b)|^q}{2\lambda(1)} \int_0^1 \lambda(t) dt \right]^{1/q} \quad (58)$$

$$\times \left[\frac{1}{b-a} \int_a^b \left| |f(t)| - \frac{1}{b-a} \int_a^b |f(s)| ds \right|^p dt \right]^{1/p}.$$

In particular, if $|f'|^2$ is λ -convex integrable on $[a, b]$ and λ is integrable on $[0, 1]$, then

$$D^2(f), E^2(f) \leq (b-a) \left[\frac{|f'(a)|^2 + |f'(b)|^2}{2\lambda(1)} \int_0^1 \lambda(t) dt \right]^{1/2} D(f), \quad (59)$$

and

$$G^2(f) \leq (b-a) \left[\frac{|f'(a)|^2 + |f'(b)|^2}{2\lambda(1)} \int_0^1 \lambda(t) dt \right]^{1/2} D(|f|). \quad (60)$$

The first inequality in (59) is equivalent to:

$$D(f) \leq (b-a) \left[\frac{|f'(a)|^2 + |f'(b)|^2}{2\lambda(1)} \int_0^1 \lambda(t) dt \right]^{1/2}. \quad (61)$$

Remark 3.8. Let $f : I \rightarrow \mathbb{C}$ be a differentiable function on \mathring{I} , $[a, b] \subset \mathring{I}$ and $g : [a, b] \rightarrow \mathbb{C}$ is an integrable function on $[a, b]$. Assume that $p, q > 1$ and $\frac{1}{p} + \frac{1}{q} = 1$.

If $|f'|^q$ is convex on $[a, b]$, then

$$|C(f, g)| \leq (b-a) \left[\frac{|f'(a)|^q + |f'(b)|^q}{4} \right]^{1/q} \quad (62)$$

$$\times \left[\frac{1}{b-a} \int_a^b \left| g(t) - \frac{1}{b-a} \int_a^b g(s) ds \right|^p dt \right]^{1/p}.$$

If $|f'|^q$ is of P -type on $[a, b]$, then

$$|C(f, g)| \leq (b-a) \left[\frac{|f'(a)|^q + |f'(b)|^q}{2} \right]^{1/q} \quad (63)$$

$$\times \left[\frac{1}{b-a} \int_a^b \left| g(t) - \frac{1}{b-a} \int_a^b g(s) ds \right|^p dt \right]^{1/p}.$$

If $|f'|^q$ is Breckner s -convex on $[a, b]$, for $s \in (0, 1)$, then

$$|C(f, g)| \leq (b-a) \left[\frac{|f'(a)|^q + |f'(b)|^q}{2(s+1)} \right]^{1/q} \times \left[\frac{1}{b-a} \int_a^b \left| g(t) - \frac{1}{b-a} \int_a^b g(s) ds \right|^p dt \right]^{1/p}. \tag{64}$$

If $|f'|^q$ is of s -Godunova-Levin type, with $s \in [0, 1)$, then

$$|C(f, g)| \leq (b-a) \left[\frac{|f'(a)|^q + |f'(b)|^q}{2(1-s)} \right]^{1/q} \times \left[\frac{1}{b-a} \int_a^b \left| g(t) - \frac{1}{b-a} \int_a^b g(s) ds \right|^p dt \right]^{1/p}. \tag{65}$$

4. Application for Riemann-Stieltjes Integral

The following representation is of interest in itself. The result was firstly obtained in [27] (see also [29]). For the sake a completeness we give here a short proof as well.

Lemma 4.1. *If $v : [a, b] \rightarrow \mathbb{C}$ is continuous (of bounded variation) on $[a, b]$ and $h : [a, b] \rightarrow \mathbb{C}$ is of bounded variation (continuous) on $[a, b]$, then we have the identity*

$$\frac{v(b) \int_a^b (t-a) dh(t) + v(a) \int_a^b (b-t) dh(t)}{b-a} - \int_a^b v(t) dh(t) = \int_a^b h(t) dv(t) - \frac{v(b) - v(a)}{b-a} \int_a^b h(t) dt. \tag{66}$$

Proof. Integrating by parts in the Riemann-Stieltjes integral we have

$$\begin{aligned} & \frac{v(b) \int_a^b (t-a) dh(t) + v(a) \int_a^b (b-t) dh(t)}{b-a} - \int_a^b v(t) dh(t) \\ &= \int_a^b \left[\frac{v(b)(t-a) + v(a)(b-t)}{b-a} - v(t) \right] dh(t) \\ &= \left[\frac{(t-a)v(b) + (b-t)v(a)}{b-a} - v(t) \right] h(t) \Big|_a^b \\ & \quad - \int_a^b h(t) d \left[\frac{(t-a)v(b) + (b-t)v(a)}{b-a} - v(t) \right] = \end{aligned} \tag{67}$$

$$\begin{aligned} &= [v(b) - v(a)]h(b) - [v(a) - v(a)]h(a) \\ &\quad - \int_a^b h(t) \left[\frac{v(b) - v(a)}{b - a} dt - dv(t) \right] \\ &= \int_a^b h(t) dv(t) - \frac{v(b) - v(a)}{b - a} \int_a^b h(t) dt \end{aligned}$$

and the identity is proven. □

We can provide now the following application for Riemann-Stieltjes integral:

Proposition 4.2. *If $v : I \rightarrow \mathbb{C}$ is twice differentiable on the interior of the interval I denoted \mathring{I} and $[a, b] \subset \mathring{I}$. If $|v''|$ is λ -convex integrable on $[a, b]$ and λ is integrable on $[0, 1]$, then for $h : [a, b] \rightarrow \mathbb{C}$ integrable on $[a, b]$, we have the inequalities*

$$\begin{aligned} &\left| \frac{v(b) \int_a^b (t - a) dh(t) + v(a) \int_a^b (b - t) dh(t)}{b - a} - \int_a^b v(t) dh(t) \right| \tag{68} \\ &\leq \frac{|v''(a)| + |v''(b)|}{2\lambda(1)} (b - a) \int_0^1 \lambda(t) dt \int_a^b \left| h(t) - \frac{1}{b - a} \int_a^b h(s) ds \right| dt \end{aligned}$$

Proof. From (66) we have

$$\begin{aligned} &\frac{v(b) \int_a^b (t - a) dh(t) + v(a) \int_a^b (b - t) dh(t)}{b - a} - \int_a^b v(t) dh(t) \tag{69} \\ &= \int_a^b h(t) v'(t) dt - \frac{v(b) - v(a)}{b - a} \int_a^b h(t) dt = (b - a)C(v', h). \end{aligned}$$

Since $|v''|$ is λ -convex integrable on $[a, b]$, then by applying Theorem 3.1 for $f = v'$ and $g = h$ we deduce the desired result (68). □

Remark 4.3. If $v : I \rightarrow \mathbb{C}$ is twice differentiable on \mathring{I} , $[a, b] \subset \mathring{I}$ and $g : [a, b] \rightarrow \mathbb{C}$ is an integrable function on $[a, b]$.

If $|v''|$ is convex on $[a, b]$, then

$$\begin{aligned} &\left| \frac{v(b) \int_a^b (t - a) dh(t) + v(a) \int_a^b (b - t) dh(t)}{b - a} - \int_a^b v(t) dh(t) \right| \tag{70} \\ &\leq \frac{|v''(a)| + |v''(b)|}{4} (b - a) \int_a^b \left| h(t) - \frac{1}{b - a} \int_a^b h(s) ds \right| dt. \end{aligned}$$

If $|v''|$ is of P -type on $[a, b]$, then

$$\begin{aligned} & \left| \frac{v(b) \int_a^b (t-a) dh(t) + v(a) \int_a^b (b-t) dh(t)}{b-a} - \int_a^b v(t) dh(t) \right| \quad (71) \\ & \leq \frac{|v''(a)| + |v''(b)|}{2} (b-a) \int_a^b \left| h(t) - \frac{1}{b-a} \int_a^b h(s) ds \right| dt. \end{aligned}$$

If $|v''|$ is Breckner s -convex on $[a, b]$, for $s \in (0, 1)$, then

$$\begin{aligned} & \left| \frac{v(b) \int_a^b (t-a) dh(t) + v(a) \int_a^b (b-t) dh(t)}{b-a} - \int_a^b v(t) dh(t) \right| \quad (72) \\ & \leq \frac{|v''(a)| + |v''(b)|}{2(s+1)} (b-a) \int_a^b \left| h(t) - \frac{1}{b-a} \int_a^b h(s) ds \right| dt. \end{aligned}$$

If $|v''|$ is of s -Godunova-Levin type, with $s \in [0, 1)$, then

$$\begin{aligned} & \left| \frac{v(b) \int_a^b (t-a) dh(t) + v(a) \int_a^b (b-t) dh(t)}{b-a} - \int_a^b v(t) dh(t) \right| \quad (73) \\ & \leq \frac{|v''(a)| + |v''(b)|}{2(1-s)} (b-a) \int_a^b \left| h(t) - \frac{1}{b-a} \int_a^b h(s) ds \right| dt. \end{aligned}$$

Similar results may be stated if $|f'|^q$ is λ -convex integrable on $[a, b]$ and λ is integrable on $[0, 1]$. However the details are not provided here.

5. Applications for Selfadjoint Operators

We denote by $\mathcal{B}(H)$ the Banach algebra of all bounded linear operators on a complex Hilbert space $(H; \langle \cdot, \cdot \rangle)$. Let $A \in \mathcal{B}(H)$ be selfadjoint and let φ_λ be defined for all $\lambda \in \mathbb{R}$ as follows

$$\varphi_\lambda(s) := \begin{cases} 1, & \text{for } -\infty < s \leq \lambda, \\ 0, & \text{for } \lambda < s < +\infty. \end{cases}$$

Then for every $\lambda \in \mathbb{R}$ the operator

$$E_\lambda := \varphi_\lambda(A) \quad (74)$$

is a projection which reduces A .

The properties of these projections are collected in the following fundamental result concerning the spectral representation of bounded selfadjoint operators in Hilbert spaces, see for instance [56, p. 256]:

Theorem 5.1 (Spectral Representation Theorem). *Let A be a bounded self-adjoint operator on the Hilbert space H and let $m = \min \{ \lambda \mid \lambda \in Sp(A) \} =: \min Sp(A)$ and $M = \max \{ \lambda \mid \lambda \in Sp(A) \} =: \max Sp(A)$. Then there exists a family of projections $\{E_\lambda\}_{\lambda \in \mathbb{R}}$, called the spectral family of A , with the following properties*

- a) $E_\lambda \leq E_{\lambda'}$ for $\lambda \leq \lambda'$;
- b) $E_{m-0} = 0, E_M = I$ and $E_{\lambda+0} = E_\lambda$ for all $\lambda \in \mathbb{R}$;
- c) We have the representation

$$A = \int_{m-0}^M \lambda dE_\lambda. \quad (75)$$

More generally, for every continuous complex-valued function φ defined on \mathbb{R} and for every $\varepsilon > 0$ there exists a $\delta > 0$ such that

$$\left\| \varphi(A) - \sum_{k=1}^n \varphi(\lambda'_k) [E_{\lambda_k} - E_{\lambda_{k-1}}] \right\| \leq \varepsilon \quad (76)$$

whenever

$$\begin{cases} \lambda_0 < m = \lambda_1 < \dots < \lambda_{n-1} < \lambda_n = M, \\ \lambda_k - \lambda_{k-1} \leq \delta \text{ for } 1 \leq k \leq n, \\ \lambda'_k \in [\lambda_{k-1}, \lambda_k] \text{ for } 1 \leq k \leq n \end{cases} \quad (77)$$

this means that

$$\varphi(A) = \int_{m-0}^M \varphi(\lambda) dE_\lambda, \quad (78)$$

where the integral is of Riemann-Stieltjes type.

Corollary 5.2. *With the assumptions of Theorem 5.1 for A, E_λ and φ we have the representations*

$$\varphi(A)x = \int_{m-0}^M \varphi(\lambda) dE_\lambda x \text{ for all } x \in H \quad (79)$$

and

$$\langle \varphi(A)x, y \rangle = \int_{m-0}^M \varphi(\lambda) d \langle E_\lambda x, y \rangle \text{ for all } x, y \in H. \quad (80)$$

In particular,

$$\langle \varphi(A)x, x \rangle = \int_{m-0}^M \varphi(\lambda) d \langle E_\lambda x, x \rangle \text{ for all } x \in H. \quad (81)$$

Moreover, we have the equality

$$\|\varphi(A)x\|^2 = \int_{m-0}^M |\varphi(\lambda)|^2 d\|E_\lambda x\|^2 \text{ for all } x \in H. \tag{82}$$

The next result shows that it is legitimate to talk about "the" spectral family of the bounded selfadjoint operator A since it is uniquely determined by the requirements a), b) and c) in Theorem 5.1, see for instance [56, p. 258]:

Theorem 5.3. *Let A be a bounded selfadjoint operator on the Hilbert space H and let $m = \min Sp(A)$ and $M = \max Sp(A)$. If $\{F_\lambda\}_{\lambda \in \mathbb{R}}$ is a family of projections satisfying the requirements a), b) and c) in Theorem 5.1, then $F_\lambda = E_\lambda$ for all $\lambda \in \mathbb{R}$ where E_λ is defined by (74).*

By the above two theorems, the spectral family $\{E_\lambda\}_{\lambda \in \mathbb{R}}$ uniquely determines and in turn is uniquely determined by the bounded selfadjoint operator A .

We have:

Theorem 5.4. *Let A be a bounded selfadjoint operator on the Hilbert space H and let $m = \min \{\lambda \mid \lambda \in Sp(A)\} =: \min Sp(A)$ and $M = \max \{\lambda \mid \lambda \in Sp(A)\} =: \max Sp(A)$. Consider also the spectral family $\{E_\lambda\}_{\lambda \in \mathbb{R}}$ of A .*

If $f : I \rightarrow \mathbb{C}$ is twice differentiable on \mathring{I} and $[m, M] \subset \mathring{I}$, $|f''|$ is λ -convex integrable on $[m, M]$ and λ is integrable on $[0, 1]$, then we have the inequalities

$$\begin{aligned} & \left| \left\langle \left[\frac{f(m)(M1_H - A) + f(M)(A - m1_H)}{M - m} \right] x, y \right\rangle - \langle f(A)x, y \rangle \right| \tag{83} \\ & \leq \frac{|f''(m)| + |f''(M)|}{2\lambda(1)} (M - m) \int_0^1 \lambda(t) dt \\ & \times \int_{m-0}^M \left| \langle E_t x, y \rangle - \frac{1}{M - m} \int_{m-0}^M \langle E_s x, y \rangle ds \right| dt \\ & \leq \frac{|f''(m)| + |f''(M)|}{4\lambda(1)} (M - m)^2 \|x\| \|y\| \int_0^1 \lambda(t) dt \end{aligned}$$

for any $x, y \in H$.

Proof. Let $x, y \in H$ and consider $h : \mathbb{R} \rightarrow \mathbb{C}$, $h(t) := \langle E_t x, y \rangle$.

If we use the first inequality in (68) for the interval $[m - \varepsilon, M]$ with small

$\varepsilon > 0$, we have

$$\begin{aligned} & \left| \frac{f(M) \int_{m-\varepsilon}^M (t-m+\varepsilon) d \langle E_t x, y \rangle + f(m-\varepsilon) \int_{m-\varepsilon}^M (M-t) d \langle E_t x, y \rangle}{M-m+\varepsilon} \right. \\ & \left. - \int_{m-\varepsilon}^M f(t) d \langle E_t x, y \rangle \right| \\ & \leq \frac{|f''(m-\varepsilon)| + |f''(M)|}{2\lambda(1)} (M-m+\varepsilon)^2 \int_0^1 \lambda(t) dt \\ & \times \frac{1}{M-m+\varepsilon} \int_{m-\varepsilon}^M \left| \langle E_t x, y \rangle - \frac{1}{M-m+\varepsilon} \int_{m-\varepsilon}^M \langle E_s x, y \rangle ds \right| dt. \end{aligned} \tag{84}$$

Taking the limit over $\varepsilon \rightarrow 0+$ and using the Spectral representation theorem, we have

$$\begin{aligned} & \left| \left\langle \left[\frac{f(m)(M1_H - A) + f(M)(A - m1_H)}{M-m} \right] x, y \right\rangle - \langle f(A)x, y \rangle \right| \\ & \leq \frac{|f''(m)| + |f''(M)|}{2\lambda(1)} (M-m)^2 \int_0^1 \lambda(t) dt \\ & \times \frac{1}{M-m} \int_{m-0}^M \left| \langle E_t x, y \rangle - \frac{1}{M-m} \int_{m-0}^M \langle E_s x, y \rangle ds \right| dt \end{aligned} \tag{85}$$

for any $x, y \in H$.

By the Schwarz inequality in H we have that

$$\begin{aligned} & \int_{m-0}^M \left| \langle E_t x, y \rangle - \frac{1}{M-m} \int_{m-0}^M \langle E_s x, y \rangle ds \right| dt \\ & = \int_{m-0}^M \left| \left\langle \left[E_t x - \frac{1}{M-m} \int_{m-0}^M E_s x ds \right], y \right\rangle \right| dt \\ & \leq \|y\| \int_{m-0}^M \left\| E_t x - \frac{1}{M-m} \int_{m-0}^M E_s x ds \right\| dt \end{aligned} \tag{86}$$

for any $x, y \in H$.

On utilizing the Cauchy-Buniakovski-Schwarz integral inequality we may state that

$$\begin{aligned} & \int_{m-0}^M \left\| E_t x - \frac{1}{M-m} \int_{m-0}^M E_s x ds \right\| dt \\ & \leq (M-m)^{1/2} \left(\int_{m-0}^M \left\| E_t x - \frac{1}{M-m} \int_{m-0}^M E_s x ds \right\|^2 dt \right)^{1/2} \end{aligned} \tag{87}$$

for any $x \in H$.

Observe that the following equalities of interest hold and they can be easily proved by direct calculations

$$\begin{aligned} & \frac{1}{M-m} \int_{m-0}^M \left\| E_t x - \frac{1}{M-m} \int_{m-0}^M E_s x ds \right\|^2 dt \\ &= \frac{1}{M-m} \int_{m-0}^M \|E_t x\|^2 dt - \left\| \frac{1}{M-m} \int_{m-0}^M E_s x ds \right\|^2 \end{aligned} \tag{88}$$

and

$$\begin{aligned} & \frac{1}{M-m} \int_{m-0}^M \|E_t x\|^2 dt - \left\| \frac{1}{M-m} \int_{m-0}^M E_s x ds \right\|^2 \\ &= \frac{1}{M-m} \int_{m-0}^M \left\langle E_t x - \frac{1}{M-m} \int_{m-0}^M E_s x ds, E_t x - \frac{1}{2} x \right\rangle dt \end{aligned} \tag{89}$$

for any $x \in H$.

By (87), (88) and (89) we get

$$\begin{aligned} & \int_{m-0}^M \left\| E_t x - \frac{1}{M-m} \int_{m-0}^M E_s x ds \right\| dt \\ & \leq (M-m)^{1/2} \left(\int_{m-0}^M \left\langle E_t x - \frac{1}{M-m} \int_{m-0}^M E_s x ds, E_t x - \frac{1}{2} x \right\rangle dt \right)^{1/2} \end{aligned} \tag{90}$$

for any $x \in H$.

On making use of the Schwarz inequality in H we also have

$$\begin{aligned} & \int_{m-0}^M \left\langle E_t x - \frac{1}{M-m} \int_{m-0}^M E_s x ds, E_t x - \frac{1}{2} x \right\rangle dt \\ & \leq \int_{m-0}^M \left\| E_t x - \frac{1}{M-m} \int_{m-0}^M E_s x ds \right\| \left\| E_t x - \frac{1}{2} x \right\| dt \\ & = \frac{1}{2} \|x\| \int_{m-0}^M \left\| E_t x - \frac{1}{M-m} \int_{m-0}^M E_s x ds \right\| dt, \end{aligned} \tag{91}$$

where we used the fact that E_t are projectors, and in this case we have

$$\left\| E_t x - \frac{1}{2} x \right\|^2 = \|E_t x\|^2 - \langle E_t x, x \rangle + \frac{1}{4} \|x\|^2 = \frac{1}{4} \|x\|^2$$

for any $t \in [m, M]$ for any $x \in H$.

From (90) and (91) we get

$$\begin{aligned} & \int_{m-0}^M \left\| E_t x - \frac{1}{M-m} \int_{m-0}^M E_s x ds \right\| dt \\ & \leq (M-m)^{1/2} \left(\frac{1}{2} \|x\| \int_{m-0}^M \left\| E_t x - \frac{1}{M-m} \int_{m-0}^M E_s x ds \right\| dt \right)^{1/2}, \end{aligned} \tag{92}$$

which is clearly equivalent with the following inequality of interest in itself

$$\int_{m-0}^M \left\| E_t x - \frac{1}{M-m} \int_{m-0}^M E_s x ds \right\| dt \leq \frac{1}{2} \|x\| (M-m) \tag{93}$$

for any $x \in H$.

From (86) we then get

$$\frac{1}{M-m} \int_{m-0}^M \left| \langle E_t x, y \rangle - \frac{1}{M-m} \int_{m-0}^M \langle E_s x, y \rangle ds \right| dt \leq \frac{1}{2} \|x\| \|y\|$$

for any $x, y \in H$. □

Remark 5.5. If $|f''|$ is convex on $[m, M]$, then we have the inequalities

$$\begin{aligned} & \left| \left\langle \left[\frac{f(m)(M1_H - A) + f(M)(A - m1_H)}{M-m} \right] x, y \right\rangle - \langle f(A)x, y \rangle \right| \tag{94} \\ & \leq \frac{|f''(m)| + |f''(M)|}{4} (M-m) \\ & \times \int_{m-0}^M \left| \langle E_t x, y \rangle - \frac{1}{M-m} \int_{m-0}^M \langle E_s x, y \rangle ds \right| dt \\ & \leq \frac{|f''(m)| + |f''(M)|}{8} (M-m)^2 \|x\| \|y\|, \end{aligned}$$

for any $x, y \in H$.

Example 5.6. a) Let A be a bounded selfadjoint operator on the Hilbert space H and $m = \min \{ \lambda \mid \lambda \in Sp(A) \}$ and $M = \max \{ \lambda \mid \lambda \in Sp(A) \} =: \max Sp(A)$. Consider also the spectral family $\{E_\lambda\}_{\lambda \in \mathbb{R}}$ of A . Then by Theorem 5.4 we have for $f(t) = t^p, p \geq 3$ that

$$\begin{aligned} & \left| \left\langle \left[\frac{m^p(M1_H - A) + M^p(A - m1_H)}{M-m} \right] x, y \right\rangle - \langle A^p x, y \rangle \right| \tag{95} \\ & \leq p(p-1) \frac{m^{p-2} + M^{p-2}}{4} (M-m) \\ & \times \int_{m-0}^M \left| \langle E_t x, y \rangle - \frac{1}{M-m} \int_{m-0}^M \langle E_s x, y \rangle ds \right| dt \\ & \leq p(p-1) \frac{m^{p-2} + M^{p-2}}{8} (M-m)^2 \|x\| \|y\| \end{aligned}$$

for any $x, y \in H$.

b) With the assumptions of a) and if $m > 0$, then by Theorem 5.4 we have for $f(t) = \ln t$, that

$$\begin{aligned} & \left| \left\langle \left[\frac{\ln m (M1_H - A) + \ln M (A - m1_H)}{M - m} \right] x, y \right\rangle - \langle \ln Ax, y \rangle \right| \quad (96) \\ & \leq \frac{m^2 + M^2}{4m^2M^2} (M - m) \int_{m-0}^M \left| \langle E_t x, y \rangle - \frac{1}{M - m} \int_{m-0}^M \langle E_s x, y \rangle ds \right| dt \\ & \leq \frac{m^2 + M^2}{8m^2M^2} (M - m)^2 \|x\| \|y\| \end{aligned}$$

for any $x, y \in H$.

Similar results may be stated if $|f'|^q$ is λ -convex integrable on $[a, b]$ and λ is integrable on $[0, 1]$. However the details are not provided here.

REFERENCES

- [1] M. Alomari - M. Darus, *The Hadamard's inequality for s-convex function*, Int. J. Math. Anal. (Ruse) 2 (13-16) (2008), 639–646.
- [2] M. Alomari - M. Darus, *Hadamard-type inequalities for s-convex functions*, Int. Math. Forum 3 (37-40) (2008), 1965–1975.
- [3] G. A. Anastassiou, *Univariate Ostrowski inequalities, revisited*, Monatsh. Math. 135 (3) (2002), 175–189.
- [4] N. S. Barnett - P. Cerone - S. S. Dragomir - M. R. Pinheiro - A. Sofo, *Ostrowski type inequalities for functions whose modulus of the derivatives are convex and applications*, Inequality Theory and Applications, Vol. 2 (Chinju/Masan, 2001), 19–32, Nova Sci. Publ., Hauppauge, NY, 2003. Preprint: RGMIA Res. Rep. Coll. 5 (2002), No. 2, Art. 1.
- [5] E. F. Beckenbach, *Convex functions*, Bull. Amer. Math. Soc. 54 (1948), 439–460.
- [6] M. Bombardelli - S. Varošanec, *Properties of h-convex functions related to the Hermite-Hadamard-Fejér inequalities*. Comp. Math. Appl. 58 (9) (2009), 1869–1877.
- [7] W. W. Breckner, *Stetigkeitsaussagen für eine Klasse verallgemeinerter konvexer Funktionen in topologischen linearen Räumen*, (German) Publ. Inst. Math. (Beograd) (N.S.) 23 (37) (1978), 13–20.
- [8] W. W. Breckner - G. Orbán, *Continuity properties of rationally s-convex mappings with values in an ordered topological linear space*. Universitatea "Babeş-Bolyai", Facultatea de Matematica, Cluj-Napoca, 1978.
- [9] P. Cerone - S. S. Dragomir, *Midpoint-type rules from an inequalities point of view*, Ed. G. A. Anastassiou, Handbook of Analytic-Computational Methods in Applied Mathematics, CRC Press, New York. 135–200.

- [10] P. Cerone - S. S. Dragomir, *New bounds for the three-point rule involving the Riemann-Stieltjes integrals*, in *Advances in Statistics Combinatorics and Related Areas*, C. Gulati, et al. (Eds.), World Science Publishing, 2002, 53–62.
- [11] P. Cerone - S. S. Dragomir, *New bounds for the Čebyšev functional*, *Appl. Math. Lett.* 18 (2005), 603–611.
- [12] P. Cerone - S. S. Dragomir, *A refinement of the Grüss inequality and applications*, *Tamkang J. Math.* 38 (1) (2007), 37–49. Preprint available at RGMIA Res. Rep. Coll., 5 (2) (2002), Art. 14.
- [13] P. Cerone - S. S. Dragomir - J. Roumeliotis, *Some Ostrowski type inequalities for n -time differentiable mappings and applications*, *Demonstratio Mathematica*, 32 (2) (1999), 697–712.
- [14] P. L. Chebychev, *Sur les expressions approximatives des intégrals définis par les autres prises entre les même limites*, *Proc. Math. Soc. Charkov* 2 (1882), 93–98.
- [15] X.-L. Cheng - J. Sun, *Note on the perturbed trapezoid inequality*, *J. Inequal. Pure Appl. Math.* 3 (2) (2002), Art. 29.
- [16] G. Cristescu, *Hadamard type inequalities for convolution of h -convex functions*, *Ann. Tiberiu Popoviciu Semin. Funct. Equ. Approx. Convexity* 8 (2010), 3–11.
- [17] S. S. Dragomir, *Ostrowski's inequality for monotonous mappings and applications*, *J. KSIAM* 3 (1) (1999), 127–135.
- [18] S. S. Dragomir, *The Ostrowski's integral inequality for Lipschitzian mappings and applications*, *Comp. Math. Appl.* 38 (1999), 33–37.
- [19] S. S. Dragomir, *On the Ostrowski's inequality for Riemann-Stieltjes integral*, *Korean J. Appl. Math.* 7 (2000), 477–485.
- [20] S. S. Dragomir, *On the Ostrowski's inequality for mappings of bounded variation and applications*, *Math. Ineq. & Appl.* 4 (1) (2001), 33–40.
- [21] S. S. Dragomir, *On the Ostrowski inequality for Riemann-Stieltjes integral $\int_a^b f(t) du(t)$ where f is of Hölder type and u is of bounded variation and applications*, *J. KSIAM* 5 (1) (2001), 35–45.
- [22] S. S. Dragomir, *Ostrowski type inequalities for isotonic linear functionals*, *J. Inequal. Pure & Appl. Math.* 3 (5) (2002), Art. 68.
- [23] S. S. Dragomir, *An inequality improving the first Hermite-Hadamard inequality for convex functions defined on linear spaces and applications for semi-inner products*, *J. Inequal. Pure Appl. Math.* 3 (2) (2002), Art. 31.
- [24] S. S. Dragomir, *An inequality improving the second Hermite-Hadamard inequality for convex functions defined on linear spaces and applications for semi-inner products*, *J. Inequal. Pure Appl. Math.* 3 (3) (2002), Art. 35.
- [25] S. S. Dragomir, *A survey on Cauchy-Buniakowski-Schwarz's type discrete inequality*, *J. Ineq. Pure & Appl. Math.* 4 (3) (2003), Art. 63.
- [26] S. S. Dragomir, *An Ostrowski like inequality for convex functions and applications*, *Revista Math. Complutense*, 16 (2) (2003), 373–382.

- [27] S. S. Dragomir, *Inequalities of Grüss type for the Stieltjes integral and applications*, Kragujevac J. Math. 26 (2004), 89–112.
- [28] S. S. Dragomir, *Bounds for the normalised Jensen functional*, Bull. Austral. Math. Soc. 74 (2006), 471–478.
- [29] S. S. Dragomir, *Inequalities for Stieltjes integrals with convex integrators and applications*, Appl. Math. Lett. 20 (2007), 123–130.
- [30] S. S. Dragomir, *Čebyšev's type inequalities for functions of selfadjoint operators in Hilbert spaces*, Preprint RGMIA Res. Rep. Coll. 11(e) (2008), Art. 9.
- [31] S. S. Dragomir, *Grüss' type inequalities for functions of selfadjoint operators in Hilbert spaces*, Preprint RGMIA Res. Rep. Coll. 11(e) (2008), Art. 11.
- [32] S. S. Dragomir, *Inequalities for the Čebyšev functional of two functions of selfadjoint operators in Hilbert spaces*, Preprint RGMIA Res. Rep. Coll. 11(e) (2008).
- [33] S. S. Dragomir, *New Grüss' type inequalities for functions of bounded variation and applications*, Appl. Math. Lett. 25 (10) (2012), 1475–1479.
- [34] S. S. Dragomir, *Operator Inequalities of the Jensen, Čebyšev and Grüss Type*, Springer Briefs in Mathematics, Springer, New York, 2012.
- [35] S. S. Dragomir, *Operator Inequalities of Ostrowski and Trapezoidal Type*, Springer Briefs in Mathematics. Springer, New York, 2012.
- [36] S. S. Dragomir, *Inequalities of Hermite-Hadamard type for h -convex functions on linear spaces*, Preprint RGMIA Res. Rep. Coll. 16 (2013), Art. 75.
- [37] S. S. Dragomir, *Inequalities of Hermite-Hadamard type for λ -convex functions on linear spaces*, Preprint RGMIA Res. Rep. Coll. 17 (2014), Art. 13.
- [38] S. S. Dragomir, *Discrete inequalities of Jensen type for λ -convex functions on linear spaces*, Preprint RGMIA Res. Rep. Coll. 17 (2014), Art. 15.
- [39] S. S. Dragomir, *Integral inequalities of Jensen type for λ -convex functions*, Preprint RGMIA Res. Rep. Coll. 17 (2014), Art. 18.
- [40] S. S. Dragomir - P. Cerone - J. Roumeliotis - S. Wang, *A weighted version of Ostrowski inequality for mappings of Hölder type and applications in numerical analysis*, Bull. Math. Soc. Sci. Math. Roumanie 42 (90) (4) (1999), 301–314.
- [41] S. S. Dragomir - S. Fitzpatrick, *The Hadamard inequalities for s -convex functions in the second sense*, Demonstratio Math. 32 (4) (1999), 687–696.
- [42] S. S. Dragomir - S. Fitzpatrick, *The Jensen inequality for s -Breckner convex functions in linear spaces*, Demonstratio Math. 33 (1) (2000), 43–49.
- [43] S. S. Dragomir - B. Mond, *On Hadamard's inequality for a class of functions of Godunova and Levin*, Indian J. Math. 39 (1) (1997), 1–9.
- [44] S. S. Dragomir - C. E. M. Pearce, *On Jensen's inequality for a class of functions of Godunova and Levin*, Period. Math. Hungar. 33 (2) (1996), 93–100.
- [45] S. S. Dragomir - C. E. M. Pearce, *Quasi-convex functions and Hadamard's inequality*, Bull. Austral. Math. Soc. 57 (1998), 377–385.

- [46] S. S. Dragomir - J. Pečarić - L. Persson, *Some inequalities of Hadamard type*, Soochow J. Math. 21 (3) (1995), 335–341.
- [47] S. S. Dragomir - J. Pečarić - L. Persson, *Properties of some functionals related to Jensen's inequality*, Acta Math. Hungarica 70 (1996), 129–143.
- [48] S. S. Dragomir - Th. M. Rassias (Eds), *Ostrowski Type Inequalities and Applications in Numerical Integration*, Kluwer Academic Publisher, 2002.
- [49] S. S. Dragomir - S. Wang, *A new inequality of Ostrowski's type in L_1 -norm and applications to some special means and to some numerical quadrature rules*, Tamkang J. of Math. 28 (1997), 239–244.
- [50] S. S. Dragomir - S. Wang, *Applications of Ostrowski's inequality to the estimation of error bounds for some special means and some numerical quadrature rules*, Appl. Math. Lett. 11 (1998), 105–109.
- [51] S. S. Dragomir - S. Wang, *A new inequality of Ostrowski's type in L_p -norm and applications to some special means and to some numerical quadrature rules*, Indian J. of Math., 40 (3) (1998), 245–304.
- [52] A. El Farissi, *Simple proof and refinement of Hermite-Hadamard inequality*, J. Math. Ineq. 4 (3) (2010), 365–369.
- [53] T. Furuta - J. Mičić Hot - J. Pečarić - Y. Seo, *Mond-Pečarić Method in Operator Inequalities. Inequalities for Bounded Selfadjoint Operators on a Hilbert Space*, Element, Zagreb, 2005.
- [54] E. K. Godunova - V. I. Levin, *Inequalities for functions of a broad class that contains convex, monotone and some other forms of functions*, (Russian) Numerical mathematics and mathematical physics (Russian), 138–142, 166, Moskov. Gos. Ped. Inst., Moscow, 1985
- [55] G. Grüss, *Über das maximum des absoluten Betrages von $\frac{1}{b-a} \int_a^b f(x)g(x)dx - \frac{1}{(b-a)^2} \int_a^b f(x)dx \cdot \int_a^b g(x)dx$* , Math. Z. 39 (1934), 215–226.
- [56] G. Helmborg, *Introduction to Spectral Theory in Hilbert Space*, John Wiley & Sons, Inc. New York, 1969.
- [57] H. Hudzik - L. Maligranda, *Some remarks on s -convex functions*, Aequationes Math. 48 (1) (1994), 100–111.
- [58] E. Kikianty - S. S. Dragomir, *Hermite-Hadamard's inequality and the p -HH-norm on the Cartesian product of two copies of a normed space*, Math. Inequal. Appl. 13 (1) (2010), 1–32.
- [59] U. S. Kirmaci - M. Klaričić Bakula - M. E. Özdemir - J. Pečarić, *Hadamard-type inequalities for s -convex functions*, Appl. Math. Comput. 193 (1) (2007), 26–35.
- [60] M. A. Latif, *On some inequalities for h -convex functions*, Int. J. Math. Anal. (Ruse) 4 (29-32) (2010), 1473–1482.
- [61] A. Lupaş, *The best constant in an integral inequality*, Mathematica (Cluj) 15 (38) (2) (1973), 219–222.

- [62] D. S. Mitrinović - I. B. Lacković, *Hermite and convexity*, Aequationes Math. 28 (1985), 229–232.
- [63] D. S. Mitrinović - J. E. Pečarić, *Note on a class of functions of Godunova and Levin*, C. R. Math. Rep. Acad. Sci. Canada 12 (1) (1990), 33–36.
- [64] A. Matković - J. E. Pečarić - I. Perić, *A variant of Jensen's inequality of Mercer's type for operators with applications*, Linear Algebra Appl. 418 (2-3) (2006), 551–564.
- [65] D. S. Mitrinović - J. E. Pečarić - A. M. Fink, *Inequalities Involving Functions and their Integrals and Derivatives*, Kluwer Academic Publishers, Dordrecht, 1991.
- [66] D. S. Mitrinović - J. E. Pečarić - A. M. Fink, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, Dordrecht, 1993.
- [67] B. Mond - J. E. Pečarić, *Convex inequalities in Hilbert spaces*, Houston J. Math., 19 (1993), 405–420.
- [68] B. Mond - J. E. Pečarić, *Classical inequalities for matrix functions*, Utilitas Math. 46 (1994), 155–166.
- [69] A. M. Ostrowski, *On an integral inequality*, Aequationes Math. 4 (1970), 358–373.
- [70] J. E. Pečarić - J. Mičić - Y. Seo, *Inequalities between operator means based on the Mond-Pečarić method*, Houston J. Math. 30 (1) (2004), 191–207.
- [71] C. E. M. Pearce - A. M. Rubinov, *P-functions, quasi-convex functions, and Hadamard-type inequalities*, J. Math. Anal. Appl. 240 (1) (1999), 92–104.
- [72] J. E. Pečarić - S. S. Dragomir, *On an inequality of Godunova-Levin and some refinements of Jensen integral inequality*, Itinerant Seminar on Functional Equations, Approximation and Convexity (Cluj-Napoca, 1989), 263–268, Preprint, 89-6, Univ. "Babeş-Bolyai", Cluj-Napoca, 1989.
- [73] J. Pečarić - S. S. Dragomir, *A generalization of Hadamard's inequality for isotonic linear functionals*, Radovi Mat. (Sarajevo) 7 (1991), 103–107.
- [74] M. Radulescu - S. Radulescu - P. Alexandrescu, *On the Godunova-Levin-Schur class of functions*, Math. Inequal. Appl. 12 (4) (2009), 853–862.
- [75] M. Z. Sarikaya - A. Saglam - H. Yildirim, *On some Hadamard-type inequalities for h-convex functions*, J. Math. Inequal. 2 (3) (2008), 335–341.
- [76] E. Set - M. E. Özdemir - M. Z. Sarikaya, *New inequalities of Ostrowski's type for s-convex functions in the second sense with applications*, Facta Univ. Ser. Math. Inform. 27 (1) (2012), 67–82.
- [77] M. Z. Sarikaya - E. Set - M. E. Özdemir, *On some new inequalities of Hadamard type involving h-convex functions*, Acta Math. Univ. Comenian. (N.S.) 79 (2) (2010), 265–272.
- [78] M. S. Slater, *A companion inequality to Jensen's inequality*, J. Approx. Theory 32 (1984), 160–166.

- [79] M. Tunç, *Ostrowski-type inequalities via h -convex functions with applications to special means*, J. Inequal. Appl. 2013 (2013):326, 10 pp.
- [80] S. Varošanec, *On h -convexity*, J. Math. Anal. Appl. 326 (1) (2007), 303–311.

SILVESTRU SEVER DRAGOMIR

*Mathematics, College of Engineering & Science
Victoria University, PO Box 14428
Melbourne City, MC 8001, Australia.*

*School of Computational & Applied Mathematics,
University of the Witwatersrand,
Private Bag 3, Johannesburg 2050, South Africa
e-mail: sever.dragomir@vu.edu.au*