

VICTORIA UNIVERSITY
MELBOURNE AUSTRALIA


New Inequalities of Hermite-Hadamard Type for Log-Convex Functions

This is the Published version of the following publication

Dragomir, Sever S (2017) New Inequalities of Hermite-Hadamard Type for Log-Convex Functions. *Khayyam Journal of Mathematics*, 3 (2). 98 - 115.
ISSN 2423-4788

The publisher's official version can be found at
http://www.kjm-math.org/article_47458.html
Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/36633/>



Khayyam Journal of Mathematics

emis.de/journals/KJM
kjm-math.org

NEW INEQUALITIES OF HERMITE-HADAMARD TYPE FOR LOG-CONVEX FUNCTIONS

SILVESTRU SEVER DRAGOMIR^{1,2}

Communicated by J. Brzdęk

ABSTRACT. Some new inequalities of Hermite-Hadamard type for log-convex functions defined on real intervals are given.

1. INTRODUCTION

A function $f : I \rightarrow [0, \infty)$ is said to be *log-convex* or *multiplicatively convex* if $\log f$ is convex, or, equivalently, if for all $x, y \in I$ and $t \in [0, 1]$ one has the inequality:

$$f(tx + (1-t)y) \leq [f(x)]^t [f(y)]^{1-t}. \quad (1.1)$$

We note that if f and g are convex and g is increasing, then $g \circ f$ is convex; moreover, since $f = \exp(\log f)$, it follows that a log-convex function is convex, but the converse may not necessarily be true. This follows directly from (1.1) because, by the *arithmetic-geometric mean inequality*, we have

$$[f(x)]^t [f(y)]^{1-t} \leq tf(x) + (1-t)f(y)$$

for all $x, y \in I$ and $t \in [0, 1]$.

Let us recall the *Hermite-Hadamard inequality*

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2}, \quad (1.2)$$

where $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ is a convex function on the interval I , $a, b \in I$ and $a < b$.

For related results, see [1]-[22], [25]-[28], [29]-[39] and [40]-[51].

Date: Received: 11 April 2017; Revised: 11 June 2017; Accepted: 24 June 2017.

2010 Mathematics Subject Classification. Primary 26D15; Secondary 25D10.

Key words and phrases. Convex functions, integral inequalities, log-convex functions.

Note that if we apply the above inequality for the log-convex functions $f : I \rightarrow (0, \infty)$, we have that

$$\ln \left[f \left(\frac{a+b}{2} \right) \right] \leq \frac{1}{b-a} \int_a^b \ln f(x) dx \leq \frac{\ln f(a) + \ln f(b)}{2}, \quad (1.3)$$

from which we get

$$f \left(\frac{a+b}{2} \right) \leq \exp \left[\frac{1}{b-a} \int_a^b \ln f(x) dx \right] \leq \sqrt{f(a) f(b)}, \quad (1.4)$$

which is an inequality of Hermite-Hadamard's type for log-convex functions.

By using simple properties of log-convex functions, Dragomir and Mond proved in 1998 the following result [31].

Theorem 1.1. *Let $f : I \rightarrow [0, \infty)$ be a log-convex mapping on I and $a, b \in I$ with $a < b$. Then one has the inequality:*

$$f \left(\frac{a+b}{2} \right) \leq \frac{1}{b-a} \int_a^b \sqrt{f(x) f(a+b-x)} dx \leq \sqrt{f(a) f(b)}. \quad (1.5)$$

The inequality between the first and the second term in (1.5) may be improved as follows [31]. A different upper bound for the middle term in (1.5) can be also provided.

Theorem 1.2. *Let $f : I \rightarrow (0, \infty)$ be a log-convex mapping on I and $a, b \in I$ with $a < b$. Then one has the inequalities:*

$$\begin{aligned} f \left(\frac{a+b}{2} \right) &\leq \exp \left[\frac{1}{b-a} \int_a^b \ln f(x) dx \right] \\ &\leq \frac{1}{b-a} \int_a^b \sqrt{f(x) f(a+b-x)} dx \\ &\leq \frac{1}{b-a} \int_a^b f(x) dx \leq L(f(a), f(b)), \end{aligned} \quad (1.6)$$

where $L(p, q)$ is the logarithmic mean of the strictly positive real numbers p, q , i.e.,

$$L(p, q) := \frac{p-q}{\ln p - \ln q} \text{ if } p \neq q \text{ and } L(p, p) := p.$$

The last inequality in (1.6) was obtained in a different context in [41].

As shown in [57], the following result also holds.

Theorem 1.3. *Let $f : I \rightarrow (0, \infty)$ be a log-convex mapping on I and $a, b \in I$ with $a < b$. Then one has the inequalities:*

$$f \left(\frac{a+b}{2} \right) \leq \left(\frac{1}{b-a} \int_a^b \sqrt{f(x)} dx \right)^2 \leq \frac{1}{b-a} \int_a^b f(x) dx. \quad (1.7)$$

The following result improving the classical first Hermite-Hadamard inequality for differentiable log-convex functions also hold [15].

Theorem 1.4. Let $f : I \rightarrow (0, \infty)$ be a differentiable log-convex function on the interval of real numbers $\overset{\circ}{I}$ (the interior of I) and $a, b \in \overset{\circ}{I}$ with $a < b$. Then the following inequalities hold:

$$\begin{aligned} & \frac{\frac{1}{b-a} \int_a^b f(x) dx}{f\left(\frac{a+b}{2}\right)} \\ & \geq L \left(\exp \left[\frac{f'\left(\frac{a+b}{2}\right)}{f\left(\frac{a+b}{2}\right)} \left(\frac{b-a}{2}\right) \right], \exp \left[-\frac{f'\left(\frac{a+b}{2}\right)}{f\left(\frac{a+b}{2}\right)} \left(\frac{b-a}{2}\right) \right] \right) \geq 1. \end{aligned} \quad (1.8)$$

The second Hermite-Hadamard inequality can be improved as follows [15].

Theorem 1.5. Let $f : I \rightarrow \mathbb{R}$ be as in Theorem 1.4. Then we have the inequality:

$$\begin{aligned} & \frac{\frac{f(a)+f(b)}{2}}{\frac{1}{b-a} \int_a^b f(x) dx} \geq 1 + \log \left[\frac{\int_a^b f(x) dx}{\int_a^b f(x) \exp \left[\frac{f'(x)}{f(x)} \left(\frac{a+b}{2} - x\right) \right] dx} \right] \\ & \geq 1 + \log \left[\frac{\frac{1}{b-a} \int_a^b f(x) dx}{f\left(\frac{a+b}{2}\right)} \right] \geq 1. \end{aligned} \quad (1.9)$$

Motivated by the above results, we establish in this paper some new inequalities for log-convex functions, some of them improving earlier results. Applications for special means are also provided.

2. NEW INEQUALITIES

The following refinement of the Hermite-Hadamard inequality holds.

Lemma 2.1. Let $h : [a, b] \rightarrow \mathbb{R}$ be a convex function and $a = x_0 < x_1 < \dots < x_{n-1} < x_n = b$ an arbitrary division of $[a, b]$ with $n \geq 2$. Then

$$\begin{aligned} h\left(\frac{a+b}{2}\right) & \leq \frac{1}{b-a} \sum_{i=0}^{n-1} h\left(\frac{x_i + x_{i+1}}{2}\right) (x_{i+1} - x_i) \\ & \leq \frac{1}{b-a} \int_a^b h(x) dx \\ & \leq \frac{1}{b-a} \sum_{i=0}^{n-1} \frac{h(x_i) + h(x_{i+1})}{2} (x_{i+1} - x_i) \leq \frac{h(a) + h(b)}{2}. \end{aligned} \quad (2.1)$$

The inequality (2.1) was obtained in 1994 as a particular case of a more general result (see [14] and also mentioned in [34, p. 22]). For a direct proof, see the recent paper [27].

Theorem 2.2. Let $f : [a, b] \rightarrow (0, \infty)$ be a log-convex function on $[a, b]$ and $a = x_0 < x_1 < \dots < x_{n-1} < x_n = b$ an arbitrary division of $[a, b]$ with $n \geq 1$.

Then

$$\begin{aligned}
f\left(\frac{a+b}{2}\right) &\leq \prod_{i=1}^{n-1} \left[f\left(\frac{x_i+x_{i+1}}{2}\right) \right]^{\frac{x_{i+1}-x_i}{b-a}} \\
&\leq \exp\left(\frac{1}{b-a} \int_a^b \ln f(x) dx\right) \\
&\leq \prod_{i=1}^{n-1} \left[\sqrt{f(x_i)f(x_{i+1})} \right]^{\frac{x_{i+1}-x_i}{b-a}} \leq \sqrt{f(a)f(b)}.
\end{aligned} \tag{2.2}$$

Proof. If we write the inequality (2.1) for the function $h = \ln f$, then we get

$$\begin{aligned}
\ln f\left(\frac{a+b}{2}\right) &\leq \frac{1}{b-a} \sum_{i=0}^{n-1} (x_{i+1}-x_i) \ln f\left(\frac{x_i+x_{i+1}}{2}\right) \\
&\leq \frac{1}{b-a} \int_a^b \ln f(x) dx \\
&\leq \frac{1}{b-a} \sum_{i=0}^{n-1} \frac{\ln f(x_i) + \ln f(x_{i+1})}{2} (x_{i+1}-x_i) \leq \frac{\ln f(a) + \ln f(b)}{2}.
\end{aligned}$$

This inequality is equivalent to

$$\begin{aligned}
\ln f\left(\frac{a+b}{2}\right) &\leq \ln \left(\prod_{i=1}^{n-1} \left[f\left(\frac{x_i+x_{i+1}}{2}\right) \right]^{\frac{x_{i+1}-x_i}{b-a}} \right) \\
&\leq \frac{1}{b-a} \int_a^b \ln f(x) dx \\
&\leq \ln \left(\prod_{i=1}^{n-1} \left[\sqrt{f(x_i)f(x_{i+1})} \right]^{\frac{x_{i+1}-x_i}{b-a}} \right) \leq \ln \sqrt{f(a)f(b)}.
\end{aligned} \tag{2.3}$$

This inequality is of interest in itself.

If we take the exponential in (2.3), we get the desired result (2.2). \square

Corollary 2.3. *Let $f : [a, b] \rightarrow (0, \infty)$ be a log-convex function on $[a, b]$ and $x \in [a, b]$, then*

$$\begin{aligned}
f\left(\frac{a+b}{2}\right) &\leq \left[f\left(\frac{a+x}{2}\right) \right]^{\frac{x-a}{b-a}} \left[f\left(\frac{x+b}{2}\right) \right]^{\frac{b-x}{b-a}} \\
&\leq \exp\left(\frac{1}{b-a} \int_a^b \ln f(x) dx\right) \\
&\leq \left[\sqrt{f(a)} \right]^{\frac{x-a}{b-a}} \sqrt{f(x)} \left[\sqrt{f(b)} \right]^{\frac{b-x}{b-a}} \leq \sqrt{f(a)f(b)}
\end{aligned} \tag{2.4}$$

and, equivalently

$$\begin{aligned}
\ln f\left(\frac{a+b}{2}\right) &\leq \frac{x-a}{b-a} \ln f\left(\frac{a+x}{2}\right) + \frac{b-x}{b-a} \ln f\left(\frac{x+b}{2}\right) \\
&\leq \frac{1}{b-a} \int_a^b \ln f(x) dx \\
&\leq \frac{1}{2} \left[\ln f(x) + \frac{(x-a) \ln f(a) + (b-x) \ln f(b)}{b-a} \right] \\
&\leq \frac{\ln f(a) + \ln f(b)}{2}.
\end{aligned} \tag{2.5}$$

Remark 2.4. If we take in (2.5) $x = \frac{a+b}{2}$, then we get

$$\begin{aligned}
\ln f\left(\frac{a+b}{2}\right) &\leq \frac{1}{2} \left[\ln f\left(\frac{3a+b}{4}\right) + \ln f\left(\frac{a+3b}{4}\right) \right] \\
&\leq \frac{1}{b-a} \int_a^b \ln f(x) dx \\
&\leq \frac{1}{2} \left[\ln f\left(\frac{a+b}{2}\right) + \frac{\ln f(a) + \ln f(b)}{2} \right] \leq \frac{\ln f(a) + \ln f(b)}{2}.
\end{aligned} \tag{2.6}$$

From the second inequality in (2.6) we get

$$\begin{aligned}
0 &\leq \frac{1}{b-a} \int_a^b \ln f(x) dx - \ln f\left(\frac{a+b}{2}\right) \\
&\leq \frac{\ln f(a) + \ln f(b)}{2} - \frac{1}{b-a} \int_a^b \ln f(x) dx,
\end{aligned}$$

which shows that the integral term in (1.3) is closer to the left side than to the right side of that inequality.

We also have the particular inequalities:

$$\begin{aligned}
&\ln f\left(\frac{a+b}{2}\right) \\
&\leq \frac{1}{\sqrt{b} + \sqrt{a}} \left[\sqrt{a} \ln f\left(\frac{\sqrt{a}(\sqrt{a} + \sqrt{b})}{2}\right) + \sqrt{b} \ln f\left(\frac{\sqrt{b}(\sqrt{a} + \sqrt{b})}{2}\right) \right] \\
&\leq \frac{1}{b-a} \int_a^b \ln f(y) dy \\
&\leq \frac{1}{2} \left[\frac{\sqrt{b} \ln f(b) + \sqrt{a} \ln f(a)}{\sqrt{b} + \sqrt{a}} + \ln f(\sqrt{ab}) \right] \leq \frac{\ln f(a) + \ln f(b)}{2}
\end{aligned} \tag{2.7}$$

and

$$\begin{aligned}
& \ln f\left(\frac{a+b}{2}\right) \\
& \leq \frac{1}{a+b}a \ln f\left(a\frac{3a+b}{2(a+b)}\right) + \frac{1}{a+b}b \ln f\left(b\frac{a+3b}{2(a+b)}\right) \\
& \leq \frac{1}{b-a} \int_a^b \ln f(y) dy \\
& \leq \frac{1}{2} \left[\frac{b \ln f(b) + a \ln f(a)}{a+b} + \ln f\left(\frac{2ab}{a+b}\right) \right] \leq \frac{\ln f(a) + \ln f(b)}{2}.
\end{aligned} \tag{2.8}$$

The following reverses of the Hermite-Hadamard inequality hold [23, 24].

Lemma 2.5. *Let $h : [a, b] \rightarrow \mathbb{R}$ be a convex function on $[a, b]$. Then*

$$\begin{aligned}
0 & \leq \frac{1}{8} \left[h_+\left(\frac{a+b}{2}\right) - h_-\left(\frac{a+b}{2}\right) \right] (b-a) \\
& \leq \frac{h(a) + h(b)}{2} - \frac{1}{b-a} \int_a^b h(x) dx \\
& \leq \frac{1}{8} [h_-(b) - h_+(a)] (b-a)
\end{aligned} \tag{2.9}$$

and

$$\begin{aligned}
0 & \leq \frac{1}{8} \left[h_+\left(\frac{a+b}{2}\right) - h_-\left(\frac{a+b}{2}\right) \right] (b-a) \\
& \leq \frac{1}{b-a} \int_a^b h(x) dx - h\left(\frac{a+b}{2}\right) \\
& \leq \frac{1}{8} [h_-(b) - h_+(a)] (b-a).
\end{aligned} \tag{2.10}$$

The constant $\frac{1}{8}$ is best possible in all inequalities from (2.9) and (2.10).

In the case of log-convex functions we have:

Theorem 2.6. *Let $f : [a, b] \rightarrow (0, \infty)$ be a log-convex function on $[a, b]$. Then*

$$\begin{aligned}
1 & \leq \exp\left(\frac{1}{8} \left[\frac{f_+\left(\frac{a+b}{2}\right) - f_-\left(\frac{a+b}{2}\right)}{f\left(\frac{a+b}{2}\right)} \right] (b-a)\right) \\
& \leq \frac{\sqrt{f(a)f(b)}}{\exp\left(\frac{1}{b-a} \int_a^b \ln f(x) dx\right)} \\
& \leq \exp\left(\frac{1}{8} \left[\frac{f_-(b)}{f(b)} - \frac{f_+(a)}{f(a)} \right] (b-a)\right)
\end{aligned} \tag{2.11}$$

and

$$\begin{aligned}
1 &\leq \exp \left(\frac{1}{8} \left[\frac{f_+ \left(\frac{a+b}{2} \right) - f_- \left(\frac{a+b}{2} \right)}{f \left(\frac{a+b}{2} \right)} \right] (b-a) \right) \\
&\leq \frac{\exp \left(\frac{1}{b-a} \int_a^b \ln f(x) dx \right)}{f \left(\frac{a+b}{2} \right)} \\
&\leq \exp \left(\frac{1}{8} \left[\frac{f_-(b)}{f(b)} - \frac{f_+(a)}{f(a)} \right] (b-a) \right).
\end{aligned} \tag{2.12}$$

Proof. If we write the inequality (2.9) for the convex function $h = \ln f$

$$\begin{aligned}
0 &\leq \frac{1}{8} \left[\frac{f_+ \left(\frac{a+b}{2} \right) - f_- \left(\frac{a+b}{2} \right)}{f \left(\frac{a+b}{2} \right)} \right] (b-a) \\
&\leq \frac{\ln f(a) + \ln f(b)}{2} - \frac{1}{b-a} \int_a^b \ln f(x) dx \\
&\leq \frac{1}{8} \left[\frac{f_-(b)}{f(b)} - \frac{f_+(a)}{f(a)} \right] (b-a)
\end{aligned}$$

that is equivalent to

$$\begin{aligned}
0 &\leq \ln \left[\exp \left(\frac{1}{8} \left[\frac{f_+ \left(\frac{a+b}{2} \right) - f_- \left(\frac{a+b}{2} \right)}{f \left(\frac{a+b}{2} \right)} \right] (b-a) \right) \right] \\
&\leq \ln \left(\frac{\sqrt{f(a)f(b)}}{\exp \left(\frac{1}{b-a} \int_a^b \ln f(x) dx \right)} \right) \\
&\leq \ln \left[\exp \left(\frac{1}{8} \left[\frac{f_-(b)}{f(b)} - \frac{f_+(a)}{f(a)} \right] (b-a) \right) \right].
\end{aligned}$$

By taking the exponential in this inequality, we get the desired result (2.11).

The inequality (2.12) follows from (2.10). \square

We also have the following result.

Theorem 2.7. *Let $f : [a, b] \rightarrow (0, \infty)$ be a log-convex function on $[a, b]$ and $a = x_0 < x_1 < \dots < x_{n-1} < x_n = b$ be an arbitrary division of $[a, b]$ with $n \geq 1$. Then*

$$\begin{aligned}
\exp \left[\frac{1}{b-a} \int_a^b \ln f(x) dx \right] &\leq \frac{1}{b-a} \sum_{i=0}^{n-1} \int_{x_i}^{x_{i+1}} \sqrt{f(x) f(x_i + x_{i+1} - x)} dx \\
&\leq \frac{1}{b-a} \int_a^b f(x) dx.
\end{aligned} \tag{2.13}$$

Proof. Observe that we have

$$\begin{aligned} & \exp \left[\frac{1}{b-a} \int_a^b \ln f(x) dx \right] \\ &= \exp \left[\frac{1}{b-a} \sum_{i=0}^{n-1} \int_{x_i}^{x_{i+1}} \ln f(x) dx \right] \\ &= \exp \left[\sum_{i=0}^{n-1} \frac{x_{i+1} - x_i}{b-a} \left(\frac{1}{x_{i+1} - x_i} \int_{x_i}^{x_{i+1}} \ln f(x) dx \right) \right]. \end{aligned} \tag{2.14}$$

Since $\sum_{i=0}^{n-1} \frac{x_{i+1} - x_i}{b-a} = 1$, by Jensen's inequality for the convex function \exp we have

$$\begin{aligned} & \exp \left[\sum_{i=0}^{n-1} \frac{x_{i+1} - x_i}{b-a} \left(\frac{1}{x_{i+1} - x_i} \int_{x_i}^{x_{i+1}} \ln f(x) dx \right) \right] \\ & \leq \sum_{i=0}^{n-1} \frac{x_{i+1} - x_i}{b-a} \exp \left(\frac{1}{x_{i+1} - x_i} \int_{x_i}^{x_{i+1}} \ln f(x) dx \right). \end{aligned} \tag{2.15}$$

Utilising the inequality (1.6) on each of the intervals $[x_i, x_{i+1}]$ for $i \in \{0, \dots, n-1\}$ we have

$$\begin{aligned} & \exp \left[\frac{1}{x_{i+1} - x_i} \int_{x_i}^{x_{i+1}} \ln f(x) dx \right] \\ & \leq \frac{1}{x_{i+1} - x_i} \int_{x_i}^{x_{i+1}} \sqrt{f(x) f(x_i + x_{i+1} - x)} dx \\ & \leq \frac{1}{x_{i+1} - x_i} \int_{x_i}^{x_{i+1}} f(x) dx, \end{aligned} \tag{2.16}$$

for any $i \in \{0, \dots, n-1\}$.

If we multiply the inequality (2.16) by $\frac{x_{i+1} - x_i}{b-a}$ and sum over i from 0 to $n-1$, then we get

$$\begin{aligned} & \sum_{i=0}^{n-1} \frac{x_{i+1} - x_i}{b-a} \exp \left(\frac{1}{x_{i+1} - x_i} \int_{x_i}^{x_{i+1}} \ln f(x) dx \right) \\ & \leq \frac{1}{b-a} \sum_{i=0}^{n-1} \int_{x_i}^{x_{i+1}} \sqrt{f(x) f(x_i + x_{i+1} - x)} dx \leq \frac{1}{b-a} \sum_{i=0}^{n-1} \int_{x_i}^{x_{i+1}} f(x) dx \\ & = \frac{1}{b-a} \int_a^b f(x) dx. \end{aligned} \tag{2.17}$$

Making use of (2.14), (2.15) and (2.17) we get the desired result (2.13). □

Corollary 2.8. *Let $f : [a, b] \rightarrow (0, \infty)$ be a log-convex function on $[a, b]$ and $y \in [a, b]$, then*

$$\begin{aligned} & \exp \left[\frac{1}{b-a} \int_a^b \ln f(x) dx \right] \\ & \leq \frac{1}{b-a} \left[\int_a^y \sqrt{f(x) f(a+y-x)} dx + \int_y^b \sqrt{f(x) f(b+y-x)} dx \right] \\ & \leq \frac{1}{b-a} \int_a^b f(x) dx. \end{aligned} \quad (2.18)$$

We define the p -logarithmic mean as

$$L_p(a, b) := \begin{cases} \left[\frac{b^{p+1} - a^{p+1}}{(p+1)(b-a)} \right]^{\frac{1}{p}}, & \text{with } a \neq b \\ a, & \text{if } a = b \end{cases}$$

for $p \neq 0, -1$ and $a, b > 0$.

The following result also holds.

Theorem 2.9. *Let $f : [a, b] \rightarrow (0, \infty)$ be a log-convex function on $[a, b]$. Then for any $p > 0$ we have the inequality*

$$\begin{aligned} f\left(\frac{a+b}{2}\right) & \leq \exp \left[\frac{1}{b-a} \int_a^b \ln f(x) dx \right] \\ & \leq \left(\frac{1}{b-a} \int_a^b f^p(x) f^p(a+b-x) dx \right)^{\frac{1}{2p}} \\ & \leq \left(\frac{1}{b-a} \int_a^b f^{2p}(x) dx \right)^{\frac{1}{2p}} \\ & \leq \begin{cases} [L_{2p-1}(f(a), f(b))]^{1-\frac{1}{2p}} [L(f(a), f(b))]^{\frac{1}{2p}}, & p \neq \frac{1}{2}; \\ L(f(a), f(b)), & p = \frac{1}{2}. \end{cases} \end{aligned} \quad (2.19)$$

If $p \in (0, \frac{1}{2})$, then we have

$$\begin{aligned} f\left(\frac{a+b}{2}\right) & \leq \exp \left[\frac{1}{b-a} \int_a^b \ln f(x) dx \right] \\ & \leq \left(\frac{1}{b-a} \int_a^b f^p(x) f^p(a+b-x) dx \right)^{\frac{1}{2p}} \\ & \leq \left(\frac{1}{b-a} \int_a^b f^{2p}(x) dx \right)^{\frac{1}{2p}} \leq \frac{1}{b-a} \int_a^b f(x) dx. \end{aligned} \quad (2.20)$$

Proof. If f is a log-convex function on $[a, b]$, then f^{2p} is log-convex on $[a, b]$ for $p > 0$ and by (1.6) we have

$$\begin{aligned} f^{2p}\left(\frac{a+b}{2}\right) &\leq \exp\left[\frac{1}{b-a}\int_a^b \ln f^{2p}(x) dx\right] \\ &\leq \frac{1}{b-a}\int_a^b f^p(x) f^p(a+b-x) dx \\ &\leq \frac{1}{b-a}\int_a^b f^{2p}(x) dx \leq L(f^{2p}(a), f^{2p}(b)). \end{aligned} \quad (2.21)$$

Taking the power $\frac{1}{2p}$ in (2.21) we get

$$\begin{aligned} f\left(\frac{a+b}{2}\right) &\leq \exp\left[\frac{1}{b-a}\int_a^b \ln f(x) dx\right] \\ &\leq \left(\frac{1}{b-a}\int_a^b f^p(x) f^p(a+b-x) dx\right)^{\frac{1}{2p}} \\ &\leq \left(\frac{1}{b-a}\int_a^b f^{2p}(x) dx\right)^{\frac{1}{2p}} \leq [L(f^{2p}(a), f^{2p}(b))]^{\frac{1}{2p}}. \end{aligned} \quad (2.22)$$

Observe that, for $p \neq \frac{1}{2}$,

$$\begin{aligned} [L(f^{2p}(a), f^{2p}(b))]^{\frac{1}{2p}} &= \left[\frac{f^{2p}(a) - f^{2p}(b)}{\ln f^{2p}(a) - \ln f^{2p}(b)}\right]^{\frac{1}{2p}} \\ &= \left[\frac{f^{2p}(a) - f^{2p}(b)}{2p(f(a) - f(b))} \frac{f(a) - f(b)}{\ln f(a) - \ln f(b)}\right]^{\frac{1}{2p}} \\ &= \left[\frac{f^{2p}(a) - f^{2p}(b)}{2p(f(a) - f(b))}\right]^{\frac{1}{2p}} \left[\frac{f(a) - f(b)}{\ln f(a) - \ln f(b)}\right]^{\frac{1}{2p}} \\ &= [L_{2p-1}(f(a), f(b))]^{1-\frac{1}{2p}} [L(f(a), f(b))]^{\frac{1}{2p}} \end{aligned}$$

and by (2.22) we get the desired result (2.19).

The last inequality in (2.20) follows by the following integral inequality for power $q \in (0, 1)$, namely

$$\frac{1}{b-a}\int_a^b f^q(x) dx \leq \left(\frac{1}{b-a}\int_a^b f(x) dx\right)^q,$$

that follows by Jensen's inequality for concave functions. \square

Remark 2.10. If we take in (2.19) $p = 1$, then we get

$$\begin{aligned} f\left(\frac{a+b}{2}\right) &\leq \exp\left[\frac{1}{b-a}\int_a^b \ln f(x) dx\right] \\ &\leq \left(\frac{1}{b-a}\int_a^b f(x)f(a+b-x) dx\right)^{\frac{1}{2}} \\ &\leq \left(\frac{1}{b-a}\int_a^b f^2(x) dx\right)^{\frac{1}{2}} \\ &\leq [A(f(a), f(b))]^{\frac{1}{2}} [L(f(a), f(b))]^{\frac{1}{2}}. \end{aligned} \quad (2.23)$$

If we take $p = \frac{1}{4}$ in (2.20), then we get

$$\begin{aligned} f\left(\frac{a+b}{2}\right) &\leq \exp\left[\frac{1}{b-a}\int_a^b \ln f(x) dx\right] \\ &\leq \left(\frac{1}{b-a}\int_a^b \sqrt[4]{f(x)f(a+b-x)} dx\right)^2 \\ &\leq \left(\frac{1}{b-a}\int_a^b \sqrt{f(x)} dx\right)^2 \leq \frac{1}{b-a}\int_a^b f(x) dx. \end{aligned} \quad (2.24)$$

This improves the inequality (1.7).

3. RELATED INEQUALITIES

In this section, we establish some related results for log-convex functions.

Theorem 3.1. *Let $f : [a, b] \rightarrow (0, \infty)$ be a log-convex function on $[a, b]$. Then for any $x \in [a, b]$ we have*

$$\begin{aligned} f(b)(b-x) + f(a)(x-a) - \int_a^b f(y) dy \\ \geq \int_a^b f(y) \ln f(y) dy - \ln f(x) \int_a^b f(y) dy. \end{aligned} \quad (3.1)$$

In particular,

$$\begin{aligned} \frac{f(b) + f(a)}{2} - \frac{1}{b-a} \int_a^b f(y) dy \\ \geq \frac{1}{b-a} \int_a^b f(y) \ln f(y) dy - \ln f\left(\frac{a+b}{2}\right) \frac{1}{b-a} \int_a^b f(y) dy, \end{aligned} \quad (3.2)$$

$$\begin{aligned} \frac{f(b)\sqrt{b} + f(a)\sqrt{a}}{\sqrt{b} + \sqrt{a}} - \frac{1}{b-a} \int_a^b f(y) dy \\ \geq \frac{1}{b-a} \int_a^b f(y) \ln f(y) dy - \ln f(\sqrt{ab}) \frac{1}{b-a} \int_a^b f(y) dy \end{aligned} \quad (3.3)$$

and

$$\begin{aligned} & \frac{f(b)b + f(a)a}{a+b} - \frac{1}{b-a} \int_a^b f(y) dy \\ & \geq \frac{1}{b-a} \int_a^b f(y) \ln f(y) dy - \ln f\left(\frac{2ab}{a+b}\right) \frac{1}{b-a} \int_a^b f(y) dy. \end{aligned} \quad (3.4)$$

Proof. Since the function $\ln f$ is convex on $[a, b]$, by the gradient inequality we have

$$\ln f(x) - \ln f(y) \geq \frac{f'_+(y)}{f(y)} (x - y), \quad (3.5)$$

for any $x \in [a, b]$ and $y \in (a, b)$.

If we multiply (3.5) by $f(y) > 0$ and integrate on $[a, b]$ over y we get

$$\begin{aligned} & \ln f(x) \int_a^b f(y) dy - \int_a^b f(y) \ln f(y) dy \\ & \geq \int_a^b f'_+(y) (x - y) dy = f(y) (x - y)|_a^b + \int_a^b f(y) dy \\ & = f(b) (x - b) + f(a) (a - x) + \int_a^b f(y) dy, \end{aligned}$$

which is equivalent to (3.1).

The inequality (3.2) follows by (3.1) on taking $x = \frac{a+b}{2}$.

If we take in (3.1) $x = \sqrt{ab}$, then we get

$$\begin{aligned} & f(b) \sqrt{b} (\sqrt{b} - \sqrt{a}) + f(a) \sqrt{a} (\sqrt{b} - \sqrt{a}) - \int_a^b f(y) dy \\ & \geq \int_a^b f(y) \ln f(y) dy - \ln f(\sqrt{ab}) \int_a^b f(y) dy, \end{aligned}$$

which is equivalent to (3.3).

If we take in (3.1) $x = \frac{2ab}{a+b}$, then we get

$$\begin{aligned} & f(b) b \left(\frac{b-a}{a+b}\right) + f(a) a \left(\frac{b-a}{a+b}\right) - \int_a^b f(y) dy \\ & \geq \int_a^b f(y) \ln f(y) dy - \ln f\left(\frac{2ab}{a+b}\right) \int_a^b f(y) dy, \end{aligned}$$

which is equivalent to (3.4). □

Corollary 3.2. *Let $f : [a, b] \rightarrow (0, \infty)$ be a log-convex function on $[a, b]$. Then*

$$\begin{aligned} & \frac{f(b) + f(a)}{2} - \frac{1}{b-a} \int_a^b f(y) dy \\ & \geq \int_a^b f(y) \ln f(y) dy - \int_a^b f(y) dy \frac{1}{b-a} \int_a^b \ln f(y) dy \geq 0. \end{aligned} \quad (3.6)$$

Proof. If we take the integral mean over x in (3.1), then we get

$$\begin{aligned} & \frac{1}{b-a} \int_a^b [f(b)(b-x) + f(a)(x-a)] dx - \int_a^b f(y) dy \\ & \geq \int_a^b f(y) \ln f(y) dy - \int_a^b f(y) dy \frac{1}{b-a} \int_a^b \ln f(x) \end{aligned}$$

and since

$$\frac{1}{b-a} \int_a^b [f(b)(b-x) + f(a)(x-a)] dx = \frac{f(b) + f(a)}{2} - \frac{1}{b-a} \int_a^b f(y) dy,$$

then the first inequality in (3.6) is proved.

Since \ln is an increasing function on $(0, \infty)$, we have

$$(f(x) - f(y)) (\ln f(x) - \ln f(y)) \geq 0,$$

for any $x, y \in [a, b]$, showing that the functions f and $\ln f$ are synchronous on $[a, b]$.

By making use of the Čebyšev integral inequality for synchronous functions $g, h : [a, b] \rightarrow \mathbb{R}$, namely

$$\frac{1}{b-a} \int_a^b g(x) h(x) dx \geq \frac{1}{b-a} \int_a^b g(x) dx \frac{1}{b-a} \int_a^b h(x) dx,$$

we have

$$\frac{1}{b-a} \int_a^b f(x) \ln f(x) dx \geq \frac{1}{b-a} \int_a^b f(x) dx \frac{1}{b-a} \int_a^b \ln f(x) dx,$$

which proves the last part of (3.6). \square

The inequality (3.6) improves the well know result for convex functions

$$\frac{f(b) + f(a)}{2} \geq \frac{1}{b-a} \int_a^b f(y) dy.$$

We have:

Corollary 3.3. *Let $f : [a, b] \rightarrow (0, \infty)$ be a log-convex function on $[a, b]$. If $f(a) \neq f(b)$ and*

$$\alpha_f := \frac{\int_a^b f'(y) y dy}{\int_a^b f'(y) dy} = \frac{bf(b) - af(a) - \int_a^b f(y) dy}{f(b) - f(a)} \in [a, b], \quad (3.7)$$

then

$$\ln f(\alpha_f) \geq \frac{\int_a^b f(y) \ln f(y) dy}{\int_a^b f(y) dy}. \quad (3.8)$$

Proof. Follows from (3.1) by observing that

$$f(b)(b - \alpha_f) + f(a)(\alpha_f - a) = \int_a^b f(y) dy.$$

\square

Remark 3.4. We observe that if $f : [a, b] \rightarrow (0, \infty)$ is nondecreasing with $f(a) \neq f(b)$, the condition (3.7) is satisfied.

We also have:

Corollary 3.5. *Let $f : [a, b] \rightarrow (0, \infty)$ be a log-convex function on $[a, b]$. Then*

$$\begin{aligned} f(b) \left(b - \frac{\int_a^b y f(y) dy}{\int_a^b f(y) dy} \right) + f(a) \left(\frac{\int_a^b y f(y) dy}{\int_a^b f(y) dy} - a \right) - \int_a^b f(y) dy \\ \geq \int_a^b f(y) \ln f(y) dy - \int_a^b f(y) dy \ln f \left(\frac{\int_a^b y f(y) dy}{\int_a^b f(y) dy} \right) \geq 0. \end{aligned} \quad (3.9)$$

Proof. The first inequality follows by (3.1) on taking

$$x = \frac{\int_a^b y f(y) dy}{\int_a^b f(y) dy} \in [a, b]$$

since $f(y) > 0$ for any $y \in [a, b]$.

By Jensen's inequality for the convex function $\ln f$ and the positive weight f we have

$$\frac{\int_a^b f(y) \ln f(y) dy}{\int_a^b f(y) dy} \geq f \left(\frac{\int_a^b f(y) y dy}{\int_a^b f(y) dy} \right),$$

which proves the second inequality in (3.9). \square

4. APPLICATIONS

The function $f : (0, \infty) \rightarrow (0, \infty)$, $f(t) = \frac{1}{t}$ is log-convex on $(0, \infty)$. If we use the inequality (2.2) for this function, then we have

$$\begin{aligned} A(a, b) &\geq \prod_{i=1}^{n-1} [A(x_i, x_{i+1})]^{\frac{x_{i+1}-x_i}{b-a}} \geq I(a, b) \\ &\geq \prod_{i=1}^{n-1} [G(x_i, x_{i+1})]^{\frac{x_{i+1}-x_i}{b-a}} \geq G(a, b), \end{aligned} \quad (4.1)$$

for any $a = x_0 < x_1 < \dots < x_{n-1} < x_n = b$ an arbitrary division of $[a, b]$ with $n \geq 1$.

In particular, we have

$$\begin{aligned} A(a, b) &\geq [A(a, x)]^{\frac{x-a}{b-a}} [A(x, b)]^{\frac{b-x}{b-a}} \\ &\geq I(a, b) \geq \sqrt{a^{\frac{x-a}{b-a}} x b^{\frac{b-x}{b-a}}} \geq G(a, b), \end{aligned} \quad (4.2)$$

for any $x \in [a, b]$.

If we use the inequalities (2.11) and (2.12) for $f : (0, \infty) \rightarrow (0, \infty)$, $f(t) = \frac{1}{t}$, then we have

$$(1 \leq) \frac{I(a, b)}{G(a, b)} \leq \exp \left(\frac{1}{8} \frac{(b-a)^2}{ab} \right) \quad (4.3)$$

and

$$(1 \leq) \frac{A(a, b)}{I(a, b)} \leq \exp\left(\frac{1}{8} \frac{(b-a)^2}{ab}\right). \quad (4.4)$$

If we use the inequality (3.6) for $f : (0, \infty) \rightarrow (0, \infty)$, $f(t) = \frac{1}{t}$, then we have

$$L(a, b) - H(a, b) \geq (b-a) H(a, b) \ln\left(\frac{I(a, b)}{G(a, b)}\right) (\geq 0). \quad (4.5)$$

The interested reader may apply the above inequalities for other log-convex functions such as $f(t) = \frac{1}{t^p}$, $p > 0$, $t > 0$, $f(t) = \exp g(t)$, with g any convex function on an interval, etc. The details are omitted.

REFERENCES

1. M. Alomari and M. Darus, *The Hadamard's inequality for s -convex function*. Int. J. Math. Anal. (Ruse), **2** (2008), no. 13-16, 639–646.
2. M. Alomari and M. Darus, *Hadamard-type inequalities for s -convex functions*. Int. Math. Forum, **3** (2008), no. 37-40, 1965–1975.
3. G.A. Anastassiou, *Univariate Ostrowski inequalities, revisited*. Monatsh. Math., **135** (2002), no. 3, 175–189.
4. G.D. Anderson, M.K. Vamanamurthy and M. Vuorinen, *Generalized convexity and inequalities*, J. Math. Anal. Appl., **335** (2007) 1294–1308.
5. N.S. Barnett, P. Cerone, S.S. Dragomir, M.R. Pinheiro, and A. Sofo, *Ostrowski type inequalities for functions whose modulus of the derivatives are convex and applications*. Inequality Theory and Applications, **Vol. 2** (Chinju/Masan, 2001), 19–32, Nova Sci. Publ., Hauppauge, NY, 2003. Preprint: *RGMIA Res. Rep. Coll.* **5** (2002), No. 2, Art. 1 [Online <http://rgmia.org/papers/v5n2/Paperwapp2q.pdf>].
6. E.F. Beckenbach, *Convex functions*, Bull. Amer. Math. Soc., **54**(1948), 439–460.
7. M. Bombardelli and S. Varošanec, *Properties of h -convex functions related to the Hermite-Hadamard-Fejér inequalities*. Comput. Math. Appl., **58** (2009), no. 9, 1869–1877.
8. W.W. Breckner, *Stetigkeitsaussagen für eine Klasse verallgemeinerter konvexer Funktionen in topologischen linearen Räumen*. (German) Publ. Inst. Math. (Beograd) (N.S.), **23(37)** (1978), 13–20.
9. W.W. Breckner and G. Orbán, *Continuity properties of rationally s -convex mappings with values in an ordered topological linear space*. Universitatea "Babeş-Bolyai", Facultatea de Matematica, Cluj-Napoca, 1978. viii+92 pp.
10. P. Cerone and S.S. Dragomir, *Midpoint-type rules from an inequalities point of view*, Ed. G. A. Anastassiou, Handbook of Analytic-Computational Methods in Applied Mathematics, CRC Press, New York. 135-200.
11. P. Cerone and S.S. Dragomir, *New bounds for the three-point rule involving the Riemann-Stieltjes integrals*, in Advances in Statistics Combinatorics and Related Areas, C. Gulati, et al. (Eds.), World Science Publishing, 2002, 53-62.
12. P. Cerone, S.S. Dragomir and J. Roumeliotis, *Some Ostrowski type inequalities for n -time differentiable mappings and applications*, Demonstratio Mathematica, **32** (1999), no. 2, 697–712.
13. G. Cristescu, *Hadamard type inequalities for convolution of h -convex functions*. Ann. Tiberiu Popoviciu Semin. Funct. Equ. Approx. Convexity, **8** (2010), 3–11.
14. S.S. Dragomir, *Some remarks on Hadamard's inequalities for convex functions*, Extracta Math., **9** (2) (1994), 88-94.
15. S.S. Dragomir, *Refinements of the Hermite-Hadamard integral inequality for log-convex functions*, Austral. Math. Soc. Gaz., **28** (2001), no. 3, 129–134.

16. S.S. Dragomir, *Ostrowski's inequality for monotonous mappings and applications*, J. KSIAM, **3** (1999), no. 1, 127-135.
17. S.S. Dragomir, *The Ostrowski's integral inequality for Lipschitzian mappings and applications*, Comp. Math. Appl., **38** (1999), 33-37.
18. S.S. Dragomir, *On the Ostrowski's inequality for Riemann-Stieltjes integral*, Korean J. Appl. Math., **7** (2000), 477-485.
19. S.S. Dragomir, *On the Ostrowski's inequality for mappings of bounded variation and applications*, Math. Ineq. Appl., **4**(1) (2001), 33-40.
20. S.S. Dragomir, *On the Ostrowski inequality for Riemann-Stieltjes integral $\int_a^b f(t) du(t)$ where f is of Hölder type and u is of bounded variation and applications*, J. KSIAM, **5**(1) (2001), no. 1, 35-45.
21. S.S. Dragomir, *Ostrowski type inequalities for isotonic linear functionals*, J. Inequal. Pure Appl. Math., **3** (2002), no. 5, Art. 68.
22. S.S. Dragomir, *An inequality improving the first Hermite-Hadamard inequality for convex functions defined on linear spaces and applications for semi-inner products*. J. Inequal. Pure Appl. Math., **3** (2002), no. 2, Article 31, 8 pp.
23. S.S. Dragomir, *An inequality improving the first Hermite-Hadamard inequality for convex functions defined on linear spaces and applications for semi-inner products*, J. Inequal. Pure and Appl. Math., **3** (2002), no. 2, Art. 31.
24. S.S. Dragomir, *An inequality improving the second Hermite-Hadamard inequality for convex functions defined on linear spaces and applications for semi-inner products*, J. Inequal. Pure and Appl. Math., **3** (2002), no. 3, Art. 35.
25. S.S. Dragomir, *An Ostrowski like inequality for convex functions and applications*, Revista Math. Complutense, **16** (2003), no. 2, 373-382.
26. S.S. Dragomir, *Operator Inequalities of Ostrowski and Trapezoidal Type*. Springer Briefs in Mathematics. Springer, New York, 2012. x+112 pp. ISBN: 978-1-4614-1778-1
27. S.S. Dragomir, *Some new inequalities of Hermite-Hadamard type for GA-convex functions*, Preprint RGMIA Res. Rep. Coll., **18** (2015), Art 33. [<http://rgmia.org/papers/v18/v18a33.pdf>].
28. S.S. Dragomir, P. Cerone, J. Roumeliotis and S. Wang, *A weighted version of Ostrowski inequality for mappings of Hölder type and applications in numerical analysis*, Bull. Math. Soc. Sci. Math. Roumanie, **42(90)** (1999), 4, 301-314.
29. S.S. Dragomir and S. Fitzpatrick, *The Hadamard inequalities for s -convex functions in the second sense*. Demonstratio Math. **32** (1999), no. 4, 687-696.
30. S.S. Dragomir and S. Fitzpatrick, *The Jensen inequality for s -Breckner convex functions in linear spaces*. Demonstratio Math., **33** (2000), no. 1, 43-49.
31. S.S. Dragomir and B. Mond, *Integral inequalities of Hadamard's type for log-convex functions*, Demonstratio Math., **31** (1998), no. 2, 354-364.
32. S.S. Dragomir and C.E.M. Pearce, *On Jensen's inequality for a class of functions of Godunova and Levin*. Period. Math. Hungar., **33** (1996), no. 2, 93-100.
33. S.S. Dragomir and C.E.M. Pearce, *Quasi-convex functions and Hadamard's inequality*, Bull. Austral. Math. Soc., **57** (1998), 377-385.
34. S.S. Dragomir and C.E.M. Pearce, *Selected Topics on Hermite-Hadamard Inequalities and Applications*, RGMIA Monographs, Victoria University, 2000.
35. S.S. Dragomir, J. Pečarić and L. Persson, *Some inequalities of Hadamard type*. Soochow J. Math. **21** (1995), no. 3, 335-341.
36. S.S. Dragomir and Th. M. Rassias (Eds), *Ostrowski Type Inequalities and Applications in Numerical Integration*, Kluwer Academic Publisher, 2002.
37. S.S. Dragomir and S. Wang, *A new inequality of Ostrowski's type in L_1 -norm and applications to some special means and to some numerical quadrature rules*, Tamkang J. of Math., **28**(1997), 239-244.

38. S.S. Dragomir and S. Wang, *Applications of Ostrowski's inequality to the estimation of error bounds for some special means and some numerical quadrature rules*, Appl. Math. Lett., **11** (1998), 105-109.
39. S.S. Dragomir and S. Wang, *A new inequality of Ostrowski's type in L_p -norm and applications to some special means and to some numerical quadrature rules*, Indian J. of Math., **40** (1998), no. 3, 245-304.
40. A. El Farissi, *Simple proof and refinement of Hermite-Hadamard inequality*, J. Math. Ineq., **4** (2010), no. 3, 365-369.
41. P.M. Gill, C.E.M. Pearce and J. Pečarić, *Hadamard's inequality for r -convex functions*, Journal of Mathematical Analysis and Applications., **215** (1997), no. 2, 461-470.
42. E.K. Godunova and V.I. Levin, *Inequalities for functions of a broad class that contains convex, monotone and some other forms of functions*. (Russian) Numerical mathematics and mathematical physics (Russian), 138-142, 166, Moskov. Gos. Ped. Inst., Moscow, 1985
43. H. Hudzik and L. Maligranda, *Some remarks on s -convex functions*. Aequationes Math., **48** (1994), no. 1, 100-111.
44. E. Kikianty and S.S. Dragomir, *Hermite-Hadamard's inequality and the p -HH-norm on the Cartesian product of two copies of a normed space*, Math. Inequal. Appl., **13** (2010), no. 1, 1-32.
45. U.S. Kirmaci, M. Klaričić Bakula, M.E. Özdemir and J. Pečarić, *Hadamard-type inequalities for s -convex functions*. Appl. Math. Comput., **193** (2007), no. 1, 26-35.
46. M.A. Latif, *On some inequalities for h -convex functions*. Int. J. Math. Anal. (Ruse), **4** (2010), no. 29-32, 1473-1482.
47. D.S. Mitrinović and I. B. Lacković, *Hermite and convexity*, Aequationes Math. **28** (1985), 229-232.
48. D.S. Mitrinović and J. E. Pečarić, *Note on a class of functions of Godunova and Levin*. C. R. Math. Rep. Acad. Sci. Canada, **12** (1990), no. 1, 33-36.
49. M.A. Noor, K.I. Noor and M.U. Awan, *Some inequalities for geometrically-arithmetically h -convex functions*, Creat. Math. Inform. **23** (2014), No. 1, 91 - 98.
50. C.E.M. Pearce and A.M. Rubinov, *P -functions, quasi-convex functions, and Hadamard-type inequalities*. J. Math. Anal. Appl. **240** (1999), no. 1, 92-104.
51. J.E. Pečarić and S.S. Dragomir, *On an inequality of Godunova-Levin and some refinements of Jensen integral inequality*. Itinerant Seminar on Functional Equations, Approximation and Convexity (Cluj-Napoca, 1989), 263-268, Preprint, 89-6, Univ. "Babeş-Bolyai", Cluj-Napoca, 1989.
52. J. Pečarić and S.S. Dragomir, *A generalization of Hadamard's inequality for isotonic linear functionals*, Radovi Mat. (Sarajevo), **7** (1991), 103-107.
53. M. Radulescu, S. Radulescu and P. Alexandrescu, *On the Godunova-Levin-Schur class of functions*. Math. Inequal. Appl., **12** (2009), no. 4, 853-862.
54. M.Z. Sarikaya, A. Saglam, and H. Yildirim, *On some Hadamard-type inequalities for h -convex functions*. J. Math. Inequal., **2** (2008), no. 3, 335-341.
55. E. Set, M.E. Özdemir and M.Z. Sarikaya, *New inequalities of Ostrowski's type for s -convex functions in the second sense with applications*. Facta Univ. Ser. Math. Inform., **27** (2012), no. 1, 67-82.
56. M.Z. Sarikaya, E. Set and M.E. Özdemir, *On some new inequalities of Hadamard type involving h -convex functions*. Acta Math. Univ. Comenian. (N.S.), **79** (2010), no. 2, 265-272.
57. W.T. Sulaiman, *Refinements to Hadamard's inequality for log-convex functions*. Applied Mathematics, **2011**, 2, 899-903.
58. M. Tunç, *Ostrowski-type inequalities via h -convex functions with applications to special means*. J. Inequal. Appl., **2013**, 2013:326.
59. S. Varošanec, *On h -convexity*. J. Math. Anal. Appl., **326** (2007), no. 1, 303-311.

60. X.-M. Zhang, Y.-M. Chu and X.-H. Zhang, *The Hermite-Hadamard type inequality of GA-convex functions and its application*, Journal of Inequalities and Applications, **2010**, Article ID 507560, 11 pages.

¹MATHEMATICS, COLLEGE OF ENGINEERING & SCIENCE, VICTORIA UNIVERSITY, PO BOX 14428, MELBOURNE CITY, MC 8001, AUSTRALIA.

E-mail address: sever.dragomir@vu.edu.au

²DST-NRF CENTRE OF EXCELLENCE IN THE MATHEMATICAL AND STATISTICAL SCIENCES, SCHOOL OF COMPUTATIONAL & APPLIED MATHEMATICS, UNIVERSITY OF THE WITWATERSRAND, PRIVATE BAG 3, JOHANNESBURG 2050, SOUTH AFRICA.