



VICTORIA UNIVERSITY
MELBOURNE AUSTRALIA

Addressing RFID misreadings to better infer bee hive activity

This is the Published version of the following publication

Susanto, Ferry, Gillard, T, De Souza, P, Vincent, B, Budi, S, Almeida, A, Pessin, G, Arruda, H, Williams, RN, Engelke, U, Marendy, P, Hirsch, P and He, Jing (2018) Addressing RFID misreadings to better infer bee hive activity. IEEE Access, 6. 31935 - 31949. ISSN 2169-3536

The publisher's official version can be found at
<https://ieeexplore.ieee.org/document/8375089>

Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/37782/>

Received April 26, 2018, accepted May 27, 2018, date of publication June 7, 2018, date of current version June 29, 2018.

Digital Object Identifier 10.1109/ACCESS.2018.2844181

Addressing RFID Misreadings to Better Infer Bee Hive Activity

FERRY SUSANTO^{1,2,3}, THOMAS GILLARD^{1,4}, PAULO DE SOUZA, JR.¹, BENITA VINCENT¹,
SETIA BUDI^{1,5,6}, AURO ALMEIDA⁷, GUSTAVO PESSIN⁸, HELDER ARRUDA⁸,
RAYMOND N. WILLIAMS^{1,5}, ULRICH ENGELKE¹, PETER MARENDY^{1,5},
PASCAL HIRSCH^{1,5}, AND JING HE^{2,3}

¹Data61, Commonwealth Scientific and Industrial Research Organization, Sandy Bay, TAS 7005, Australia

²Institute of Information Technology, Nanjing University of Finance and Economics, Nanjing 210003, China

³School of Software and Electrical Engineering, Swinburne University of Technology, Melbourne, VIC 3122, Australia

⁴School of Life and Environmental Sciences, The University of Sydney, Sydney, NSW 2006, Australia

⁵School of Engineering and ICT, University of Tasmania, Sandy Bay, TAS 7005, Australia

⁶Faculty of Information Technology, Maranatha Christian University, Bandung 40164, Indonesia

⁷Land and Water, Commonwealth Scientific and Industrial Research Organization, University of Tasmania, Hobart, TAS 7005, Australia

⁸Sustainable Development, Vale Institute of Technology, Belem 955, Brazil

Corresponding author: Jing He (lotusjing@gmail.com)

This work was supported in part by the National Natural Science Foundation of Innovative Research Groups Science Foundation of China under Grant 51221004, in part by the ARC DECRA and ARC Discovery Projects under Grant DE130100911 and Grant DP130101327, in part by the NSFC Funding under Grant 61332013, in part by the International Science and Technology Cooperation Projects under Grant 2016D10008, Grant 2013DFG12810, and Grant 2013C24027, in part by the Municipal Natural Science Foundation of Ningbo under Grant 2015A610119, in part by the Guangzhou Science and Technology Project under Grant 2016201604030034, in part by the Major Projects of Natural Science Research in Jiangsu Higher Education Institutions under Grant 14KJA520001, in part by the Jiangsu Production and Research Project under BY2015010-05, in part by the Natural Science Foundation of Zhejiang Province under Grant Y16F020002, in part by the Ningbo Natural Science Fund under Grant 2015A610119, in part by the International Science and Technology Cooperation Projects of Ningbo under Grant 2016D10008, and in part by the project of Research And Development of Intelligent Resource Allocation and Sharing Platform for Marine Electronic Information Industry under Grant 2017GY116.

ABSTRACT This paper proposes a method to address misreadings and consequent inadequacy of radio-frequency identification data for social insect monitoring. Six-month worth field experiment data were collected to demonstrate the application of the method. The data are transformed into a linear combination of the Gaussian model and curve-fitted using an evolutionary algorithm. This results show that the proposed method allows us to improve the quality of data that infer honey bee behavior at the colony level.

INDEX TERMS Apis mellifera, RFID, optimization, genetic algorithm, curve fitting, data quality.

I. INTRODUCTION

Pollinators play an integral role in food production, responsible for 1/3 of all commercial crop pollination, and with an estimated value of up to USD\$ 200 billion per annum [1], [2]. Honey bees, bumble bees, and some European stingless bees are exploited commercially as they are easily managed, and live in colonies with large numbers of individuals. Yet European honey bee (*Apis mellifera*) colony numbers in Europe and North America especially have been reported to decline over the last century [3]. Bee population decline has a detrimental impact on food security, floral biodiversity and abundance; the consequential impacts for the landscape as a result of bee losses have come to the attention of researchers.

In addition to their value in agriculture, honey bees are used as a model organism for a range of studies in

neurobiology, including cognition, perception, vision, genetics, and behavior [4]–[6]. Understanding the behavior of social insects at an individual level is very challenging, as each colony consists of a large number of individuals which, within the same caste, are highly similar visually. The effective collection of data pertaining to the location of individuals requires the unique identification of individuals by marking or fitting identifiers to each target individual.

Tagging insects with microdots [7], color patterns [8] and QR codes [9] is commonplace in entomological research. von Frisch [10], the Nobel Prize winner in Physiology or Medicine for 1973, painted bees to reveal the algorithm associated with the waggle dance of honey bees. This approach is also known to have been used by Darwin *et al.* [11]. While extremely useful and relatively

cheap, the use of microdots, QR codes and painting insects are time-consuming and labor intensive procedures, requiring either direct human observation or image recording and sophisticated processing.

More recently radio-frequency identification (RFID) devices have been applied to social insects [12]–[14]. In fact, RFID has become a popular tool in entomological research in the last decade and is being widely used in bee research groups in Europe [12], [15]–[20], North America [21], [22], China [23], [24], and Australia [25].

The main advantages of this technology are that thousands of bees in a single hive can be tagged so as to be individually identifiable, by a small number of operators and relatively quickly. Readings can be recorded constantly and without excessive human intervention. However, RFID technology is known to fail when collecting information on bees passing through gates (e.g. hives or feeders). This is likely to be caused by the short-range of reach of the antennas, and bees flying too fast in and out of the hives and feeders. Furthermore, the orientation and spatial positioning of bees as they enter hives could also cause misreadings due to the polarization of some RFID tags. A number of studies have reported the success rate of such systems when reading insect RFID tags, with results varying depending on, for example, experimental setup, reader capabilities, RFID tag (i.e. antenna) size, duty cycle, and the power of the electromagnetic signal. Here, we categorize the performance of the reader as: low (i.e. less than 80% success readings) [26], medium (i.e. between 80% and 90%) [21], [27], or high (i.e. greater than 90%) [12], [18] read efficiency. As a consequence of a low reading success rate, it becomes difficult to interpret what behavior an individual bee was involved in and the duration of that bee being inside and/or outside the hive.

Other concerns associated with tagging small insects are the additional weight of the tag, and the alterations necessary to the hive entry to accommodate readers and antennas. Because most social insects have a short adult lifespan, typically from a few days to a couple of months, studying them with the assistance of tags requires regular visits to colonies. As a consequence hives are opened regularly, changing the internal environment, stressing the colony as a whole and killing some of the insects in the process. Some insects are extremely sensitive to odors emitted by glues, and therefore insects fitted with tags may be attacked by other individuals.

Despite these limitations, RFID tagging of insects is becoming the most practical available tool to investigate individual behavior in a colony on a large scale. Using RFIDs is now more popular and, despite missing some readings, the behavior of the colony can be reasonably well captured in those experiments using electronic tags.

In this paper, we estimate the number of bees engaged in three different behaviors (e.g. by the entry, short mission, and foraging) by applying a classification method to the entire colony's activity data as tagged bee pass readers mounted at the hive entrance. This work addresses the key limitation of one popular method in the electronic tagging of social

insects: lost readings do not allow accurate interpretation of individual behavior.

The main contribution of this paper is the development of a method that allows the estimation of how many active bees in a hive will likely be foraging at a given moment (e.g. 3:30pm) or within a given time period (e.g. between 10 am and 2 pm). Here, “foraging” refers to those activities bees undertake in order to search for and collect resources for the colony, mainly food resources such as nectar and pollen, as well as hive resources including water and resin. The activities include searching for new food sources as well as exploiting current ones that the bee knows about, either as a result of previously visiting the source or by observing a waggle dance performed by another bee that has visited the source.

The paper is structured in the following manner: Section 2 describes the methodology and Section 3 presents the experimental results obtained. A discussion and conclusion will be provided in Section 4 and Section 5, respectively.

II. MATERIALS AND METHODS

A. FIELD EXPERIMENT

The dataset utilized for this work was obtained from a field experiment conducted in Tasmania, Australia. We set up four beehives with a radio-frequency identification (RFID) reader installed at each hive's entrance, as illustrated in Figure 1a. We visited the hives on a regular basis (e.g. once or twice a week) to tag bees with RFID tags (Section II-B). Using this setup, the bee passes through the hive entrance and is detected at a particular reader (Figure 1b); the individual detected, and the date and time of detection are recorded. The data is organized in individual daily CSV files, based on UTC time.

B. BEE TAGGING

Adult worker bees were tagged at the hive using $2.5 \times 2.5 \times 0.4$ mm RFID tags (Hitachi Chemical, Japan) secured to the thorax using cyanoacrylate super glue (Cyberbond LLC, Batavia, Illinois, USA). Each tag weights 2.4mg, one-third of an adult honey bee's maximum foraging weigh. Live bees were restrained against the honeycomb using a modified dissection probe, by applying gentle pressure between the thorax and abdomen. The tag was applied on the thorax between the wings, ensuring that both wing pairs were free before releasing the bee. The bee was then observed to be able to fly prior to proceeding to the next bee. Each tag is coded in hexadecimal format with a unique bee identification number, consisting of a range of parameters including the country in which the experiment is taking place, the hive number within an apiary the bee originated from, and the bee's species, strain, and caste.

C. CLASSIFICATION OF BEE BEHAVIORS

Bee activity is, for the purpose of this paper, the detection of a bee fitted with an RFID tag passing through the entry. Bee behavior is the interpretation of what the bee



FIGURE 1. The bee experiment conducted at Geeveston, Tasmania: (a) hive entrance; and (b) RFID reader installed underneath the entrance that detects the passage of tagged bees.

was actually doing. The assessment of what behavior a bee was exhibiting usually needs to take into account several successive RFID detections.

The daily activity of bees is greatly affected by external factors, especially the weather, and in particular temperature, precipitation, solar radiation, and wind speed [28]. If, for example, the hive becomes too hot, bees can leave the hive and “beard” at the entrance, or use their wings to ventilate at the entry of the hive. If the temperature is too low (i.e. typically below 10°C) bees will not leave the hive, and instead form a cluster on the comb of the brood nest to maintain the optimal brood temperature of 34.5°C . Bees do not fly during storms or during high wind speeds. If the external temperature is mild and on a sunny, calm day, bees will likely be found foraging in large numbers.

Either by observation or experience, honey bee activity data can be classified here into four behavior categories:

By The Entry (BTE): Bees classified as being “by the entry” are those with successive detections of the same bee by an RFID antenna at a maximum time interval between successive readings of less than three minutes. For example, a bee could be by the entry for 30 minutes, and it will be classified as BTE provided successive readings are within three minutes or less. This behavior is usually associated with hive maintenance, including cleaning and control of hive temperature, defense, or after returning from a foraging trip [29]–[31].

Short Mission (SM): Bees engaged in short missions are those with successive detections intervals between three and six minutes. This means the bees left the entry for a period of time no longer than six minutes. Bees engaged in short missions are those believed to be making short orientation or defecation flights, inspecting the surroundings, or engaged in defense activity [32].

Foraging (FG): Bees will be classified as foraging when the gap between successive detections is longer than six minutes. During the day, a bee may be detected many times and, in most cases, the time intervening between the first and last detections of the day will be

considered foraging, except when successive readings indicate the bee is by the entry or on short missions. Bees with recorded first and last detections will only be considered as foraging between sunrise and sunset. Foraging periods almost certainly incorporate periods of time when bees return from the field and stay inside the hive before going out again. Foraging is a crucial behavior of bees and can be associated with different roles: *scouts*, which spontaneously search for new food/water sources; *exploiters* and *water carriers* are individuals that make repetitive flights to food and water sources, respectively; *recruits* are individuals that search for food sources with a prior awareness of the approximate location of the source after observing a waggle dance [32]–[34]. In principle, using RFID technology, bees should be detected at every instance of leaving and returning to the hive and this would provide some insight into the duration bees spent foraging and how long they stay inside the hive before leaving again. However, misreadings of RFID systems make this task practically impossible. A way to overcome this difficulty is to associate the sporadic detection of bees to a foraging behavior. Therefore, the bee will be engaged in foraging activities for a long period of time, comprising several missions. Rather than recording each mission as a discrete event, the overall behavior is defined as a foraging role.

Departed bees (DB): Bees that leave the hive and never return, either because they die or because they swarm (including absconding) [35], [36]. Swarming was not observed in our hives during the experiment.

Table 1 presents a summary of bee behaviors as described above. For the purpose of this work, we also performed data curation to filter out erroneous data in accordance with empirical study based on field observations and an initial investigation of the data. One of the main issues relates to continuous readings with extremely short time intervals (according to our rules, this is classified as BTE). This can happen when a dead bee, with its RFID still attached, is located within the reading range of the reader. To overcome this, we configured

TABLE 1. Categorization of bee behaviors using RFID data. These behaviors reflect the duration of and time between consecutive readings and are used to report the most relevant results.

Bee Behaviour	Summary description
By the entry (BTE)	Successive readings of the same bee within less than three minutes indicates that this particular bee is near the RFID reader.
Short mission (SM)	Individual bee engaged in an activity of short duration (i.e. between three and six minutes). For example, making orientation or defecation flights and inspecting the hive surroundings.
Foraging (FG)	Individual bee engaged in activities to search for or exploit food/water sources. Example bee roles undertaking such activities are: scout, exploiter, recruit, water carrier.
Departed bee (DB)	The last detection of a bee in its lifetime.

the software in such a way that BTE reads with a duration of more than 30 minutes were discarded. Similarly, FG durations of more than six hours were omitted, as this is most likely attributable to instances of missed readings.

The classification criteria for bee behavior proposed above can be altered to accommodate other users' needs, without further limitations to the model implemented in the current work. For example, if a beekeeper or an entomologist understands their bees are at the entry for no longer than two minutes, it is possible to change the model to incorporate this observation. These behaviors are described for the European honey bee, *Apis mellifera*, and are not necessarily the same for other bee species.

D. INTERPRETING BEE POPULATION BEHAVIOR

The behaviors described above are applied to the recordings of bee activities to identify the behavior of each individual bee. After this step, a collective distribution for each bee behavior is generated. The diurnal distribution of all of the inferred behavior categories over the whole dataset resembles normal distributions (Figure 4). The resulting histogram is least-squared curve fitted using a Gaussian model (Section II-E).

The proposed method considers the overall activity as a linear combination of each type of possible activity (i.e. Gaussian Mixture Model) related to bee behavior. Therefore, the linear combination of Gaussian curves can be written as:

$$G_{ALL} = \alpha G_{BTE} + \beta G_{SM} + \theta G_{FG} \quad (1)$$

where each component of this equation is a Gaussian curve (G) expressed as:

$$G(x, BKG, I, T_{\mu}, T_{\sigma}) = I e^{-\frac{(x-T_{\mu})^2}{2 T_{\sigma}^2}} + BKG \quad (2)$$

where x is the data point (time of day in this case) to be estimated, BKG is a given background, I is the intensity, and T_{μ} and T_{σ} are the mean and the standard deviation of the distribution respectively. The parameters (α , β and θ) represent the relative number of bees involved in different behaviors

within the colony, calculated using the area under the curve for different Gaussian Probability Density Functions (PDFs).

For the purpose of this work, Gaussian parameters to be curve-fitted are as follows:

- i Background-effect (BKG): The data 'normalisation' which ensures that the PDF to be curve-fitted complies with the shape of a distribution. This is needed because brief visits to the hive entry (BTE and SM) occur regularly at night time. They are associated with bee defense or bees working to better climatise the colony (e.g., temperature or moisture control). Such events are considered a background (BKG) activity and are homogeneously distributed during the entire day and night. An example of such phenomena can be observed in Figure 4, where BTE detections occur between the hours of sunset and sunrise.
- ii Intensity (I): The height of the PDF indicating the overall probability of a particular activity taking place.
- iii The average of time in a day (T_{μ}): The time of day in which a particular bee activity is most likely to occur (highest I).
- iv The standard deviation of time of day (T_{σ}): The spread of the PDF of bee activity in a day.

Curve fitting is performed with the experimental data to determine the parameters of the Gaussian curves and their relative contribution to the overall distribution of behavior. Once the curve fitting is achieved, two key questions can be answered:

1. How many bees are performing a given behavior at a given moment of the day? This is determined by the relative intensity of each curve at the moment of interest.
2. How many bees are performing a given behavior during a given period of the day? This is determined by calculating the area under the curves during the period of interest.

E. CURVE FITTING USING GENETIC ALGORITHMS

Genetic Algorithm (GA) is a meta-heuristic method to generate a near-optimal solution for an optimization

TABLE 2. Summary mathematical representations for the data set (D).

Set Notation	Description	Representation		
		Item	Index	Example
A	Bee Activity	a_i	i, \dots, I	$A = \{a_1, a_2, \dots, a_i, \dots, a_I\}$
T	Time of day	t_j	j, \dots, J	$T = \{t_1, t_2, \dots, t_j, \dots, t_J\}$
D	Dataset	$d_{i,j}$		$D = \{\dots, d_{i,j}, \dots, d_{I,J}\}$

problem by evolving a pre-defined genetic representation (i.e. chromosome design), using natural selection process (e.g. selection, crossover, and mutation), towards a better solution. The following sub-sections discuss the requirement to perform GA in detail: (i) data notation; (ii) parameter initializations and constraints; (iii) chromosome design; and (iv) fitness function.

1) DATA NOTATION

First of all, we discuss the procedure we use to formalise the problem (Figure 4) in a mathematical way, and the notations to be used within the following sub-sections. The dataset is divided into two levels:

- i *Bee behavior*. Let $A = \{a_1, a_2, a_3\} = \{BTE, SM, FG\}$ be a list of distinct behaviors with $I = 3$ as discussed in Section II-C.
- ii *Time of day*. The time (t) in a day $T = \{t_1, t_2, \dots, t_j, \dots, t_J\}$ which is associated with its bin counts. In this work, we analyze the data in 30-minute intervals within a day resulting in $J = 24hr \div 30min = 48$ elements.

Based on the categorisation procedure above, we can now represent our dataset as: $D = \{d_{1,1}, d_{1,2}, \dots, d_{i,j}, \dots, d_{I,J}\}$ with each datum ($d_{i,j}$) representing the count/frequency of activity occurred at i^{th} activity A and j^{th} time of day T . A summary of these notations is given in Table 2.

We further denote dataset representations so that we could specify the particular category to obtain a sub-set of data from it. To illustrate this, some examples are shown below:

$$\begin{aligned}
 D &= \{d : d \in D\} \\
 &= \{d_{1,1}, d_{1,2}, \dots, d_{i,j}, \dots, d_{I,J}\} \\
 D_i &= D_{i=x} \\
 &= \{d : d \in D \wedge i = x\} \\
 &= \{d_{x,1}, d_{x,2}, \dots, d_{x,j}, \dots, d_{x,J}\} \\
 D_j &= D_{j=y} \\
 &= \{d : d \in D \wedge j = y\} \\
 &= \{d_{1,y}, d_{2,y}, \dots, d_{i,y}, \dots, d_{I,y}\}
 \end{aligned}$$

where x and y are artificial notations that depend on user input. Such representations will be used in the following sub-sections.

Since this work is based on a ‘data-driven’ modelling process, it is necessary to calculate the ‘importance’ of each

datum in order to compute the mean μ and standard deviation σ of a particular activity’s occurrence within a day. Therefore, the significance (denoted by the ‘weight’ $W = \{w_1, w_2, \dots, w_j, \dots, w_J\}$) of each datum (i.e. time index in day T) corresponds to the data availability ($d_{i,j}$) and is addressed by utilising the *weighted* mean (μ^*) and standard deviation (σ^*) equation as below:

$$\mu^*(V, W) = \frac{\sum_i^N w_i \cdot v_i}{\sum_i^N w_i} \quad (3)$$

$$\sigma^*(V, W) = \left[\frac{\sum_i^N w_i \cdot (v_i - \mu^*)^2}{\sum_i^N w_i} \right]^{\frac{1}{2}} \quad (4)$$

where V is a list of data with each datum denoted using v_i (i.e. $V = \{v_1, v_2, \dots, v_n, \dots, v_N\}$).

2) INITIAL PARAMETERS ESTIMATION AND CONSTRAINTS

The *BKG*-effect is estimated using the mean value of data D_i (i.e. i^{th} activity A) that holds the minimum Coefficient of Variation (CV) of the first n_{first} and last n_{last} data within time of day:

$$\arg \min_{n_{first}, n_{last}} CV \left((D_i)_{n_{first}} \cup (D_i)_{n_{last}} \right) \quad (5)$$

where n_{first} datum $(D_i)_{n_{first}} = \{D_{i,j=y} : y \in \mathbb{Z} \wedge y \leq n_{first}\}$ and n_{last} datum in time-of-day $(D_i)_{n_{last}} = \{D_{i,j=y} : y \in \mathbb{Z} \wedge n_{last} \leq y \leq J\}$. Thus, the BKG_i at activity i is obtained by:

$$BKG_i = \mu \left((D_i)_{n_{first}} \cup (D_i)_{n_{last}} \right) \quad (6)$$

and its constraint:

$$C(BKG_i) = \sigma \left((D_i)_{n_{first}} \cup (D_i)_{n_{last}} \right) \quad (7)$$

where $\mu()$ and $\sigma()$ corresponds to the minimised Equation 5.

Then, let time $T_{first,last} = \{t_x \in \mathbb{Z} \wedge n_{first} < y < n_{last}\}$ be a sub-set of T and its corresponding datum $U_i = \{D_{i,j=y} : y \in \mathbb{Z} \wedge n_{first} < y < n_{last}\}$. The remaining parameters for individual Gaussian G_i of distinct activity (a_i) are estimated in the following:

Estimation

$$\begin{array}{ll}
I_i & \mu(\{u : u \in U_i \wedge u \geq U_{i, \text{uniq}, (n-2)}\}) \\
(T_\mu)_i & \mu^*(\{T_{\text{first}, \text{last}}, U_i\}) \\
(T_\sigma)_i & \sigma^*(\{T_{\text{first}, \text{last}}, U_i\})
\end{array}$$

Constraint

$$\begin{array}{ll}
I_i & \sigma(\{u : u \in U_i \wedge u \geq U_{i, \text{uniq}, (n-2)}\}) \\
(T_\mu)_i & 30 \text{ min} \\
(T_\sigma)_i & \sigma^*(\{T_{\text{first}, \text{last}}, U_i\}) \div 2
\end{array}$$

where u denotes each datum within U_i and $U_{i, \text{uniq}, (n-2)}$ is the third largest ‘unique’ value within U_i (represented using an order statistic). Lastly, note that the lower and higher boundary (search space) for the optimization is in the form:

$$\begin{aligned}
C_{lo} &= \text{Estimation} - \text{Constraint} \\
C_{hi} &= \text{Estimation} + \text{Constraint}
\end{aligned} \quad (8)$$

3) GA’S CHROMOSOME DESIGN

Also called chromosome encoding and decoding, is a crucial step required to quantify the problem into an ‘individual’ for the optimization process. In this work, we designed a single individual using the following approach [37], [38]: Based on Equation 1, a complete distribution of the data consisting of $I = 3$ activities (BTE, SM, and FG) and 4 parameters (BKG, I, T_μ and T_σ , as in Equation 2) is required to generate one single Gaussian distribution. Therefore, in our case, one individual will consist of 12 elements ($3 \text{ activities} \times 4 \text{ parameters}$) with values between 0 and 1. In order to decode the value of a particular element within the individual, the following equation is employed:

$$p(x, C_{lo}, C_{hi}) = C_{lo} + x \cdot (C_{hi} - C_{lo}) \quad (9)$$

where x and p are the encoded and the decoded value of a particular Gaussian parameter respectively; C_{lo} and C_{hi} are the constraint values calculated from the previous section (Equation 8). Figure 2 demonstrates the design of a single individual with a decoding example for a parameter I of the Gaussian distribution G_i .




FIGURE 2. An illustration of the chromosome design utilized in this work, where each element within the individual holds a value between 0 and 1, and B denotes the background (BKG). An encoding and decoding example for the intensity (I_i) parameter of distribution G_i is also presented. In this case, assume that we have element I_i with value 0.7615 (encoded) within the individual which is equivalent to 205.32 (decoded) after applying Equation 9.

4) FITNESS FUNCTION

The quality of a particular individual is assessed by minimizing the sum of the chi-square (χ^2) function [37] for different

bee activities:

$$\text{fitness} = \sum_i^I \chi_i^2 \quad (10)$$

$$\chi_i^2 = \frac{1}{J - N_p} \sum_j^J \frac{(d_{i,j} - G_{i,j})^2}{d_{i,j} + 1} \quad (11)$$

where J is the number of elements in a day (see Table 2); N_p is the number of parameters to be optimized (four in this case); and $G_{i,j}$ is the estimated value using G_i at time t_j . Note that the $+1$ within the denominator on the right-side of Equation 11 is employed to avoid a divide-by-zero error, which could occur if the number of data points was extremely low.

III. EXPERIMENTAL RESULTS

This section provides the results obtained from the experiment. Figure 3 depicts the overview throughout the entire experiment, commenced on April 2nd and ended on November 11th, 2014. During the period, a total of 2,425 RFID tags were deployed; however, only 1,101 bees fitted with RFIDs were detected at least once after being tagged. Such a phenomenon can be explained by: (i) misreadings of the RFID system; (ii) the tag was not fitted properly so that the bee was able to remove it; and (iii) tags lost during the tagging process resulting from environmental conditions (e.g., on a windy day). The number of bees alive increases on the day our team members make a field visit to tag bees, and it is shown that approximately 30% of the tagged bees will be detected at least once on the following days.

Although data was collected between April and November, the analyses in the following sections were only undertaken on data collected between May and October to accommodate for the build up and decline of tagged bee numbers in the instrumented hives as shown in Figure 3, allowing for more robust results.

A. BEE ACTIVITY

Eight months of experimental data from 1,101 bees were recorded in CSV files. Bee detections were classified using the previously established rules (Section II-C) in order to assess each bee’s behavior at a given time. Once the behavior was determined, the data were grouped according to the time of day the activity occurred.

The daily distribution of bee behavior over the period of eight months is shown in Figure 4. The figure shows that bee activity starts to increase at approximately 7 am, and declines at approximately 8 pm. During that period, bees are most active between 12 pm and 1 pm. It is also observed that there are some detections before and after 7 am and 8 pm respectively which have been classified as by the entry. The analysis in this work removes Australian daylight savings time for consistency across data.

Figure 5 shows when bee activity is assigned to the category ‘departed bee’. Departed bees are those that left the hive and never returned. Occurrences after sunset and




FIGURE 3. Summary information of field visits for tagging bees and daily number of bees alive throughout the experiment. In this case, the lifespan of an individual bee (referred as ‘alive’ bee) is estimated from the first day it was tagged until the very last day of its detection.




FIGURE 4. Cumulative plot of six months experimental data, illustrating bee behavior distributions throughout the day.




FIGURE 5. Daily distribution of events associated with departed bees. These are events where bees are detected for the last time. The events occurring between dusk and dawn are likely to be of bees that died inside the hive and were transported out of the hive by other workers.

before sunrise are likely to be associated with bees that died inside the hive and were transported out by worker bees.

B. CURVE FITTING

Figure 6 shows the result for the curve fitting of each behavior (by the entry, short missions, foraging) for the entire




FIGURE 6. A visualisation produced by the curve-fitting program developed for this work that demonstrates the Gaussian PDF of distinct bee activities in a day. The x-axis shows the time of day and the y-axis is the frequency/count of activities. The dots (in different colors corresponding to the histogram values in Figure 4) represent the data to be curve fitted with BTE, SM, FG denoted in blue, green, red respectively. The solid lines are the curve fitted Gaussian PDFs for isolated activities (G_{BTE} , G_{SM} , G_{FG}) and the combined ones (G_{ALL} in black dashed-line) where the curve fitting used the optimization approach proposed in this work (Section II-E).




FIGURE 7. The proportion of bee behaviors relative to time of day.

period of the experiment. The sum of each behavior overlaps relatively well with the overall bee activity (black dots). It suggests that bees start foraging at approximately 7 am and finish at 8 pm. Around noon (between approximately 12 pm and 1 pm), most bees are involved in foraging role (e.g. exploiter, recruit, scout, water carrier); followed by by-the-entry activities (e.g. hive defense, temperature control); and lastly, on short missions (e.g. orientation flights, wandering around the nest).

The proportion of bees involved in different behaviors varies relative to time of day. Therefore, a normalisation of the curve fitted Gaussian PDF (Figure 6) is depicted in Figure 7. The normalized curve reveals that approximately 80% of the workers within the colony are engaged in BTE and $\approx 20\%$

are in SM during early morning (before sunrise) and late at night (after sunset).

C. BEE BEHAVIOUR

Once each individual bee behavior has been initially interpreted and following the curve fitting process, we can determine the proportion of bees performing a specific task at a given moment of the day or during a specific period of the day. If we consider the cohort of bees fitted with electronic tags to be representative of the entire bee population in a hive, it is possible to estimate how many bees would be engaging in, for example, foraging activities.

The result shows the proposed method is reasonable and the area under each curve should represent the number of bees




FIGURE 8. The proportion of bee colony behaviors for different months. The histogram (in yellow) presents the data availability for each month.

TABLE 3. Gaussian parameter values for each month of data reported in Figure 8. The 'Area(%)' column indicates the percentage of detections recorded of bees undertaking particular behaviors, relative to the total detections for that period.

Month		Gaussian Parameter				Area (%)
		BKG (Counts)	I (Counts)	T_{μ} (hh : mm)	T_{σ} (minutes)	
May	BTE	0.27	61.85	12:50	97	23.3
	SM	0.00	18.93	12:56	87	6.2
	FG	0.00	207.54	12:29	90	70.5
Jun	BTE	0.27	114.60	13:05	83	22.6
	SM	0.00	32.98	12:59	72	5.5
	FG	0.00	425.04	13:01	72	71.9
Jul	BTE	0.19	44.62	12:55	73	20.0
	SM	0.00	10.74	13:00	77	4.9
	FG	0.00	184.88	12:53	69	75.1
Aug	BTE	0.21	53.98	12:44	88	12.0
	SM	0.00	17.34	13:02	76	3.3
	FG	0.00	431.68	12:54	79	84.7
Sept	BTE	0.28	37.08	12:38	140	18.3
	SM	0.07	8.03	11:48	143	4.1
	FG	0.00	192.08	12:27	118	77.6
Oct	BTE	0.37	13.49	13:21	125	9.2
	SM	0.04	4.11	13:19	97	2.1
	FG	0.00	147.31	12:48	124	88.7

undertaking distinct behaviors throughout the day. Monthly proportions (from May to October) of bees foraging, in short missions or by the entry are shown in Figure 8 and values of the Gaussian parameters summarizing this data are given in Table 3. Figure 8 shows an overall increase of foraging behavior in the long term, and a decrease in by the entry behavior and short missions. Also, it shows that the data

availability dropped significantly in July, most probably due to the markedly decreased temperature during the winter period in Tasmania (as further discussed in the Discussion section).

Furthermore, based on Table 3, the standard deviation of the Gaussian parameter (T_{σ}) during winter period (i.e. June to August) is lower compared to other months

TABLE 4. Summary of the proposed bee behavioral model and its level of certainty for insects monitored under the current empirical study. The 'Threshold' column gives the cut-off points for two successive readings within the classification procedure for the raw bee detection data; whilst, the 'Duration' column indicates the range of valid bee behavior durations. For instance, BTE durations of more than 30 minutes will be omitted.

Bee Behaviour	Activities	Threshold	Duration	Certainty
By the entry	High frequency readings, defense, air conditioning of hive	$x \leq 3min$	$1sec < x \leq 30min$	High
Short mission	Orientation flights, walking around the hive	$3min < x \leq 6min$	$3min < x \leq 6min$	Medium
Foraging	Collecting/depositing food/water, scouting for new food resources	$x > 6min$	$6min < x \leq 6hr$	Low
Departed bee	Last detection of an individual, e.g. dead bee	–	–	High

(e.g. $T_{\sigma} < 80min$ overall). This was probably caused by (i) lower temperatures during the winter months that reduces bee activity, and (ii) the fact that the sun rises later and sets much earlier than in other seasons [28], [39].

IV. DISCUSSION

RFID systems where readers are installed in the field with limited power availability, operating with high reading frequency to capture every potential bee tagged leaving or returning the hive, in confined spaces like a bee-hive entry and with tags small enough to fit on bees are operationally challenging. Missing readings were inevitable and this fact makes the interpretation of each individual bee behavior very difficult. This work addresses this problem by developing a method that assigns a behavior for each bee based on roles and extrapolates that behavior for a cohort of bees doing the same activity.

Foraging behaviors were restricted to daylight hours, typically between 5 am and 8 pm; our data correlates well with nature, as bees will not forage when ambient temperature or solar radiation levels are too low. Additionally, the proportion of bees undertaking various behaviors varies over the course of the day. An increased probability of bees undertaking short missions and remaining by the entry is seen during the hours when bees are not actively engaged in foraging activities. This is likely due to forager role plasticity resulting in the reallocation of foragers to defensive or hygienic roles, or simply a matter of proportions altering as numbers of bees engaged in BTE and SM increases relative to FG (Figure 7).

Not only was activity variable over the daily cycle, but our six months of data include a distinct shift in behavior over the long term (Figure 8). The bee tagging period was commenced in April and terminated in October. In April and November tagged bee numbers within the hives were much lower than in the intermediate months as a result of tagged populations becoming established and dying out respectively (Figure 3). As a result of this, only data from May to October was included in the analysis. A significant decrease in readings was observed in July (Figure 8) due to predominantly cold weather, higher rainfall, and decreased solar radiation

when compared to the months of September through April. Furthermore, a proportional increase in foraging behaviors was observed over the course of the experiment. This may be due to an improvement in operator skill over time, resulting in more efficient tagging and a reduction in tagging-associated mortality, or adjustment of the colony to the colder winter temperature after the initial shock in July. Increased activity is expected, and observed, concurrent with the increase in temperatures into the spring months.

Bee behavior classification can be interpreted on the basis of the frequency of readings. By considering issue of misreadings in the RFID system, we are able to define the levels of certainty for the behavioral characterizations in our proposed model (Table 4).

High frequency reads are associated with the constant presence of a bee by the colony entry. This leads to a high degree of certainty about the assignment of a behavior of a bee to be 'by the entry' or 'short mission'. A departed bee is also very clear as the last recording of a bee could be confidently assigned as a bee that never returned to the hive.

The foraging behavior of bees is, by its very nature, complex. Bees could leave and return to the hive in missions lasting as long as an hour (and possibly longer) several times in a day. If the RFID system does not miss any readings, we would be able to confirm with absolute certainty when the bee left the hive, and when the same bee returned. We would also be able to say how long the bee was inside the hive between outdoor missions and for how long each mission lasted. When a single reading is missing, however, it becomes almost impossible to determine the bee's behavior at a given time with absolute certainty. With our approach we are able to estimate with some degree of confidence when the bees were engaged in foraging activities. This is possible because our rules are defined in such a way that single readings exclude BTE behaviors, and long durations between readings exclude SM behaviors, leaving only FG.

Figure 9 illustrates an example of an effort logging record throughout the lifetime of an individual bee. In this instance, this bee was tagged on 13th August 2014 at Hive 001 and last detected on 31st August 2014. It is very likely that this bee was tagged when it was very young because: (i) a 'single




FIGURE 9. Screenshot of a visualisation tool developed to analyze individual bee activity using the bee behavior classification described in Section II-C. Each row represents a day that the bee was active (i.e. detected at least once in that day); and the horizontal axis is the time of day, showing the times at which different bee behaviors occurred. Within each day, the times of sunrise (yellow) and sunset (green) are also indicated. For this example, the colored bars represent: (i) blue bar in the first row, the day the individual was tagged; (ii) red, single detection in that day; (iii) light blue, by the entry; (iv) orange, short mission; and (v) purple, foraging period.

detection' is observed on August 14th and 17th; and (ii) the bee started to forage on 18th August, despite the fact that it was tagged on August 13th. This bee is very likely to be a forager (e.g., scout, recruit, exploiter) throughout its lifetime. Note that the 'partitioned' foraging period indicates that there were detections with more than six minutes intervals between successive readings within its foraging period. This could, for example, be explained by assuming that the bee was either

out exploiting food sources during these intervals or that it was in the hive depositing nectar or pollen before undertaking further foraging activities.

V. CONCLUSIONS

The proposed method allows for a robust use of data from social insect monitoring based on RFID devices. Misreadings, which are common in RFID-based experiments, can be

better managed by combining insect behavior with activity data. The classification proposed in this paper (Section II-C) is based on results reported in the literature and on observation of our bees. However, depending on different bee species or other factors, it is possible to change the software configuration (i.e. model parameters) associated with bee activity.

Under traditional techniques, once the detection of the insect fails, RFID data become useless. The proposed method addresses this problem by assigning a given behavior for each tagged insect, then combining results for the entire tagged population using curve fitting based on genetic algorithms.

One limitation of this work is the inability to determine the number of data required in order to have a good representation of the results. For example, the curve fitting of monthly data (Section III-C) does not include April and November because those months do not have enough bee activity data for the curve-fitting purposes. Another limitation is the fact we have a small number of hives which limits the replicability of the experiments. And finally, the ideal calibration is to have another independent method to determine the activity of the bees. While we have scales in some of the hives, a camera, and an image processing technique could be used to determine when the bees leave and return as a mean to calibrate our results. Such an approach will be used in the future.

The method proposed in this paper could be used by other research groups using RFID to study social insects to better analyze the RFID data and overcome the issue of missed readings which are commonly experienced with electronic tagging. This has an important and positive implications for those using RFID data in insect behavior modelling. Study in the design of environmental sensor networks which involves animal-borne instruments as mobile sensor nodes [40] could also benefit from this work.

ACKNOWLEDGMENT

The authors would like to thank G. Allen and S. Quarrel from the Tasmanian Institute of Agriculture for their support to this paper.

REFERENCES

- [1] G. Degrandi-Hoffman, Y. Chen, E. W. Dejong, M. L. Chambers, and G. Hidalgo, "Effects of oral exposure to fungicides on honey bee nutrition and virus levels," *J. Econ. Entomol. Complete*, vol. 106, no. 6, pp. 2518–2528, 2015, doi: [10.1093/jee/fov251](#).
- [2] D. M. Bauer and I. S. Wing, "The macroeconomic cost of catastrophic pollinator declines," *Ecol. Econ.*, vol. 126, pp. 1–13, Jun. 2016, doi: [10.1016/j.ecolecon.2016.01.011](#).
- [3] D. van Engelsdorp et al., "Colony collapse disorder: A descriptive study," *PLoS ONE*, vol. 4, no. 8, p. e6481, 2009, doi: [10.1371/journal.pone.0006481](#).
- [4] M. V. Srinivasan, "Honey bees as a model for vision, perception, and cognition," *Annu. Rev. Entomol.*, vol. 55, no. 1, pp. 267–284, 2010, doi: [10.1146/annurev.ento.010908.164537](#).
- [5] A. Zayed and G. E. Robinson, "Understanding the relationship between brain gene expression and social behavior: Lessons from the honey bee," *Annu. Rev. Genetics*, vol. 46, no. 1, pp. 591–615, 2012, doi: [10.1146/annurev-genet-110711-155517](#).
- [6] T. D. Seeley, *The Wisdom of the Hive: The Social Physiology of Honey Bee Colonies*. Cambridge, MA, USA: Harvard Univ. Press, 1995. [Online]. Available: <http://opac.library.usyd.edu.au/record=2208589>
- [7] M. R. Whitehead and R. Peakall, "Microdot technology for individual marking of small arthropods," *Agricult. Forest Entomol.*, vol. 14, no. 2, pp. 171–175, 2012, doi: [10.1111/j.1461-9563.2011.00556.x](#).
- [8] J. Degen et al., "Exploratory behaviour of honeybees during orientation flights," *Animal Behav.*, vol. 102, pp. 45–57, Apr. 2015, doi: [10.1016/j.anbehav.2014.12.030](#).
- [9] D. P. Mersch, A. Crespi, and L. Keller, "Tracking individuals shows spatial fidelity is a key regulator of ant social organization," *Science*, vol. 340, no. 6136, pp. 1090–1093, 2013, doi: [10.1126/science.1234316](#).
- [10] K. von Frisch, "Decoding the language of the bee," *Science*, vol. 185, no. 4152, pp. 663–668, 1974, doi: [10.1126/science.185.4152.663](#).
- [11] C. Darwin and R. B. Freeman, *Charles Darwin on the Routes of Male Humble Bees*. London, U.K.: British Museum, 1968.
- [12] S. Streit, F. Bock, C. W. W. Pirk, and J. Tautz, "Automatic life-long monitoring of individual insect behaviour now possible," *Zoology*, vol. 106, no. 3, pp. 169–171, 2003, doi: [10.1078/0944-2006-00113](#).
- [13] A. Van Geystelen, K. Benaets, D. C. de Graaf, M. H. D. Larmuseau, and T. Wenseleers, "Track-a-Forager: A program for the automated analysis of RFID tracking data to reconstruct foraging behaviour," *Insectes Sociaux*, vol. 63, no. 1, pp. 175–183, 2016, doi: [10.1007/s00040-015-0453-z](#).
- [14] J. J. Bromenshenk, C. B. Henderson, R. A. Seccomb, P. M. Welch, S. E. Debnam, and D. R. Firth, "Bees as biosensors: Chemosensory ability, honey bee monitoring systems, and emergent sensor technologies derived from the pollinator syndrome," *Biosensors*, vol. 5, no. 4, pp. 678–711, 2015, doi: [10.3390/bios5040678](#).
- [15] I. M. M. Heidinger, M. D. Meixner, S. Berg, and R. Büchler, "Observation of the mating behavior of honey bee (*Apis mellifera* L.) queens using radio-frequency identification (RFID): Factors influencing the duration and frequency of nuptial flights," *Insects*, vol. 5, no. 3, pp. 513–527, 2014, doi: [10.3390/insects5030513](#).
- [16] C. W. Schneider, J. Tautz, B. Grünewald, and S. Fuchs, "RFID tracking of sublethal effects of two neonicotinoid insecticides on the foraging behavior of *Apis mellifera*," *PLoS ONE*, vol. 7, no. 1, p. e30023, 2012, doi: [10.1371/journal.pone.0030023](#).
- [17] R. J. Stelzer, R. Stanewsky, and L. Chittka, "Circadian foraging rhythms of bumblebees monitored by radio-frequency identification," *J. Biol. Rhythms*, vol. 25, no. 4, pp. 257–267, 2010, doi: [10.1177/0748730410371750](#).
- [18] M. Molet, L. Chittka, R. J. Stelzer, S. Streit, and N. E. Raine, "Colony nutritional status modulates worker responses to foraging recruitment pheromone in the bumblebee *Bombus terrestris*," *Behav. Ecol. Sociobiol.*, vol. 62, no. 12, pp. 1919–1926, 2008, doi: [10.1007/s00265-008-0623-3](#).
- [19] A. Decourtye et al., "Honeybee tracking with microchips: A new methodology to measure the effects of pesticides," *Ecotoxicology*, vol. 20, no. 2, pp. 429–437, 2011, doi: [10.1007/s10646-011-0594-4](#).
- [20] M. Moreau, P. Arrufat, G. Latil, and R. Jeanson, "Use of radio-tagging to map spatial organization and social interactions in insects," *J. Exp. Biol.*, vol. 214, pp. 17–21, Jan. 2011, doi: [10.1242/jeb.050526](#).
- [21] K. Ohashi, D. D'Souza, and J. D. Thomson, "An automated system for tracking and identifying individual nectar foragers at multiple feeders," *Behav. Ecol. Sociobiol.*, vol. 64, no. 5, pp. 891–897, 2010, doi: [10.1007/s00265-010-0907-2](#).
- [22] S. Sumner, E. Lucas, J. Barker, and N. Isaac, "Radio-tagging technology reveals extreme nest-drifting behavior in a eusocial insect," *Current Biol.*, vol. 17, no. 2, pp. 140–145, 2007, doi: [10.1016/j.cub.2006.11.064](#).
- [23] X. He, W. Wang, Q. Qin, Z. Zeng, S. Zhang, and A. B. Barron, "Assessment of flight activity and homing ability in Asian and European honey bee species, *Apis cerana* and *Apis mellifera*, measured with radio frequency tags," *Apidologie*, vol. 44, no. 1, pp. 38–51, 2013, doi: [10.1007/s13592-012-0156-7](#).
- [24] X.-J. He, L.-Q. Tian, X.-B. Wu, and Z.-J. Zeng, "RFID monitoring indicates honeybees work harder before a rainy day," *Insect Sci.*, vol. 23, no. 1, pp. 157–159, 2016, doi: [10.1111/1744-7917.12298](#).
- [25] C. J. Perry, E. Søvik, M. R. Myerscough, and A. B. Barron, "Rapid behavioral maturation accelerates failure of stressed honey bee colonies," *Proc. Nat. Acad. Sci. USA*, vol. 112, no. 11, pp. 3427–3432, 2015, doi: [10.1073/pnas.1422089112](#).
- [26] P. Tenczar, C. C. Lutz, V. D. Rao, N. Goldenfeld, and G. E. Robinson, "Automated monitoring reveals extreme interindividual variation and plasticity in honeybee foraging activity levels," *Animal Behav.*, vol. 95, pp. 41–48, Sep. 2014, doi: [10.1016/j.anbehav.2014.06.006](#).

- [27] E. J. H. Robinson, T. O. Richardson, A. B. Sendova-Franks, O. Feinerman, and N. R. Franks, "Radio tagging reveals the roles of corpulence, experience and social information in ant decision making," *Behav. Ecol. Sociobiol.*, vol. 63, no. 5, pp. 627–636, 2009, doi: [10.1007/s00265-008-0696-z](https://doi.org/10.1007/s00265-008-0696-z).
- [28] L. P. Polatto, J. Chaud-Netto, and V. V. Alves-Junior, "Influence of abiotic factors and floral resource availability on daily foraging activity of bees," *J. Insect Behav.*, vol. 27, no. 5, pp. 593–612, 2014, doi: [10.1007/s10905-014-9452-6](https://doi.org/10.1007/s10905-014-9452-6).
- [29] L. Bortolotti and C. Costa, "Chemical communication in the honey bee society," in *Frontiers in Neuroscience*. Boca Raton, FL, USA: CRC Press, 2014, pp. 147–210, doi: [10.1201/b16511-610.1201/b16511-6](https://doi.org/10.1201/b16511-610.1201/b16511-6).
- [30] M. Nouvian, J. Reinhard, and M. Giurfa, "The defensive response of the honeybee *Apis mellifera*," *J. Express Biol.*, vol. 219, no. 22, pp. 3505–3517, 2016, doi: [10.1242/jeb.143016](https://doi.org/10.1242/jeb.143016).
- [31] S. Boutin, M. Alburaki, P.-L. Mercier, P. Giovenazzo, and N. Derome, "Differential gene expression between hygienic and non-hygienic honeybee (*Apis mellifera* L.) hives," *BMC Genomics*, vol. 16, p. 500, Jul. 2015, doi: [10.1186/s12864-015-1714-y](https://doi.org/10.1186/s12864-015-1714-y).
- [32] E. A. Capaldi et al., "Ontogeny of orientation flight in the honeybee revealed by harmonic radar," *Nature*, vol. 403, no. 6769, pp. 537–540, Feb. 2000, doi: [10.1038/35000564](https://doi.org/10.1038/35000564).
- [33] H. de Vries and J. C. Biesmeijer, "Modelling collective foraging by means of individual behaviour rules in honey-bees," *Behav. Ecol. Sociobiol.*, vol. 44, no. 2, pp. 109–124, 1998, doi: [10.1007/s002650050522](https://doi.org/10.1007/s002650050522).
- [34] G. E. Robinson, B. A. Underwood, and C. E. Henderson, "A highly specialized water-collecting honey bee," *Apidologie*, vol. 15, no. 3, pp. 355–358, 1984, doi: [10.1051/apido:19840307](https://doi.org/10.1051/apido:19840307).
- [35] J. D. Villa, "Swarming behavior of honey bees (Hymenoptera: Apidae) in Southeastern Louisiana," *Ann. Entomol. Soc. Amer.*, vol. 97, no. 1, pp. 111–116, 2004, doi: [10.1603/0013-8746\(2004\)097\[0111:sbohbj\]2.0.co;2](https://doi.org/10.1603/0013-8746(2004)097[0111:sbohbj]2.0.co;2).
- [36] H. R. Hepburn, S. L. Reece, P. Neumann, R. F. A. Moritz, and S. E. Radloff, "Absconding in honeybees (*Apis mellifera*) in relation to queen status and mode of worker reproduction," *Insectes Sociaux*, vol. 46, no. 4, pp. 323–326, 1999, doi: [10.1007/s000400050152](https://doi.org/10.1007/s000400050152).
- [37] F. Susanto and P. de Souza, Jr., "Mössbauer spectral curve fitting combining fundamentally different techniques," *Nucl. Instrum. Methods Phys. Res. Sect. B, Beam Interact. Mater. Atoms*, vol. 385, pp. 40–45, Oct. 2016, doi: [10.1016/j.nimb.2016.08.011](https://doi.org/10.1016/j.nimb.2016.08.011).
- [38] F. Susanto, "High resolution environmental modelling application using a swarm of sensor nodes," Ph.D. dissertation, Dept. Eng. Sci., Victoria Univ., Footscray VIC, Australia, 2017. [Online]. Available: vuir.vu.edu.au/34678/1/SUSANTO%20Ferry-thesis.pdf
- [39] P. Węgrzynowicz, D. Gerula, M. Biełkowska, and B. Panasiuk, "Causes and scale of winter flights in honey bee (*Apis mellifera carnica*) colonies," *J. Apicultural Sci.*, vol. 58, no. 1, pp. 135–143, 2014, doi: [10.2478/jas-2014-0014](https://doi.org/10.2478/jas-2014-0014).
- [40] S. Budi, P. de Souza, G. Timms, F. Susanto, V. Malhotra, and P. Turner, "Mobile platform sampling for designing environmental sensor networks," *Environ. Monit. Assessment*, vol. 190, no. 3, p. 130, 2018, doi: [10.1007/s10661-018-6510-0](https://doi.org/10.1007/s10661-018-6510-0).



THOMAS GILLARD was born in Sydney, NSW, Australia, in 1993. He received the B.Sc. degree (Hons.) in veterinary science and the B.V.Sc. degree (Hons.) from The University of Sydney in 2015 and 2017, respectively.

He is currently pursuing the Ph.D. degree in honey bee reproductive diseases with the Behavior and Genetics of Social Insects Laboratory, The University of Sydney. His research experience began during his undergraduate years, when he commenced part-time work with The University of Sydney. He was a Research Assistant with the Behavior and Foraging Ecology Laboratory, The University of Sydney, where he was involved in field-based research studies and general laboratory maintenance from 2015 to 2017. He was a Research Assistant at Data61, Commonwealth Scientific and Industrial Research Organization from 2015 to 2017, where he was involved in additional experience with laboratory work and data analysis, and a large role in apiary management and bee work.

He has a considerable amount of experience operating in research laboratories, field-based research studies, bee husbandry, management, and processes within the commercial apiculture setting. His veterinary training provides him a unique perspective moving into this field, with a focus on the holistic management of disease, from recognition of risk factors and prevention of disease, through principles of diagnosis and treatment. He has a view to applying the veterinary training to challenges afflicting the apiculture industry, both locally and globally. His primary career focuses are in the field of honey bee health and disease research.

Mr. Thomas is a member of the Australian Veterinary Association, the NSW Apiarists Association, the Society for Invertebrate Pathology, the International Union for the Study of Social Insects, and the Veterinary Invertebrate Society. He received the HR Carne Award for his B.Sc. (Vet) thesis, which explored the evolution of honey bee viruses.



PAULO DE SOUZA, JR was committed to excellence in research, and successful track record in BD&C of research products. He held senior positions at government agencies, universities, and industry. He has extensive hands-on experience in managing large teams, complex operations, and large-scale projects. He is an internationally recognized and awarded researcher and a research manager in the fields of geochemistry, sensors, sensor networks, and ICT with 20 years of experience in industry and research organizations. He is a deep expertise in research and development, innovation, and business development gained by leading challenging initiatives and academic pursuits. He is passionate about translating science into meaningful outcomes for government, business, and the community. He has authored or co-authored over 200 peer-review publications, three patents, and four books published.

He is a deep expertise in research and development, innovation, and business development gained by leading challenging initiatives and academic pursuits. He is passionate about translating science into meaningful outcomes for government, business, and the community. He has authored or co-authored over 200 peer-review publications, three patents, and four books published.



FERRY SUSANTO received the Ph.D. degree from Victoria University at Australia, Australia, in 2017. His main research interests include optimization problem, environmental modeling, data science, and artificial intelligence.



BENITA VINCENT received the Ph.D. degree from the University of Tasmania. She was involved in teaching a number of practical units of the bachelor's degree in aquaculture. She currently continues her research in amoebic gill disease with Commonwealth Scientific and Industrial Research Organization, Marine and Atmospheric Research, Hobart. Her research background is in aquatic animal health and predominately focused on amoebic gill disease of Atlantic salmon.



SETIA BUDI is currently pursuing the Ph.D. degree with the University of Tasmania, Australia. His main research interests include optimization problem, environmental monitoring, data science, and computer vision.



AURO ALMEIDA was involved in several areas of research and planning in the forestry production sector in Brazil for 14 years, where he developed and applied extensive spatial modeling to predict forest growth and water use. He joined the Commonwealth Scientific and Industrial Research Organization as a Senior Research Scientist in 2006, where he has been leading and participating in projects related to modeling forest production, catchment, and water resources management and the impact of climate change in Australia, South America, and Southeast Asia.

He has been developing multiple projects in the forestry sector in Australia and South America with the development of long-term research projects with several forestry companies and governmental agencies in the fields of forestry and pasture modeling, water use and water-use efficiency, catchment management, and strategic research plan. He has been in Southeast predicting the effects of climate change on Eucalyptus and Acacia plantations growth. From 1988 to 1991, he was a Researcher with the Italian National Energy Agency, IFAD (United Nations), and Ecotherm Spa, Italy. Since 2016, he has been the Project Deputy Leader of the Sustainable Development Investment Portfolio Project, promoting the development of basin plan, water management, and capacity building for Kamala Basin in Nepal.



GUSTAVO PESSIN received the Ph.D. degree in computer science from the University of São Paulo. He did his post-doctoral research at the Massachusetts Institute of Technology. He is currently a Research Scientist with the Vale Institute of Technology for Sustainable Development. He has interest in the area of machine learning, data science, and wireless sensor networks.



HELDER ARRUDA received the master's degree in sustainable use of natural resources from the Vale Institute of Technology for Sustainable Development with a focus on data from micro-sensors installed in bees, in partnership project with Brazil and Australia. He was a Systems Analyst with the University of Amazônia and a Programmer with Grupo Reicon. He has experience in systems development and maintenance and in relational databases. He is currently an Assistant

Researcher of the research line Biodiversity and Ecosystem Services in the Vale Institute of Technology for Sustainable Development. He currently deals with scientific experiments aimed at environmental monitoring and data analysis, using sensor networks and machine learning.

RAYMOND N. WILLIAMS, photograph and biography not available at the time of publication.



ULRICH ENGELKE received the Dipl.Ing. degree in electrical engineering from RWTH Aachen University, Germany, in 2004, and the Ph.D. degree in telecommunications from the Blekinge Institute of Technology, Karlskrona, Sweden, in 2010. He held a post-doctoral position at the University of Nantes, France. He was with the Visual Experiences Group, Philips Research, The Netherlands, and Philips Color Kinetics Inc., USA. In 2016, he was with IBM Research Almaden Laboratories,

San Jose, CA, USA. He is currently with Data61, Commonwealth Scientific and Industrial Research Organization, where he is involved in the interface between humans and data. He designs and evaluates interactive systems and methods that enable an efficient path from data to insight for improved decision making. His research focuses specifically on visual analytics and information visualization techniques, immersive analytics through virtual and mixed reality interfaces, quantitative user experience, and task performance evaluation.



PETER MARENDY is currently pursuing the part-time Ph.D. degree in addition to his Commonwealth Scientific and Industrial Research Organization duties that is looking to utilize context and data analytics to provide reasoning over both legacy and streaming sensor data. He was a Software Engineer on a number of projects, including Tailored Diet Information (Food and Nutritional Sciences), Residential Scale Energy Services, the Web Interface-Energy Systems Models Project

(Energy Transformed Flagship), the Museum Robot Project (the Commonwealth Scientific and Industrial Research Organization Autonomous Systems Laboratory, Australian National Museum, and the Department of Broadband, Communications and the Digital Economy), and Enhanced Situation Awareness (Digital Productivity and Services). He was with the VizzzBees Project (Data61), which is working toward producing a visual analytics platform for large and varied data sets. He was a Tasmanian Representative for the ICT Centre Project Management Working Group. He was also required to devise strategies and advice for assisting project leaders in areas identified for improvement and identifying potential project leaders. He actively facilitated the four-monthly project reviews for TasICT and strongly involved in the science review process for the ICT Centre. He is currently with the Microsensing Group. He is currently involved in Bees with Backpacks, the Global Initiative for Honey bee Health, and the Probing Biosystems Future Science Platform-Implantables Projects. He coaches and advises project leaders on matters of management, such as tools, methodologies, processes, and procedures.



PASCAL HIRSCH is currently pursuing the Ph.D. degree in computer science with the University of Tasmania. His research interests are within the area of data science, embedded system, and signal processing.



JING HE received the Ph.D. degree from the Academy of Mathematics and System Science, Chinese Academy of Sciences, in 2006. She was with the University of Chinese Academy of Sciences, China. She is currently a Professor with the College of Engineering and Science, Victoria University. She is currently an Adjunct Professor with the University of Chinese Academy of Science and the Nanjing University of Finance and Economics. She has published over 60 research papers in refereed international journals and conference proceedings, including *Information System*, *Information Science*, the *ACM Transactions on Internet Technology*, the IEEE TRANSACTION ON KNOWLEDGE AND DATA ENGINEERING, *Plos One*, *The Computer Journal*, *Computers and Mathematics With Applications*, and *Concurrency and Computation: Practice and Experience*. She has been active in the areas of robotics programming, virtual reality, big data analytics, sensor networks, web service, and some industry field, such

as e-Health, petroleum exploration and development, and water recourse management. She is a Workshop Co-Chair of APWeb 2008, WI 2009, MCDM 2009, WISE 2012, and HIS 2012, and the Program Chair of HIS 2011, 2012, and 2013 conferences. She served on three program committees of international conferences, such as the International Conference on Computational Science (ICCS), The IEEE International Conference on Data Mining (ICDM), and the International Symposium on Knowledge and Systems Science (KSS). She has been serving as an external reviewers for several international journals and conferences, such as *Management Science*, *The Computer Journal*, the IEEE TRANSACTION ON SYSTEMS, MAN, AND CYBERNETICS, the *International Journal of Information Technology and Decision Making*, the *Journal of Management Review* (in Chinese), *Decision Support System*, *Science* (in China), WISE, ICCS, ICDM, KSS, APWeb, and PAKDD. She received research fund from ARC DECRA, ARC Discovery, ARC Linkage, the Youth Science Fund of Chinese Academy of Sciences, and the National Science Foundation of China.

...