



VICTORIA UNIVERSITY
MELBOURNE AUSTRALIA

Some new inequalities of Hermite-Hadamard type for GA-convex functions

This is the Published version of the following publication

Dragomir, Sever S (2018) Some new inequalities of Hermite-Hadamard type for GA-convex functions. *Annales Universitatis Mariae Curie-Sklodowska, sectio A – Mathematica*, 72 (1). pp. 55-68. ISSN 0365-1029

The publisher's official version can be found at
<https://journals.umcs.pl/a/article/view/7331>

Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/37901/>

SEVER S. DRAGOMIR

**Some new inequalities
of Hermite–Hadamard type
for GA -convex functions**

ABSTRACT. Some new inequalities of Hermite–Hadamard type for GA -convex functions defined on positive intervals are given. Refinements and weighted version of known inequalities are provided. Some applications for special means are also obtained.

1. Introduction. Let $J \subset (0, \infty)$ be an interval; a real-valued function $f : J \rightarrow \mathbb{R}$ is said to be GA -convex (concave) on J (see for instance [2]), if

$$(1.1) \quad f(x^{1-\lambda}y^\lambda) \leq (\geq) (1-\lambda)f(x) + \lambda f(y)$$

for all $x, y \in J$ and $\lambda \in [0, 1]$.

Since the condition (1.1) can be written as

$$(1.2) \quad \begin{aligned} & f \circ \exp((1-\lambda)\ln x + \lambda\ln y) \\ & \leq (\geq) (1-\lambda)f \circ \exp(\ln x) + \lambda f \circ \exp(\ln y), \end{aligned}$$

then we observe that $f : J \rightarrow \mathbb{R}$ is GA -convex (concave) on J if and only if $f \circ \exp$ is convex (concave) on $\ln J := \{\ln z, z \in J\}$. If $J = [a, b]$, then $\ln J = [\ln a, \ln b]$.

It is known that the function $f(x) = \ln(1+x)$ is GA -convex on $(0, \infty)$ [2].

2010 *Mathematics Subject Classification.* 26D15, 25D10.

Key words and phrases. Convex functions, integral inequalities, GA -convex functions, Hermite–Hadamard inequalities.

For real and positive values of x , the *Euler gamma* function Γ and its *logarithmic derivative* ψ , the so-called *digamma function*, are defined by

$$\Gamma(x) := \int_0^{\infty} t^{x-1} e^{-t} dt \text{ and } \psi(x) := \frac{\Gamma'(x)}{\Gamma(x)}.$$

It has been shown in [28] that the function $f : (0, \infty) \rightarrow \mathbb{R}$ defined by

$$f(x) = \psi(x) + \frac{1}{2x}$$

is *GA-concave* on $(0, \infty)$ while the function $g : (0, \infty) \rightarrow \mathbb{R}$ defined by

$$g(x) = \psi(x) + \frac{1}{2x} + \frac{1}{12x^2}$$

is *GA-convex* on $(0, \infty)$.

Example 1. If $[a, b] \subset (0, \infty)$ and the function $g : [\ln a, \ln b] \rightarrow \mathbb{R}$ is convex (concave) on $[\ln a, \ln b]$, then the function $f : [a, b] \rightarrow \mathbb{R}$, $f(t) = g(\ln t)$ is *GA-convex* (concave) on $[a, b]$.

We recall that the classical Hermite–Hadamard inequality states that

$$(1.3) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(t) dt \leq \frac{f(a) + f(b)}{2}$$

for any convex function $f : [a, b] \rightarrow \mathbb{R}$.

For related results, see [1]–[6], [8]–[11], [12]–[18] and [20]–[27].

The *identric mean* $I(a, b)$ is defined by

$$I(a, b) := \frac{1}{e} \left(\frac{b^b}{a^a} \right)^{\frac{1}{b-a}}$$

while the *logarithmic mean* is defined by

$$L(a, b) := \frac{b-a}{\ln b - \ln a}$$

where $0 < a < b$.

In [28] the authors obtained the following Hermite–Hadamard type inequality.

Theorem 1. *If $b > a > 0$ and $f : [a, b] \rightarrow \mathbb{R}$ is a differentiable *GA-convex* (concave) function on $[a, b]$, then*

$$(1.4) \quad \begin{aligned} f(I(a, b)) &\leq (\geq) \frac{1}{b-a} \int_a^b f(t) dt \\ &\leq (\geq) \frac{b-L(a, b)}{b-a} f(b) + \frac{L(a, b)-a}{b-a} f(a). \end{aligned}$$

The differentiability of the function is not necessary in Theorem 1 for the first inequality (1.4) to hold, as shown in [10].

If we take $\lambda = \frac{1}{2}$ in the definition (1.1) of GA -convex (concave) function on $[a, b]$, then we have

$$(1.5) \quad f(\sqrt{ab}) \leq (\geq) \frac{f(a) + f(b)}{2}.$$

The following refinement of (1.5), which is an inequality of Hermite–Hadamard type, holds (see [25] for an extension for GA h -convex functions):

Theorem 2. *Let $f : [a, b] \subset (0, \infty) \rightarrow \mathbb{R}$ be a GA -convex (concave) function on $[a, b]$. Then we have*

$$(1.6) \quad f(\sqrt{ab}) \leq (\geq) \frac{1}{\ln b - \ln a} \int_a^b \frac{f(t)}{t} dt \leq (\geq) \frac{f(a) + f(b)}{2}.$$

Motivated by the above results we provide in the following a refinement of (1.6) for a division of the interval $[a, b]$. We also establish a weighted version that generalizes the inequalities (1.4) and (1.6) and provides upper and lower bounds for the moments

$$\frac{1}{b-a} \int_a^b t^p f(t) dt, \text{ for } p \in \mathbb{R} \text{ with } p \neq 0, -1$$

of the GA -convex (concave) function on $[a, b]$.

2. Some refinements. In 1994, [5] (see also [17, p. 22]) we proved the following refinement of Hermite–Hadamard inequality. For the sake of completeness we give here a direct proof that is different from the one in [5].

Lemma 1. *Let $g : [c, d] \rightarrow \mathbb{R}$ be a convex function on $[c, d]$. Then for any division $c = x_0 < x_1 < \dots < x_{n-1} < x_n = d$ with $n \geq 1$ we have the inequalities*

$$(2.1) \quad \begin{aligned} g\left(\frac{c+d}{2}\right) &\leq \frac{1}{d-c} \sum_{i=0}^{n-1} (x_{i+1} - x_i) g\left(\frac{x_{i+1} + x_i}{2}\right) \\ &\leq \frac{1}{d-c} \int_c^d g(x) dx \\ &\leq \frac{1}{d-c} \sum_{i=0}^{n-1} (x_{i+1} - x_i) \frac{g(x_i) + g(x_{i+1})}{2} \\ &\leq \frac{1}{2} [g(c) + g(d)]. \end{aligned}$$

Proof. Using the Hermite–Hadamard inequality on the interval $[x_i, x_{i+1}]$, $i \in \{0, \dots, n-1\}$, we have

$$(2.2) \quad \begin{aligned} (x_{i+1} - x_i) g\left(\frac{x_{i+1} + x_i}{2}\right) &\leq \int_{x_i}^{x_{i+1}} g(x) dx \\ &\leq (x_{i+1} - x_i) \frac{g(x_i) + g(x_{i+1})}{2} \end{aligned}$$

for any $i \in \{0, \dots, n-1\}$.

Summing in (2.2) over $i \in \{0, \dots, n-1\}$ and dividing by $d - c$, we get the second and the third inequalities in (2.1).

Since for $p_i := \frac{x_{i+1} - x_i}{d - c} \geq 0$ we have $\sum_{i=0}^{n-1} p_i = 1$, then by Jensen's inequality we have

$$\begin{aligned} \frac{1}{d - c} \sum_{i=0}^{n-1} (x_{i+1} - x_i) g\left(\frac{x_{i+1} + x_i}{2}\right) &\geq g\left(\frac{1}{d - c} \sum_{i=0}^{n-1} (x_{i+1} - x_i) \left(\frac{x_{i+1} + x_i}{2}\right)\right) \\ &= g\left(\frac{1}{d - c} \sum_{i=0}^{n-1} \left(\frac{x_{i+1}^2 - x_i^2}{2}\right)\right) \\ &= g\left(\frac{d^2 - c^2}{2(d - c)}\right) = g\left(\frac{c + d}{2}\right), \end{aligned}$$

which proves the first inequality in (2.1).

For a convex function $g : [c, d] \rightarrow \mathbb{R}$ we have

$$g(x) = g\left(\frac{(d - x)c + (x - c)d}{d - c}\right) \leq \frac{(d - x)g(c) + (x - c)g(d)}{d - c}$$

for any $x \in [c, d]$.

This implies that

$$g(x_i) \leq \frac{1}{d - c} [(d - x_i)g(c) + (x_i - c)g(d)]$$

and

$$g(x_{i+1}) \leq \frac{1}{d - c} [(d - x_{i+1})g(c) + (x_{i+1} - c)g(d)]$$

for any $i \in \{0, \dots, n-1\}$.

Therefore,

$$\begin{aligned} &\sum_{i=0}^{n-1} (x_{i+1} - x_i) \frac{g(x_i) + g(x_{i+1})}{2} \\ &\leq \frac{1}{2(d - c)} \sum_{i=0}^{n-1} (x_{i+1} - x_i) [(d - x_i)g(c) \\ &\quad + (x_i - c)g(d) + (d - x_{i+1})g(c) + (x_{i+1} - c)g(d)] \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{d-c} \left[\left(d(d-c) - \frac{1}{2}(d^2 - c^2) \right) g(c) + \left(\frac{1}{2}(d^2 - c^2) - c(d-c) \right) g(d) \right] \\
&= (d-c) \frac{g(c) + g(d)}{2},
\end{aligned}$$

and the last part of (2.1) is proved. \square

Theorem 3. Let $f : [a, b] \subset (0, \infty) \rightarrow \mathbb{R}$ be a GA-convex function on $[a, b]$. Then for any division $a = t_0 < t_1 < \dots < t_{n-1} < t_n = b$ with $n \geq 1$ we have the inequalities

$$\begin{aligned}
(2.3) \quad f(\sqrt{ab}) &\leq \frac{1}{\ln b - \ln a} \sum_{i=0}^{n-1} (\ln t_{i+1} - \ln t_i) f(\sqrt{t_i t_{i+1}}) \\
&\leq \frac{1}{\ln b - \ln a} \int_a^b \frac{f(s)}{s} ds \\
&\leq \frac{1}{\ln b - \ln a} \sum_{i=0}^{n-1} (\ln t_{i+1} - \ln t_i) \frac{f(t_i) + f(t_{i+1})}{2} \\
&\leq \frac{1}{2} [f(a) + f(b)].
\end{aligned}$$

Proof. If $f : [a, b] \subset (0, \infty) \rightarrow \mathbb{R}$ is a GA-convex function on $[a, b]$, then $f \circ \exp$ is convex on $[\ln a, \ln b]$. Let $c = \ln a$, $d = \ln b$ and the division $x_i = \ln t_i$, $i \in \{0, \dots, n-1\}$ of the interval $[c, d]$.

If we write the inequality (2.1) for $g = f \circ \exp$ on the interval $[c, d]$ and for the division

$$\ln a = \ln t_0 < \ln t_1 < \dots < \ln t_{n-1} < \ln t_n = \ln b,$$

we have

$$\begin{aligned}
(2.4) \quad &(f \circ \exp) \left(\frac{\ln a + \ln b}{2} \right) \\
&\leq \frac{1}{\ln b - \ln a} \sum_{i=0}^{n-1} (\ln t_{i+1} - \ln t_i) (f \circ \exp) \left(\frac{\ln t_{i+1} + \ln t_i}{2} \right) \\
&\leq \frac{1}{\ln b - \ln a} \int_{\ln a}^{\ln b} (f \circ \exp)(x) dx \\
&\leq \frac{1}{\ln b - \ln a} \sum_{i=0}^{n-1} (\ln t_{i+1} - \ln t_i) \frac{(f \circ \exp)(\ln t_i) + (f \circ \exp)(\ln t_{i+1})}{2} \\
&\leq \frac{1}{2} [(f \circ \exp)(\ln a) + (f \circ \exp)(\ln b)],
\end{aligned}$$

that is equivalent to

$$\begin{aligned}
 f(\sqrt{ab}) &\leq \frac{1}{\ln b - \ln a} \sum_{i=0}^{n-1} (\ln t_{i+1} - \ln t_i) f(\sqrt{t_i t_{i+1}}) \\
 &\leq \frac{1}{\ln b - \ln a} \int_{\ln a}^{\ln b} (f \circ \exp)(x) dx \\
 (2.5) \quad &\leq \frac{1}{\ln b - \ln a} \sum_{i=0}^{n-1} (\ln t_{i+1} - \ln t_i) \frac{f(t_i) + f(t_{i+1})}{2} \\
 &\leq \frac{1}{2} [f(a) + f(b)].
 \end{aligned}$$

By using the change of variable $\exp(x) = t$, we have $x = \ln t$, $dx = \frac{dt}{t}$ and

$$\int_{\ln a}^{\ln b} (f \circ \exp)(x) dx = \int_a^b \frac{f(t)}{t} dt$$

and by (2.5) we get the desired result (2.3). \square

Corollary 1. Let $f : [a, b] \subset (0, \infty) \rightarrow \mathbb{R}$ be a GA-convex function on $[a, b]$. If $a \leq t \leq b$, then we have

$$\begin{aligned}
 f(\sqrt{ab}) &\leq \frac{1}{\ln b - \ln a} [(\ln t - \ln a) f(\sqrt{at}) + (\ln b - \ln t) f(\sqrt{tb})] \\
 &\leq \frac{1}{\ln b - \ln a} \int_a^b \frac{f(s)}{s} ds \\
 (2.6) \quad &\leq \frac{1}{2} \left[f(t) + \frac{(\ln t - \ln a) f(a) + (\ln b - \ln t) f(b)}{\ln b - \ln a} \right] \\
 &\leq \frac{1}{2} [f(a) + f(b)].
 \end{aligned}$$

Remark 1. Let $f : [a, b] \subset (0, \infty) \rightarrow \mathbb{R}$ be a GA-convex function on $[a, b]$. For a division $0 = \lambda_0 < \lambda_1 < \dots < \lambda_{n-1} < \lambda_n = 1$ with $n \geq 1$ of the interval $[0, 1]$, consider the division

$$(2.7) \quad t_i := a \left(\frac{b}{a} \right)^{\lambda_i} = a^{1-\lambda_i} b^{\lambda_i}, \quad i \in \{0, \dots, n-1\}$$

of the interval $[a, b]$.

Observe that

$$\begin{aligned}
 \ln t_{i+1} - \ln t_i &= \ln \left[a \left(\frac{b}{a} \right)^{\lambda_{i+1}} \right] - \ln \left[a \left(\frac{b}{a} \right)^{\lambda_i} \right] \\
 &= \ln a + \lambda_{i+1} \ln \left(\frac{b}{a} \right) - \ln a - \lambda_i \ln \left(\frac{b}{a} \right) \\
 &= (\ln b - \ln a) (\lambda_{i+1} - \lambda_i)
 \end{aligned}$$

and

$$\sqrt{t_i t_{i+1}} = \sqrt{a^{1-\lambda_i} b^{\lambda_i} a^{1-\lambda_{i+1}} b^{\lambda_{i+1}}} = a^{1-\frac{\lambda_i+\lambda_{i+1}}{2}} b^{\frac{\lambda_i+\lambda_{i+1}}{2}}$$

for any $i \in \{0, \dots, n-1\}$.

If we write the inequality (2.3) for the division (2.7) we get

$$\begin{aligned} f(\sqrt{ab}) &\leq \sum_{i=0}^{n-1} (\lambda_{i+1} - \lambda_i) f\left(a^{1-\frac{\lambda_i+\lambda_{i+1}}{2}} b^{\frac{\lambda_i+\lambda_{i+1}}{2}}\right) \\ (2.8) \quad &\leq \frac{1}{\ln b - \ln a} \int_a^b \frac{f(s)}{s} ds \\ &\leq \sum_{i=0}^{n-1} (\lambda_{i+1} - \lambda_i) \frac{f(a^{1-\lambda_i} b^{\lambda_i}) + f(a^{1-\lambda_{i+1}} b^{\lambda_{i+1}})}{2} \\ &\leq \frac{1}{2} [f(a) + f(b)], \end{aligned}$$

for any division $0 = \lambda_0 < \lambda_1 < \dots < \lambda_{n-1} < \lambda_n = 1$ with $n \geq 1$ of the interval $[0, 1]$.

If we write the inequality (2.8) for $0 < \lambda < 1$, then we get

$$\begin{aligned} f(\sqrt{ab}) &\leq (1-\lambda) f\left(a^{\frac{1-\lambda}{2}} b^{\frac{\lambda+1}{2}}\right) + \lambda f\left(a^{\frac{2-\lambda}{2}} b^{\frac{\lambda}{2}}\right) \\ (2.9) \quad &\leq \frac{1}{\ln b - \ln a} \int_a^b \frac{f(t)}{t} dt \\ &\leq \frac{1}{2} \left[f\left(a^{1-\lambda} b^{\lambda}\right) + (1-\lambda) f(b) + \lambda f(a) \right] \\ &\leq \frac{f(a) + f(b)}{2}. \end{aligned}$$

The inequality (2.9) was obtained in [10].

In the following section we establish some weighted Hermite–Hadamard type inequalities for GA -convex functions.

3. Weighted inequalities. We have the following weighted inequality:

Theorem 4. Let $f : [a, b] \subset (0, \infty) \rightarrow \mathbb{R}$ be a GA -convex function on $[a, b]$ and $w : [a, b] \rightarrow [0, \infty)$ an integrable function on $[a, b]$ with $\int_a^b w(t) dt > 0$, then

$$\begin{aligned} (3.1) \quad f\left(\exp\left(\frac{\int_a^b w(t) \ln t dt}{\int_a^b w(t) dt}\right)\right) &\leq \frac{\int_a^b w(t) f(t) dt}{\int_a^b w(t) dt} \\ &\leq \frac{\left(\ln b - \frac{\int_a^b w(t) \ln t dt}{\int_a^b w(t) dt}\right) f(a) + \left(\frac{\int_a^b w(t) \ln t dt}{\int_a^b w(t) dt} - \ln a\right) f(b)}{\ln b - \ln a}. \end{aligned}$$

Proof. Observe that for $t \in [a, b]$ we have

$$\ln t = \frac{(\ln b - \ln t) \ln a + (\ln t - \ln a) \ln b}{\ln b - \ln a}.$$

By the convexity of $f \circ \exp$ we have

$$\begin{aligned} (3.2) \quad f(t) &= (f \circ \exp)(\ln t) \\ &= (f \circ \exp) \left(\frac{(\ln b - \ln t) \ln a + (\ln t - \ln a) \ln b}{\ln b - \ln a} \right) \\ &\leq \frac{\ln b - \ln t}{\ln b - \ln a} f(a) + \frac{\ln t - \ln a}{\ln b - \ln a} f(b) \end{aligned}$$

for any $t \in [a, b]$.

If we multiply (3.2) by $w(t) \geq 0$, $t \in [a, b]$ and integrate over t on $[a, b]$, then we get

$$\begin{aligned} &\int_a^b f(t) w(t) dt \\ &\leq \frac{\ln b \int_a^b w(t) dt - \int_a^b w(t) \ln t dt}{\ln b - \ln a} f(a) + \frac{\int_a^b w(t) \ln t dt - \ln a \int_a^b w(t) dt}{\ln b - \ln a} f(b) \end{aligned}$$

and the second inequality in (3.1) is proved.

By Jensen's inequality we have

$$\frac{\int_a^b w(t) f(t) dt}{\int_a^b w(t) dt} = \frac{\int_a^b w(t) (f \circ \exp)(\ln t) dt}{\int_a^b w(t) dt} \geq (f \circ \exp) \left(\frac{\int_a^b w(t) \ln t dt}{\int_a^b w(t) dt} \right)$$

and the first part of (3.1) is proved. \square

Corollary 2 (see Theorem 2). *Let $f : [a, b] \subset (0, \infty) \rightarrow \mathbb{R}$ be a GA-convex function on $[a, b]$, then*

$$(3.3) \quad f(\sqrt{ab}) \leq \frac{1}{\ln b - \ln a} \int_a^b \frac{f(t)}{t} dt \leq \frac{f(a) + f(b)}{2}.$$

Proof. If we take $w(t) = \frac{1}{t}$ in (3.1), then we have

$$\begin{aligned} (3.4) \quad &f \left(\exp \left(\frac{\int_a^b \frac{\ln t}{t} dt}{\int_a^b \frac{1}{t} dt} \right) \right) \leq \frac{\int_a^b \frac{f(t)}{t} dt}{\int_a^b \frac{1}{t} dt} \\ &\leq \frac{\left(\ln b - \frac{\int_a^b \frac{\ln t}{t} dt}{\int_a^b \frac{1}{t} dt} \right) f(a) + \left(\frac{\int_a^b \frac{\ln t}{t} dt}{\int_a^b \frac{1}{t} dt} - \ln a \right) f(b)}{\ln b - \ln a}. \end{aligned}$$

Since

$$\int_a^b \frac{\ln t}{t} dt = \frac{1}{2} \left[(\ln b)^2 - (\ln a)^2 \right], \quad \int_a^b \frac{1}{t} dt = \ln b - \ln a,$$

then we get from (3.4)

$$\begin{aligned} f\left(\exp\left(\frac{\ln b + \ln a}{2}\right)\right) &\leq \frac{\int_a^b \frac{f(t)}{t} dt}{\ln b - \ln a} \\ &\leq \frac{(\ln b - \frac{\ln b + \ln a}{2}) f(a) + (\frac{\ln b + \ln a}{2} - \ln a) f(b)}{\ln b - \ln a} \end{aligned}$$

and the inequality (3.3) is proved. \square

Define the p -logarithmic mean as

$$L_p(a, b) := \left(\frac{1}{p+1} \frac{b^{p+1} - a^{p+1}}{b - a} \right)^{1/p}, \quad p \neq -1, 0$$

for $0 < a < b$.

Corollary 3. Let $f : [a, b] \subset (0, \infty) \rightarrow \mathbb{R}$ be a GA-convex function on $[a, b]$, then for any $p \in \mathbb{R}$ with $p \neq 0, -1$ we have

$$\begin{aligned} (3.5) \quad &L_p^p(a, b) f\left([I(a^{p+1}, b^{p+1})]^{\frac{1}{p+1}}\right) \\ &\leq \frac{1}{b-a} \int_a^b t^p f(t) dt \\ &\leq \frac{(L(a^{p+1}, b^{p+1}) - a^{p+1}) f(a) + (b^{p+1} - L(a^{p+1}, b^{p+1})) f(b)}{(p+1)(b-a)}. \end{aligned}$$

If $p = 0$, then we have

$$\begin{aligned} (3.6) \quad &f(I(a, b)) \leq \frac{1}{b-a} \int_a^b f(t) dt \\ &\leq \frac{(L(a, b) - a) f(a) + (b - L(a, b)) f(b)}{b-a}. \end{aligned}$$

Proof. If we take $w(t) = t^p$, with $p \neq 0, -1$ in (3.1), then we have

$$\begin{aligned} (3.7) \quad &f\left(\exp\left(\frac{\int_a^b t^p \ln t dt}{\int_a^b t^p dt}\right)\right) \\ &\leq \frac{\int_a^b t^p f(t) dt}{\int_a^b t^p dt} \\ &\leq \frac{\left(\ln b - \frac{\int_a^b t^p \ln t dt}{\int_a^b t^p dt}\right) f(a) + \left(\frac{\int_a^b t^p \ln t dt}{\int_a^b t^p dt} - \ln a\right) f(b)}{\ln b - \ln a}. \end{aligned}$$

We have

$$\int_a^b t^p dt = \frac{1}{p+1} (b^{p+1} - a^{p+1})$$

and

$$\begin{aligned} \int_a^b t^p \ln t dt &= \frac{1}{p+1} \int_a^b \ln t dt^{p+1} \\ &= \frac{1}{p+1} \left[t^{p+1} \ln t \Big|_a^b - \int_a^b t^p dt \right] \\ &= \frac{1}{p+1} \left(b^{p+1} \ln b - a^{p+1} \ln a - \frac{1}{p+1} (b^{p+1} - a^{p+1}) \right). \end{aligned}$$

Then

$$\begin{aligned} \frac{\int_a^b t^p \ln t dt}{\int_a^b t^p dt} &= \frac{\frac{1}{p+1} \left(b^{p+1} \ln b - a^{p+1} \ln a - \frac{1}{p+1} (b^{p+1} - a^{p+1}) \right)}{\frac{1}{p+1} (b^{p+1} - a^{p+1})} \\ &= \frac{1}{p+1} \ln I(a^{p+1}, b^{p+1}) = \ln [I(a^{p+1}, b^{p+1})]^{\frac{1}{p+1}}, \end{aligned}$$

$$\begin{aligned} \ln b - \frac{\int_a^b t^p \ln t dt}{\int_a^b t^p dt} &= \ln b - \frac{1}{p+1} \left[\frac{b^{p+1} \ln b^{p+1} - a^{p+1} \ln a^{p+1}}{b^{p+1} - a^{p+1}} - 1 \right] \\ &= \frac{1}{p+1} \left(1 - \frac{\ln b^{p+1} - \ln a^{p+1}}{b^{p+1} - a^{p+1}} a^{p+1} \right) = \frac{1}{p+1} \left(1 - \frac{a^{p+1}}{L(a^{p+1}, b^{p+1})} \right) \\ &= \frac{1}{p+1} \frac{L(a^{p+1}, b^{p+1}) - a^{p+1}}{L(a^{p+1}, b^{p+1})} \end{aligned}$$

and, similarly,

$$\frac{\int_a^b t^p \ln t dt}{\int_a^b t^p dt} - \ln a = \frac{1}{p+1} \frac{b^{p+1} - L(a^{p+1}, b^{p+1})}{L(a^{p+1}, b^{p+1})}.$$

From (3.7) we then have

$$\begin{aligned} &f \left(\exp \left(\ln [I(a^{p+1}, b^{p+1})]^{\frac{1}{p+1}} \right) \right) \\ &\leq \frac{\frac{1}{b-a} \int_a^b t^p f(t) dt}{\frac{1}{b-a} \int_a^b t^p dt} \\ &\leq \frac{\left(\frac{1}{p+1} \frac{L(a^{p+1}, b^{p+1}) - a^{p+1}}{L(a^{p+1}, b^{p+1})} \right) f(a) + \left(\frac{1}{p+1} \frac{b^{p+1} - L(a^{p+1}, b^{p+1})}{L(a^{p+1}, b^{p+1})} \right) f(b)}{\ln b - \ln a}, \end{aligned}$$

i.e.

$$\begin{aligned}
 & f\left([I(a^{p+1}, b^{p+1})]^{\frac{1}{p+1}}\right) \\
 & \leq \frac{\frac{1}{b-a} \int_a^b t^p f(t) dt}{L_p^p(a, b)} \\
 & \leq \frac{\left(\frac{1}{p+1} \frac{L(a^{p+1}, b^{p+1}) - a^{p+1}}{L(a^{p+1}, b^{p+1})}\right) f(a) + \left(\frac{1}{p+1} \frac{b^{p+1} - L(a^{p+1}, b^{p+1})}{L(a^{p+1}, b^{p+1})}\right) f(b)}{\ln b - \ln a} \\
 & = \frac{\frac{L(a^{p+1}, b^{p+1}) - a^{p+1}}{L(a^{p+1}, b^{p+1})} f(a) + \frac{b^{p+1} - L(a^{p+1}, b^{p+1})}{L(a^{p+1}, b^{p+1})} f(b)}{\ln b^{p+1} - \ln a^{p+1}}.
 \end{aligned}$$

By multiplying this inequality with $L_p^p(a, b)$, we get

$$\begin{aligned}
 & L_p^p(a, b) f\left([I(a^{p+1}, b^{p+1})]^{\frac{1}{p+1}}\right) \\
 & \leq \frac{1}{b-a} \int_a^b t^p f(t) dt \\
 & \leq \frac{b^{p+1} - a^{p+1}}{(p+1)(b-a)} \frac{\frac{L(a^{p+1}, b^{p+1}) - a^{p+1}}{L(a^{p+1}, b^{p+1})} f(a) + \frac{b^{p+1} - L(a^{p+1}, b^{p+1})}{L(a^{p+1}, b^{p+1})} f(b)}{\ln b^{p+1} - \ln a^{p+1}} \\
 & = \frac{(L(a^{p+1}, b^{p+1}) - a^{p+1}) f(a) + (b^{p+1} - L(a^{p+1}, b^{p+1})) f(b)}{(p+1)(b-a)}.
 \end{aligned}$$

If we perform the calculations in the above inequalities for $p = 0$, we get the desired inequality (3.6). We omit the details. \square

Remark 2. If we take $p = 1$ in (3.5), then we get

$$\begin{aligned}
 (3.8) \quad & f\left(\sqrt{I(a^2, b^2)}\right) \leq \frac{1}{b-a} \int_a^b t f(t) dt \\
 & \leq \frac{(A(a, b) L(a, b) - a^2) f(a) + (b^2 - A(a, b) L(a, b)) f(b)}{2(b-a)}.
 \end{aligned}$$

4. Applications. Let $q \neq 0$ and consider the convex function $g(t) = \exp(qt)$, $t \in \mathbb{R}$. Then the function $f_q : (0, \infty) \rightarrow \mathbb{R}$, $f_q(t) = g(\ln t) = \exp(q \ln t) = t^q$ is a GA -convex function on $(0, \infty)$. We observe that for $0 < a < b$ we have

$$\begin{aligned}
 \frac{1}{b-a} \int_a^b t^q dt & = \begin{cases} \frac{1}{q+1} \frac{b^{q+1} - a^{q+1}}{b-a}, & q \neq -1 \\ \frac{\ln b - \ln a}{b-a}, & q = -1 \end{cases} \\
 & = \begin{cases} L_q^q(a, b), & q \neq -1 \\ L^{-1}(a, b), & q = -1, \end{cases}
 \end{aligned}$$

where $L_q(a, b)$ ($q \neq -1$) is the q -Logarithmic mean and L is the logarithmic mean defined in the introduction.

If we write the inequality (2.8) for the GA -convex function on $(0, \infty)$, $f_q : (0, \infty) \rightarrow \mathbb{R}$, $f_q(t) = t^q$, $q \neq 0$, then we get the following result:

Proposition 1. For any division $0 = \lambda_0 < \lambda_1 < \dots < \lambda_{n-1} < \lambda_n = 1$ with $n \geq 1$ of the interval $[0, 1]$,

$$\begin{aligned}
 G^q(a, b) &\leq \sum_{i=0}^{n-1} (\lambda_{i+1} - \lambda_i) G^q(a^{2-\lambda_i-\lambda_{i+1}}, b^{\lambda_i+\lambda_{i+1}}) \\
 (4.1) \quad &\leq L(a, b) L_{q-1}^{q-1}(a, b) \\
 &\leq \sum_{i=0}^{n-1} (\lambda_{i+1} - \lambda_i) \frac{G^q(a^{2(1-\lambda_i)}, b^{2\lambda_i}) + G^q(a^{2(1-\lambda_{i+1})}, b^{2\lambda_{i+1}})}{2} \\
 &\leq A(a^q, b^q).
 \end{aligned}$$

If we write the inequality (3.5) for the GA -convex function on $(0, \infty)$, $f_q : (0, \infty) \rightarrow \mathbb{R}$, $f_q(t) = t^q$, $q \neq 0$, then:

Proposition 2. If $0 < a < b$ and $p, q \in \mathbb{R}$ with $p \neq 0, -1$, $q \neq 0$, then

$$\begin{aligned}
 (4.2) \quad &L_p^p(a, b) [I(a^{p+1}, b^{p+1})]^{\frac{q}{p+1}} \\
 &\leq L_{p+q}^{p+q}(a, b) \\
 &\leq \frac{(L(a^{p+1}, b^{p+1}) - a^{p+1}) a^q + (b^{p+1} - L(a^{p+1}, b^{p+1})) b^q}{(p+1)(b-a)}.
 \end{aligned}$$

From (3.6) and (3.8) we get:

Proposition 3. If $0 < a < b$ and $q \in \mathbb{R}$ with $q \neq 0$, then

$$(4.3) \quad I^q(a, b) \leq L_q^q(a, b) \leq \frac{(L(a, b) - a) a^q + (b - L(a, b)) b^q}{b - a},$$

and

$$\begin{aligned}
 (4.4) \quad &I^{q/2}(a^2, b^2) \leq L_{q+1}^{q+1}(a, b) \\
 &\leq \frac{(A(a, b) L(a, b) - a^2) a^q + (b^2 - A(a, b) L(a, b)) b^q}{2(b-a)}.
 \end{aligned}$$

Acknowledgement. The author would like to thank the anonymous referee for valuable suggestions that have been implemented in the final version of the manuscript.

REFERENCES

- [1] Alomari, M., Darus, M., *The Hadamard's inequality for s -convex function*, Int. J. Math. Anal. (Ruse) **2** (2008), no. 13–16, 639–646.

-
- [2] Anderson, G. D., Vamanamurthy, M. K., Vuorinen, M., *Generalized convexity and inequalities*, J. Math. Anal. Appl. **335** (2007) 1294–1308.
- [3] Beckenbach, E. F., *Convex functions*, Bull. Amer. Math. Soc. **54** (1948), 439–460.
- [4] Cristescu, G., *Hadamard type inequalities for convolution of h -convex functions*, Ann. Tiberiu Popoviciu Semin. Funct. Equ. Approx. Convexity **8** (2010), 3–11.
- [5] Dragomir, S. S., *Some remarks on Hadamard’s inequalities for convex functions*, Extracta Math. **9** (2) (1994), 88–94.
- [6] Dragomir, S. S., *An inequality improving the first Hermite–Hadamard inequality for convex functions defined on linear spaces and applications for semi-inner products*, J. Inequal. Pure Appl. Math. **3** (2) (2002), Article 31, 8 pp.
- [7] Dragomir, S. S., *An inequality improving the second Hermite–Hadamard inequality for convex functions defined on linear spaces and applications for semi-inner products*, J. Inequal. Pure Appl. Math. **3** (3) (2002), Article 35, 18 pp.
- [8] Dragomir, S. S., *An Ostrowski like inequality for convex functions and applications*, Revista Math. Complutense **16** (2) (2003), 373–382.
- [9] Dragomir, S. S., *Operator Inequalities of Ostrowski and Trapezoidal Type*, Springer, New York, 2012.
- [10] Dragomir, S. S., *Inequalities of Hermite–Hadamard type for GA-convex functions*, Preprint RGMIA Res. Rep. Coll.
- [11] Dragomir, S. S., Cerone, P., Roumeliotis J., Wang, S., *A weighted version of Ostrowski inequality for mappings of Hölder type and applications in numerical analysis*, Bull. Math. Soc. Sci. Math. Roumanie **42** (90) (4) (1999), 301–314.
- [12] Dragomir, S. S., Fitzpatrick, S., *The Hadamard inequalities for s -convex functions in the second sense*, Demonstratio Math. **32** (4) (1999), 687–696.
- [13] Dragomir, S. S., Fitzpatrick, S., *The Jensen inequality for s -Breckner convex functions in linear spaces*, Demonstratio Math. **33** (1) (2000), 43–49.
- [14] Dragomir, S. S., Mond, B., *On Hadamard’s inequality for a class of functions of Godunova and Levin*, Indian J. Math. **39** (1) (1997), 1–9.
- [15] Dragomir, S. S., Pearce, C. E. M., *On Jensen’s inequality for a class of functions of Godunova and Levin*, Period. Math. Hungar. **33** (2) (1996), 93–100.
- [16] Dragomir, S. S., Pearce, C. E. M., *Quasi-convex functions and Hadamard’s inequality*, Bull. Austral. Math. Soc. **57** (1998), 377–385.
- [17] Dragomir, S. S., Pearce, C. E. M., *Selected Topics on Hermite–Hadamard Inequalities and Applications*, RGMIA Monographs, Victoria University, 2000 [Online http://rgmia.org/monographs/hermite_hadamard.html].
- [18] Dragomir, S. S., Pečarić, J., Persson, L., *Some inequalities of Hadamard type*, Soochow J. Math. **21** (3) (1995), 335–341.
- [19] Dragomir, S. S., Rassias, Th. M., (Eds), *Ostrowski Type Inequalities and Applications in Numerical Integration*, Kluwer Academic Publisher, 2002.
- [20] El Farissi, A., *Simple proof and refinement of Hermite–Hadamard inequality*, J. Math. Ineq. **4** (3) (2010), 365–369.
- [21] Kirmaci, U. S., Klaričić Bakula, M., Özdemir, M. E., Pečarić, J., *Hadamard-type inequalities for s -convex functions*, Appl. Math. Comput. **193** (1) (2007), 26–35.
- [22] Latif, M. A., *On some inequalities for h -convex functions*, Int. J. Math. Anal. (Ruse) **4** (29–32) (2010), 1473–1482.
- [23] Mitrinović, D. S., Lacković, I. B., *Hermite and convexity*, Aequationes Math. **28** (1985), 229–232.
- [24] Mitrinović, D. S., Pečarić, J. E., *Note on a class of functions of Godunova and Levin*, C. R. Math. Rep. Acad. Sci. Canada **12** (1) (1990), 33–36.
- [25] Noor, M. A., Noor, K. L., Awan, M. U., *Some inequalities for geometrically-arithmetically h -convex functions*, Creat. Math. Inform. **23** (1) (2014), 91–98.

- [26] Pearce, C. E. M., Rubinov, A. M., *P-functions, quasi-convex functions, and Hadamard-type inequalities*, J. Math. Anal. Appl. **240** (1) (1999), 92–104.
- [27] Pečarić, J. E., Dragomir, S. S., *On an inequality of Godunova–Levin and some refinements of Jensen integral inequality*, Itinerant Seminar on Functional Equations, Approximation and Convexity (Cluj-Napoca, 1989), 263–268, Preprint, 89-6, Univ. “Babeş-Bolyai”, Cluj-Napoca, 1989.
- [28] Zhang, X.-M., Chu, Y.-M., Zhang, X.-H., *The Hermite–Hadamard type inequality of GA-convex functions and its application*, J. Inequal. Appl. vol. **2010**, Article 507560, 11 pp.

S. S. Dragomir
College of Engineering & Science
Victoria University, PO Box 14428
Melbourne City, MC 8001
Australia
e-mail: sever.dragomir@vu.edu.au
url: <http://rgmia.org/dragomir>

DST-NRF Centre of Excellence in the Mathematical and Statistical Sciences
School of Computer Science and Applied Mathematics
University of the Witwatersrand
Private Bag 3, Johannesburg 2050
South Africa

Received April 11, 2017