



VICTORIA UNIVERSITY
MELBOURNE AUSTRALIA

The Complex Interaction between the Tumor Micro-Environment and Immune Checkpoints in Breast Cancer

This is the Published version of the following publication

Barriga, Vanessa, Kuol, Nyanbol, Nurgali, Kulmira and Apostolopoulos, Vasso
(2019) The Complex Interaction between the Tumor Micro-Environment and
Immune Checkpoints in Breast Cancer. *Cancers*, 11 (8). ISSN 2072-6694

The publisher's official version can be found at
<https://www.mdpi.com/2072-6694/11/8/1205>

Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/39536/>

Article

The Complex Interaction between the Tumor Micro-Environment and Immune Checkpoints in Breast Cancer

Vanessa Barriga ^{1,2}, Nyanbol Kuol ², Kulmira Nurgali ^{2,†} and Vasso Apostolopoulos ^{2,*,†} 

¹ College of Health and Biomedicine, Victoria University, Melbourne 3030, Australia

² Institute for Health and Sport, Victoria University, Melbourne 3030, Australia

* Correspondence: vasso.apostolopoulos@vu.edu.au; Tel.: +61-3-9919-2025

† These authors contributed equally.

Received: 14 July 2019; Accepted: 12 August 2019; Published: 19 August 2019



Abstract: The progression of breast cancer and its association with clinical outcome and treatment remain largely unexplored. Accumulating data has highlighted the interaction between cells of the immune system and the tumor microenvironment in cancer progression, and although studies have identified multiple facets of cancer progression within the development of the tumor microenvironment (TME) and its constituents, there is lack of research into the associations between breast cancer subtype and staging. Current literature has provided insight into the cells and pathways associated with breast cancer progression through expression analysis. However, there is lack of co-expression studies between immune pathways and cells of the TME that form pro-tumorigenic relationships contributing to immune-evasion. We focus on the immune checkpoint and TME elements that influence cancer progression, particularly studies in molecular subtypes of breast cancer.

Keywords: breast cancer; checkpoint; checkpoint inhibitors; tumor microenvironment; tumor-associated macrophages; cancer-associated fibroblasts; tumor-associated neutrophils; IDO; PD-L1; Siglec-9

1. Introduction

Breast carcinoma remains a complex heterogeneous disorder, being the most frequent malignant disease in women worldwide, leading to premature death in women in developed countries [1]. In fact, the higher developed countries such as Europe, Australia, and the USA have the highest incidence of breast cancer compared to those from mid-developed countries including, Latin America, central Europe, and the Caribbean; the lowest incidence being in the low developed countries such as Asia and Africa [2]. In the last decade, the mortality rate of breast cancer has declined as a result of early detection and effective treatments [3,4], although, 20% of women diagnosed with breast cancer will still die from the disease [5]. One of the challenges of breast cancer research has been the characterization of the molecular changes that are associated with breast cancer progression [6]. The diagnosis of breast cancer is based on three key parameters including histological type, tumor grade, and tumor stage. Within this, there are at least 20 different histological subtypes of breast cancer, which display differences in morphology and growth patterns [7].

The main biological markers utilized in the diagnosis and treatment of breast cancer include the overexpression of estrogen receptor (ER), progesterone receptor (PR), and human epidermal growth factor receptor 2 (HER2). Based on these clinically used markers, 5 major molecular intrinsic subtypes of breast cancer have been identified: (i) luminal A (histological phenotype: ER+, PR+, HER2−, Ki67−), (ii) luminal B (ER+, PR+, HER2+/, Ki67+), (iii) HER2 enriched (ER−, PR−, HER2+),

(iv) basal-like subtype (ER−, PR−, HER2−, triple negative; associated with poor prognosis and early onset of metastasis) and (v) claudin-low tumor subtype which is commonly classified as basal-like (or normal breast-like) [8–11]. In addition to these established subtypes, breast cancer staging refers to the extent of disease and is a critical factor in the prognostic outcomes of the patients as well as for oncologists in making treatment decisions. Staging is categorized by stages 0–IV which is determined by tumor size, invasion of tumor cells into the chest wall and lymph nodes, and distant metastasis.

The proliferation of tumor cells and metastasis is influenced by immune cells present within the tumor microenvironment (TME) and their mechanisms of actions. As such, lymphocytes, fibroblasts, neutrophils, eosinophils, macrophages and myeloid-derived suppressor cells are present within the TME with altered mechanisms and influence the outcome of cancer cell growth and migration. In addition, immune-checkpoints have been identified and can be expressed on both immune cells and tumor cells. A number of negative regulatory mechanisms can inhibit the anti-tumor immune response through the expression of these immune checkpoints. An ongoing immune response can be represented by the upregulation of immune checkpoints (CTLA4, PD-1/PD-L1, IDO and members of the CD33-like siglecs i.e., siglec-7 and siglec-9, but their expression on tumor cells can also be driven by oncogenic pathways. By activating these inhibitory pathways, cancer cells are able to escape from immune cell attack by causing apoptosis of tumor-specific T cells [12–16]. Siglec-7 inhibits NK cell lysis [17] and the mucin MUC1 modulates the tumor immunological microenvironment through engagement of the lectin Siglec-9 [18]. Development of immune checkpoint inhibitors, such as anti-CTLA-4, anti-PD-1 and anti-PD-L1, bypass the immune checkpoint, with the aim of rescuing and enhancing the function of anti-tumor T cells [19].

Immunotherapy or vaccine studies to date have primarily focused on the stimulation of CD8+ T cells and/or T helper (Th)-1 pro-inflammatory responses to biological targets expressed by tumor cells [20]. For example, the high molecular weight tumor-associated glycoprotein, MUC1 (CD227) [21–25], has been used as a target for cancer immunotherapy studies in both mice [26–43] and in human clinical trials [44–46]. In particular, human clinical studies of MUC1 conjugated to the carrier mannan has been in phase I, II, III trials, and in early stage breast cancer patients these conjugates show protection against recurrence up to 18 years post-vaccination clinical follow-up [47–50]. Although showing efficacy in early stage breast cancer patients, in other clinical studies in advanced cancer patients clinical outcomes were only modest. In an attempt to improve clinical outcomes, checkpoint inhibitors are being used in combination with vaccines/immunotherapeutics. In addition, it is important to understand the role of other immune cell infiltrates within the TME, which will aid in the design of new improved cancer immunotherapeutics. Herein, we focus on immune cells within the TME and immune-checkpoints associated with breast cancer progression.

2. Tumor Microenvironment

Advances in the understanding of the TME in pre-invasive and invasive breast cancer have demonstrated strong evidence to suggest that the TME and associated molecules as well as the infiltration of immune cells, soluble factors and altered extracellular matrix, are involved in promoting tumor growth and metastasis [51,52]. These critical elements of the TME have initiated the identification of new breast cancer markers, such as biological, immunological and immunosuppressive which are associated with tumor development and progression [53]. More recently, there are implications that the nervous system also modulates immune responses in suppressing tumor growth, or, enhances tumors ability to evade immune surveillance leading to metastasis [12–14,54]. Over the past decades of cancer research, focus has predominately fallen into the complex interactions of the TME. Its contribution to cancer progression relies heavily on the residents within the TME including, tumor-associated macrophages (TAMs), cancer-associated fibroblasts (CAF), endothelial cells, pericytes, leukocytes, tumor-infiltrating lymphocytes (TILs), cytokine milieu and the extracellular matrix (ECM) (Figure 1, Table 1). It is important to note that tumor cells can efficiently recruit stromal cells (e.g., fibroblasts), immune cells and vascular cells by secreting growth factors, cytokines and chemokines. These cells

build the microenvironment by releasing growth promoting signals and intermediate metabolites as well as remodel tissue structure. The signaling between the cancer cells and the TME stimulates or inhibits proliferation and metastatic capability [55–58]. The tumor not only manages to escape from the host immune system, but it effectively benefits from infiltrating cells by modifying their functions to create the microenvironment favorable to tumor progression [59].

2.1. Epithelial-Mesenchymal Transition

Epithelial-mesenchymal transition (EMT) is the ability of cancer cells to leave their epithelial state and acquire characteristics of mesenchymal cells. EMT plays a fundamental role in tumor progression and metastasis and involves a number of signaling pathways such as transforming growth factor beta (TGF β), nuclear factor- κ B (NF- κ B), Wnt, Notch and others [60]. EMT markers such as *Snail*, *Twist-1* and *Lox* are transcription factors that induce EMT which have also been shown to be expressed by human breast tumors [61,62]. *Snail* is known to repress *E-cadherin* (a key molecule involved in epithelial adhesion) expression, a hallmark of EMT, and has been linked to the migratory and invasive phenotype of breast cancer cells [63]. *Twist* is a helix loop helix protein involved in the downregulation of the epithelial genes such as *E-cadherin*, *claudins*, *occludins* and stimulates a number of the mesenchymal genes including, *N-cadherin* and *fibronectin* [64]. *Lox* is involved in remodeling the extracellular matrix facilitating invasion and metastasis [65]. In addition, circulatory tumor cells and highly enriched for characteristics of mesenchymal cells showing loss of *E-cadherin* and gain of *N-cadherin* [66,67]. Studies show that CXC chemokine ligand 8 (CXCL8 or IL-8) promotes the EMT of human breast cancer cells via the formation of the *TWIST1-p65* complex that activates transcription of NF- κ B and increases the binding affinity of p65 to CXCL8 [68]. Studies of EMT markers in triple negative breast cancer (TNBC) and non-TNBC subtypes showed the overexpression of *Lox* to be significantly higher in TNBC subtype compared to non-TNBC subtypes. However, the expressions of *Snail* and *Twist* showed no difference between TNBC and non-TNBC subtypes [62]. Although the expression of *Lox* cannot be linked to prognosis it may be a possible target for future TNBC systemic therapies (Figure 1, Table 1). In addition, immune infiltrates in TNBC is associated with good prognosis although this is not the case for ER+ tumors.




Figure 1. Tumor-associated immune cells in the tumor microenvironment (TME) of breast cancer models.

Within the TME there is an array of resident cells contributing to the progression and metastasis of breast cancer cells. The different residents and their associated secretory elements including stimulatory growth factors, chemokines and cytokines are shown. The expression of these residents within the TME of breast cancer patients may aid in discovering new markers associated with specific subtype leading to earlier diagnosis and better clinical outcome. [ARG1. Arginase 1; CAF. Cancer-associated Fibroblast; CD163. Macrophage scavenger receptor; CCL2; Chemokine Ligand 2; CXCL8-CXCR1/2. Chemokine Ligand 8-Chemokine Receptor 1 & 2; CXCL12. Chemokine Ligand 12; COX-2. cyclooxygenase-2; EMT. Epithelial-Mesenchymal Transition; ER– Estrogen Receptor Negative; FGF. Fibroblast Growth Factor; GM-CSF. Granulocyte-Macrophage Colony-Stimulating Factor; HGF. Hepatocyte Growth Factor; IDO. Indoleamine-2, 3-Dioxygenase; IL. Interleukin; iDC. immature Dendritic Cells; mDC. mature Dendritic Cells; MDSCs. Myeloid-derived suppressor cells; M2 TAMs. M2 subtype Tumor-associated Macrophage; NK CELL. Natural Killer; N2 TAN. N2 Subtype Tumor-associated Neutrophil; PDGF-R. Platelet-Derived Growth Factor Receptor; PD-1. Programmed cell death protein 1; PGE₂. Prostaglandin E₂; ROS. Reactive Oxygen Species; SDF-1. Stromal cell-derived factor-1; TGF- β . Transforming Growth Factor-beta; Th Cells. T-helper cells; TILs. Tumor-Infiltrating Lymphocytes; TREG. T-regulatory cells; VEGF. Vascular endothelial growth factor].

2.2. Tumor-Infiltrating Lymphocytes

Immune cells present within the tumor include those mediating adaptive immune responses, such as T cells, dendritic cells and B cells, as well as effectors of the innate immune responses, i.e., macrophages, neutrophils, eosinophils and natural killer cells [69]. In fact, premalignant ductal breast carcinoma in situ shows an increase in lymphocyte infiltration, with predominant cells being, activated T cells, B cells, and, the immune suppressive regulatory T cells (Treg; CD4+CD25+ Forkhead box protein 3 (Foxp3)+ cells) [70]. Tumor-infiltrating lymphocytes (TILs) are largely CD8+ T cells, CD4+ T helper (Th) cells and Treg cells [69,71]. Among CD4+ T cells present in the tumor, a subset of CD4+CD25^{high}Foxp3+ Treg cells are able to suppress proliferation of other T cells within the microenvironment through contact-dependent mechanisms, or anti-inflammatory cytokine (IL-10 and TGF β) secretion [72,73]. Cytotoxic CD8+ T-cells have the ability to kill cancer cells via secretion of pro-inflammatory interferon- γ (IFN γ) and granzyme-perforin complex. In addition, the activation and maturation of CD8+ T-cells is also modulated by IFN γ secreted by CD4+ Th1 cells and specific tumor-associated antigens processed by dendritic cells [74]. In relation to breast cancer subtypes, ER– cancers show higher number of TILs compared to ER+ cancers [75]. In addition, the overall number of T cells, B cells, macrophages and myeloid-derived suppressor cells (MDSC) are also higher in ER– compared to ER+ breast cancers [75]. Thus, the number of TILs within the tumor microenvironment may not only help distinguish subtype but can be used to identify markers associated with progression and metastasis from the primary tumor (Figure 1, Table 1).

2.3. Cancer-Associated Fibroblasts

Fibroblast cells are important contributors to the development of the extracellular matrix and they secrete a number of factors including collagen and cytokines, which aids in the structural framework of the extracellular matrix [76]. Within the TME and surrounding space, accumulation of different immune and regulatory cells may stimulate or inhibit tumor growth (Figure 1, Table 1). Fibroblasts represent the majority of stromal cells within the TME. Commonly, activated fibroblasts inhibit early stages of tumor progression through production of fibroblast factors and IL-6 occurring in gap junctions [77]. However, epithelial cells, endothelial cells, and cancer cells have been associated with altering fibroblasts into cancer-associated fibroblasts (CAFs) [78]. CAFs have shown to secrete various growth factors and cytokines associated with promoting breast cancer proliferation and metastasis [79]. In fact, fibroblast growth factor, human growth factor, tenascin, thrombospondin-1, TGF β and stromal cell-derived factor 1 (SDF-1 or CXCL12), are found in cancer cells including breast cancer, at sites of chronic inflammation produced by CAFs [80–82]. The secretion of high levels of TGF β by tumor cells causes migration of fibroblasts to the TME initiating trans-differentiation of fibroblasts to CAFs [83]. Platelet-derived growth

factor can also indirectly recruit myofibroblasts by stimulating TGF β secretion from macrophages [84]. Thus, CAFs have the potential to promote growth and angiogenesis, remodel the ECM, and direct cell-to-cell interaction. Furthermore, CAFs have been associated with breast cancer subtype, ER+, TNBC and HER2+ [85]. Gene expression analysis have indicated that CAFs within the tumor stroma may have subtype-specific gene expression profiles, with CAFs increasing the invasive properties of the HER2+ subtype. These studies show the importance of CAFs in cancer progression and their ability to alter the normal function of fibroblasts and their anti-tumorigenic abilities. In addition, further studies are warranted to bridge the gap in knowledge as CAFs may hold potential prognostic value and could be used to distinguish breast cancer subtype and stage.

2.4. Tumor-Associated Macrophages

Macrophage cells are large phagocytic cells found at sites of inflammation and engulf foreign antigens and cancer-associated proteins to stimulate the adaptive immune responses. Macrophages (and dendritic cells) have been the target for cancer immunotherapy and vaccines studies [86–90]. Macrophages also infiltrate the breast cancer microenvironment to initiate an inflammatory response but differentiate into tumor-associated macrophages (TAMs) resembling M2 macrophages (Figure 1). TAMs are re-programmed to inhibit lymphocyte functions through the secretion of inhibitory cytokines including IL-10, prostaglandins and reactive oxygen species (ROS) [91,92]. ROS activates TGF β which inhibits cell growth by arresting cells in the G1 phase of the cell cycle leading to either terminal differentiation or induction of apoptosis [93]. TAMs also produce extracellular matrix-degrading enzymes such as metalloproteinases (MMPs). The proteolytic cleavage of TGF β from latency-associated peptide regulates the secretion, expression and activation of MMP2, MMP3, MMP9 and MMP13, which facilitate tumor cell migration, degradation of blood vessel basal membranes and, metastasis [94,95]. M2 macrophages express arginase 1 (ARG1), anti-inflammatory cytokines, and proteases that support their pro-oncogenic functions [19]. Increased numbers of CD163+ M2 macrophages associate with unfavorable prognosis linked to non-luminal and basal-like breast cancer subtype [96,97]. In addition, vascular endothelial growth factor (VEGF) secretion by TAM stimulate tumor angiogenesis, promoting its invasiveness and metastatic potential [98–100]. TAMs are also able to transform cancer cells to undergo EMT which subsequently enhances the invasion and metastasis of breast cancer cells [101–103]. Similarly, mesenchymal-like breast cancer cells can activate macrophages to a TAM-like phenotype by granulocyte macrophage-colony stimulating factor 1 (GM-CSF1) [101].

TAMs are also associated with the regulation of programmed death receptor-1 (PD-1) and its ligand (L; PD-L1) expression in the tumor microenvironment of TNBC. IFN γ , IL-1 β , tumor TNF α , TGF β , IL-6, and IL-18 are key to the functionality of TAM [104,105]. The expression of PD-L1 on tumor cells might be induced by the secretion of IFN γ via Janus kinase/signal transducer and activator of transcription 3 (JAK/STAT3) and the phosphatidylinositol 3-kinase (PI3K)/protein kinase B (Akt) signaling pathways, which has been shown in lung cancer [106]. TGF β increases the suppressive ability of TAMs by differentiating TAMs into M2 macrophages and upregulation of PD-L1 leads to tumor escape [107]. Tumor-derived IL-18 increases the immunosuppressive properties on NK subsets inducing their PD-1 expression which is correlated with poor prognosis of TNBC patients [108]. This mechanism can lead to inactivation of TILs in the TME and cancer immunotherapy and vaccines studies, and would, therefore, not be effective in such an environment.

2.5. Tumor-Associated Neutrophils

Neutrophils are polymorphonuclear cells whose main function in the innate immune response is the first line of defense and protection against fungal and bacterial infections. Neutrophils are abundantly found in human blood and migrate to a number of tissues from the blood circulation. In recent years it has been noted that neutrophils are also present within the TME with both pro- and anti-tumorigenic properties [109]. In the TME, cytokines initiate polarization of tumor-associated neutrophils (TANs) causing them to differentiate into either pro-inflammatory/anti-tumorigenic (N1

phenotype) or, anti-inflammatory/pro-tumorigenic (N2 phenotype) (Figure 1, Table 1). Their migration from the blood circulation into the TME is stimulated by IL-8 (CXCL8-CXCR1/2 axis) expression by tumor cells [109]. Neutrophils in their normal state do not secrete oncostatin M however, upon interaction with cancer cells oncostatin M becomes highly expressed in TANs. Oncostatin M exhibits an inhibitory effect on cell proliferation of breast cancer cell lines exerting a pro-inflammatory response by inducing adhesion and chemotaxis of neutrophils [110,111]. Conversely, oncostatin M has also been shown to promote tumor progression by enhancing angiogenesis and metastasis to breast cancer cell lines (MDA-MB-23 and T47D) [112]. Neutrophils isolated from healthy human volunteers display no expression of oncostatin M but upon co-culture with MDA-MB-231 or T47D cells, oncostatin M expression is significantly increased [112]. Furthermore, production of GM-CSF1 by breast cancer cells and cell-cell contact are also necessary for neutrophils to release oncostatin M which has also been noted to concurrently increase VEGF and invasive capacity of cell lines [112].

TANs associate with aggressive breast cancer phenotype, facilitates angiogenesis, promotes mutagenesis and suppresses the immune system, leading to poor prognosis of patients [113,114]. The presence of TAN in breast cancer has also been associated with the clinical subtypes of breast cancer indicating that there is a preferential chemotaxis of neutrophils dependent on the subtype [115]. In fact, TANs predominate in TNBC compared to non-TNBC, suggesting that factors associated with the TME may have direct or indirect effects on neutrophil production and chemotaxis into the TME [115]. In addition, TGF β which is highly expressed in TNBC has been shown to be a major contributor to neutrophil chemotaxis, however, TGF β may also induce a pro-tumorigenic N2 phenotype [116–118]. The presence of TANs in the TME of breast cancer patients highlights their involvement in tumor growth and metastasis. However, not much is known regarding their presence and their correlation to the range of breast cancer subtypes, and whether they impact in cancer immunotherapy studies.

2.6. Tumor-Associated Eosinophils

The primary function of eosinophils was believed to be to fight parasitic and bacterial infections, and were involved in the pathogenesis of inflammatory diseases such as allergic asthma and chronic obstructive pulmonary disease with high secretion of IL-5 and other eosinophilia granules [119]. More recently, evidence suggests that eosinophils are abundant in inflammatory bowel disease although their role within the intestinal tissues is not clear [120]. Blood eosinophils migrate to tissues at sites of infection. In the human breast, eosinophils are crucial for mammary gland development where they are interlaced within the terminal end buds of the breast together with a complex stroma of fibroblasts and macrophages; once terminal end buds convert into terminal end ducts, fibroblasts disappear [121,122]. Eosinophils have also been observed within the TME of cancer, including colon, ovarian, prostate and lung carcinomas [123] (Figure 1). Of interest, eosinophils have been noted at the edge of breast cancer biopsy wounds, suggesting that breast cancer biopsies may trigger the recruitment of inflammatory cells including eosinophils [124]. In addition, low eosinophil counts in the peripheral blood of breast cancer patients is a major risk factor of breast cancer recurrence [123]. The presence and amount of mast cells and tumor-associated eosinophils was determined in patients diagnosed with invasive breast cancer of different stages [125]. In that study, it was noted that eosinophils were also not present within the TME. Although a positive correlation of improved prognosis with tumor-associated eosinophils has been shown in a number of solid cancers, in Hodgkin's lymphoma their presence in the TME display a poor prognosis [123]. There is conflicting data on the presence of eosinophils within the TME of breast cancer which requires further studies to understand their role and interaction with other immune cells in the TME and their interaction with cancer cells. The little or lack of eosinophil presence in breast cancers compared to other solid tumors, could be associated with the current methods utilized or different interactions between eosinophils and breast cancer that are yet not defined. Whether eosinophils contribute to immune suppression or immune stimulation in breast cancer immunotherapy studies is also unknown.

2.7. Myeloid-Derived Suppressor Cells

Myeloid-derived suppressor cells (MDSCs) are a diverse population of myeloid progenitor cells which have been shown to play a key role in chronic inflammation and cancer development [126]. In cancer, MDSCs (CD34+CD33+CD13+CD15(−)) promote tumor cell growth and suppresses immune cell function through production of arginase 1 (ARG1) which synergizes with inducible nitric oxide synthase (iNOS) to increase superoxide and nitric oxide (NO) production, thus, reducing lymphocyte function [127,128] (Figure 1, Table 1). Studies on the levels and phenotypes of myeloid cells in peripheral blood and TME of breast cancer patients revealed high levels of tumor-infiltrating myeloid cells, including granulocytes and immature cells lacking expression markers for fully differentiated monocytes or granulocytes [129]. However, the expansion was not reflected in the peripheral blood or in samples from non-breast cancer patients. In addition, the presence of ARG1 expression, important in T cell suppression, reinforces the immunosuppressive ability of MDSCs [129]. Furthermore, MDSCs isolated from breast cancer tissues show high expression of indole amine 2,3 dioxygenase (IDO), an enzyme responsible for the catabolism of tryptophan. This depletion of tryptophan by IDO in the TME produces kynurenine based by-products that lead to inhibition of T cell proliferation and induces T cell apoptosis [130].

3. Immune Checkpoint Molecules and Breast Cancer

Immune-editing involves the process of malignant cell progression based on cancer cell and immune cell interactions in three stages: (i) Elimination: where cancer cells are eliminated following immuno-surveillance involving both innate and adaptive immune cell infiltration including, NK cells, NK-T cells, T cells, and increased pro-inflammatory cytokines in the TME; (ii) Equilibrium: the balance between anti-tumor and tumor-promoting factors in that transformed cells are held in control but are not eliminated following immuno-surveillance; and (iii) Escape: where modifications to tumor cells themselves shape disease progression, as seen in breast cancer, where escape mechanisms include reduced expression of major histocompatibility complex class I [39] and/or co-stimulatory molecules and increased expression of immunosuppressive factors [16]. Immunosuppressive factors involved in the evasion of immune-surveillance play influential roles in cancer progression. Immune checkpoints of inhibitory pathways are crucial for the immune system to maintain self-tolerance and modulate immune responses in order to minimize damage. Immune checkpoints (such as CTLA4, PD-1/PD-L1, IDO, and Siglec-9) are initiated by ligand-receptor interactions and can be blocked by antibodies or modulated by recombinant forms of ligands or receptors [131] (Figure 2, Table 2). It is important to understand immune checkpoint molecules and their role in breast cancer and numerous clinical trials are assessing the efficacy of immune checkpoint inhibitors (via monoclonal antibodies) in cancer therapies either alone or in combination with other cancer therapies.




Figure 2. Immune checkpoints of immunosuppressive actions associated with breast cancer. Immune checkpoints of inhibitory pathways are fundamental in the immune system to maintain self-tolerance and modulate immune responses. In breast cancer, some of these immune checkpoints and immunosuppressive factors have been associated with subtype specificity through their expressions on breast cancer cells. The different immune cells and their ligand-receptor interactions and secreted stimulatory growth factors, chemokines and cytokines are shown. The expression of these immune markers in the TME in breast cancer may or not be subtype-specific but are important in circumventing immune recognition or to immobilize effector T cells. Thus, the expression of these ligands and receptors may be associated with breast cancer stage and clinical outcome. (AKT. serine/threonine kinase or protein kinase B; ARG1. Arginase 1; Bcl-xL. B-cell lymphoma-xtra large; CAFs. Cancer-associated fibroblasts; CTLA4. cytotoxic T-lymphocyte-associated protein 4; ER– Estrogen Receptor Negative; GM-CSF. Granulocyte-Macrophage Colony-Stimulating Factor; HER2+. Human Epidermal Growth Factor Receptor 2; IDO. Indoleamine-2,3-dioxygenase; IFN-γ. interferon gamma; IFN-γR. interferon gamma receptor; IL. Interleukin; JAK. Janus kinase; iNOS. Inducible nitric oxide synthase; MDSC. Myeloid-derived suppressor cells; MHC. major histocompatibility complex; MMP. Matrix Metalloproteinases; mTOR. Mammalian target of rapamycin; MUC. Mucins; NK. natural killer; NF-κB. nuclear factor-κB; NLR. Neutrophil-lymphocyte ratio; NO. Nitric Oxide; PI3K. PI3K, phosphoinositide 3-kinase; PD-1. programmed death-1; PD-L1. Programmed death-ligand1; PGE₂. Prostaglandin E₂; PR–. Progesterone Receptor Negative; ROS. Reactive Oxygen Species; SHP. Src homology protein-tyrosine phosphatase; Siglec 9. Sialic acid-binding lectins 9; STAT. Signal transducer and activator of transcription; TAMs. Tumor-associated macrophages; TANs. Tumor-associated neutrophils; TGF-β. Transforming Growth Factor-beta; TIL. Tumor-Infiltrating Lymphocytes; TNBC. Triple Negative Breast Cancer; Treg. regulatory T cell; TCR. T cell receptor; VEGF. Vascular endothelial growth factor).

3.1. Cytotoxic T Lymphocyte-Associated Protein 4

Cytotoxic T lymphocyte-associated protein 4 (CTLA-4, CD152), expressed on the surface of activated T cells and a subset of Treg cells, functions as an immune checkpoint molecule which can downregulate T cells and inhibit anti-tumor responses [132]. CTLA-4 interacts with the cell surface immune stimulatory markers CD80 and CD86 on dendritic cells, resulting in the activation of dendritic cells and CD4/CD8 T cells. Treg cells (CD4+Foxp3+) are able to block this interaction resulting in decreased dendritic cell activation, inhibition of cytokine production (IL-2), T cell cycle arrest and suppression of CD8+ T cell proliferation [132]. One approach cancer cells have evolved to escape

from the immune system, is via the expression of CTLA-4 on their surface and in a limited number of studies, it has been shown to decrease survival in non-small lung cancer and nasopharyngeal cancer patients [133]. In vitro, co-culture of monocyte-derived human dendritic cells activated with lipopolysaccharide and CTLA-4+ breast cancer cell line, resulted in decreased expression of MHC class II, and costimulatory cell surface molecules (CD40, CD80, CD83, CD86) on dendritic cells and inability to activate T cells [133]. The addition of anti-CTLA-4 monoclonal antibody was able to reverse dendritic cell suppression and activate CD8+ T cells and induce apoptosis of CTLA-4+ breast cancer cells. Thus, CTLA-4 blockade is a promising approach for therapeutic studies in breast cancer patients, with or without combination with immunotherapy. Further, CTLA-4 expression by T cells is associated with better prognosis, whereas the expression of CTLA-4 on tumor cells is associated with poor prognosis [134].

3.2. Programmed Cell Death Protein 1

Programmed cell death protein 1 (PD-1, CD279) which is homologous to CD28, is an inhibitory immune signaling molecule and regulates adaptive immune responses. PD-1 is not expressed on circulating T cells but is expressed on activated T cells via TGF β , IL-2, IL-7, IL-15 and IL-21. PD-1 is also expressed by activated NK cells, B cells, monocytes, dendritic cells, myeloid cells, and thymocytes. The ligands (L) for PD-1 are PD-L1 (B7-H1, CD274) and PD-L2 (B7-DC, CD273) which are expressed on dendritic cells and monocytes and are upregulated by IFN γ , GM-CSF, vascular endothelial growth factor, VEGF, lipopolysaccharide, IL-10 and IL-14 [133]. The interaction between PD-1 and its ligands PD-L1 or PD-L2 leads to decreased T cell activity and its expression has been associated with poor prognosis [135]. In breast cancer, in particular in TNBC subtype, PD-L1 is upregulated which suppresses tumor-infiltrating T cells [136] resulting in higher Ki-67+ cells, basal-like subtypes, and distant metastasis [137]. Furthermore, analysis of PD-L1 expression in breast cancer cell lines showed higher PD-L1 expression in basal and mesenchymal cell lines than luminal cell lines [138]. Clinical samples from breast cancer patients, show upregulated expression of PD-L1 and is associated with poor prognostic features: ductal type, large tumor size, high grade, ER-, PR-, ERB-B2 receptor tyrosine kinase 2 (ERBB2)+, high proliferation rate and aggressive molecular subtypes (basal and ERBB2-enriched) [138]. In the last decade a number of clinical trials have been conducted to block PD-L1/PD-1 axis on non-small cell lung carcinoma and melanoma [135]. In fact, as of March 2019 the FDA approved an anti-PD-L1 antibody, in combination with chemotherapy, for the treatment of triple-negative, metastatic breast cancer for patients whose tumors expressed PD-L1.

3.3. Indoleamine 2,3-Dioxygenase

Indoleamine 2,3-dioxygenase (IDO) is an enzyme which catalyzes the oxidative break-down of tryptophan via kynurenine pathway in the presence of IFN γ [139]. Tryptophan is a vital amino acid for cell survival, and a lack of tryptophan in the TME leads to inhibition of T cell proliferation, thus, IDO exerts an immunosuppressive effect allowing tumor cells to escape from immune cells [130,139,140]. There are a number of mechanisms by which IDO influences the anergy of T cell activity within the TME. Firstly, by inhibiting mammalian target of rapamycin complex 1 and protein kinase C, both of which are regulators of glucokinase (GLK1), an important glucose regulator [141,142]. Secondly, the expression of IDO on cancer cells can activate the general control non-depressible-2 (GCN2) inducing a stress response in cells. GCN2 is stimulated by tryptophan transfer RNA (tRNA) in cells from accumulated uncharged tryptophan as a result of tryptophan degradation via IDO. Stimulation of GCN2 by the tryptophan tRNA alters protein translation and, prevents activation of T cells and promotes Treg cell differentiation [143–145]. Thirdly, the production of kynurenine, resulting from catabolism of tryptophan, activates transcription of the aryl hydrocarbon receptor (AHR), leading to differentiation of Foxp3+ Treg cells which suppress of anti-tumor responses [146,147]. Expression of high levels of IDO by cancer cells has been correlated with poor prognosis and reduced overall survival in patients with solid tumors including, breast cancer [140]. In fact, microvesicles released by malignant tumors,

in breast cancer patients contained a significantly high expression of IDO compared to normal breast tissues or benign tumor samples [148,149]. In addition, IDO expression is higher in advanced stages of breast cancer and mainly expressed in the TNBC subgroup [148,149]. The expression of IDO in TNBC has also shown to correlate with high expressions of PD-L1 in invasive primary breast carcinomas which highlights the need for combination therapies over single ones [150].

3.4. Sialic Acid-Binding Immunoglobulin-Type Lectin

Sialic acid-binding immunoglobulin-type lectins (Siglecs) are expressed on the surface of immune cells and to date, 15 different Siglecs have been identified. In particular Siglec-7 on NK cells and Siglec-9 on neutrophils, dendritic cells, monocytes, are associated with anti-tumor immunity [151]. In particular, Siglec-9 interacts with sialic acids on cancer cells (i.e., with MUC1, a high molecular weight glycoprotein expressed on adenocarcinoma cells including breast cancer) leading to immune suppression [152]. The mucous barrier that protects the epithelia contains secreted and transmembrane mucins. These membrane-bound mucins have important biological roles in cell-cell and cell-matrix interactions [20,25]. Human adenocarcinomas overexpress tumor-associated transmembrane mucins, such as MUC1 and MUC4, and are associated with tumor progression by enhancing their role in cell growth and survival [153,154]. Thus, mucins have been identified as potential prognostic and therapeutic targets for breast cancer development. In fact, vaccines/immunotherapeutic strategies targeting MUC1 have shown promise in pre-clinical animal models and in human clinical trials, with immune cell activation and clinical responses [24,25,42,48,155–161]. MUC1 is cleaved into N- and C-terminal subunits (MUC1-N and MUC1-C) which form a heterodimeric complex expressed at the cell membrane [162]. Studies have shown that MUC1-C functions as an oncoprotein by interacting with epidermal growth factor receptor (EGFR), ERbB2, and other tyrosine kinase receptors [163]. MUC1 also contributes to the malignant phenotype of cancer cells by binding to β -catenin, blocking phosphorylation and degradation [163,164]. In fact, the interaction between Siglec-9 and MUC1 via sialylated O-glycans, aids in cell growth by stimulating the recruitment of β -catenin [152]. In breast cancer, overexpression of MUC1 is associated with poor prognosis [165], suggesting that Siglec-9 may play an important role in the progression of breast cancer. Hence, studies that evaluate the complex interaction between Siglec-9-MUC1 and breast cancer, as well as the mechanism of this interaction in vitro and in vivo are warranted. In addition, human clinical studies utilizing MUC1 as a target should include blockade of Siglec-9 for improved clinical outcomes to patients with breast and other cancers. In fact, at the American Association for Cancer Research Meeting in Chicago April 2018, it was announced that first-in-class antibodies against Siglec-9 immune checkpoints are being developed for cancer immunotherapy (Benac, O., et al., conference presentation).

Table 1. Cells within the tumor microenvironment and their role in breast cancer.

Cell Type	Mechanisms	Model	Detection	Ref
Epithelial-mesenchymal transition (EMT) (via transcription factors <i>Snail</i> , <i>Twist-1</i> and <i>Lox</i> and markers vimentin and N-cadherin)	Involved in tumor progression and metastasis through signaling pathways such as TGF β , NF- κ B, Wnt, Notch			[60,61]
	<i>Lox</i> important in extracellular matrix (ECM) \rightarrow invasion and metastasis \uparrow <i>Lox</i> in TNBC	Triple negative breast cancer (TNBC) and non-TNBC human samples	Immunohistochemistry (IHC)	[62]
	NF- κ B \rightarrow \uparrow <i>Snail</i> \rightarrow TNF- α induced EMT	MCF-7 Cell line	Cell migration (CM) Real time polymerase chain reaction (RT-PCR) Western Blot (WB)	[166]
	MSC's (Mesenchymal stem cells) \uparrow metastasis through facilitation of EMT	MDA-MB-231, T47D and SK-Br ₃ cell lines	Low-density array RT-PCR Gene expression and proliferation assays	[167]

Table 1. Cont.

Cell Type	Mechanisms	Model	Detection	Ref
Immune cells Tumor-infiltrating Lymphocytes (TILs), T-cells (Tregs: CD4, CD25, FOXP3; CD8+; CD4+ Th cells)	Suppress T-cell proliferation Induce tumor cell death via IFN- γ and granzyme-perforin molecules			[74]
	naïve CD4+ T cell recruitment \rightarrow \downarrow Immunosuppression	MDA-MB-231 cell line Primary breast carcinoma	IHC and immunofluorescence staining, Flow cytometry (FC), Migration assay	[70]
CD4+ naïve T cell		Female NOD/scid mice Humanized mice NOD/SCID/IL2ry ^{null} (NSG)	qRT-PCR, Binding assays, WB	
Tumor-associated	TAMs are re-programmed to inhibit lymphocyte functions through release of inhibitory cytokines such as IL-10, prostaglandins or reactive oxygen species (ROS)			[91,92]
macrophage (TAM)	\uparrow CD163+ in non-luminal and basal-like breast cancer	Human tumor tissue	IHC with CD163	[96]
	breast cancer cell-secreted factors modulate macrophage differentiation to M2 status	Human tumor tissue Cell line MCF-7, MDA-MB231 and T47D	IHC FC ELISA Zymography	[168]
	\uparrow CD163+ in tumor stroma of TNBC	Human Luminal A and Triple Neg/basal-like tissue	IHC Gene Expression	[169]
Cancer-associated Fibroblast (CAF)	Shown to secrete various growth factors and cytokines associated with promoting breast cancer proliferation			[79]
	CAFs derived from Her2+ breast cancers \rightarrow \uparrow actin cytoskeleton and integrin signaling	Breast tumors sub grouped according to receptor expression	IHC Gene Expression	[85]
	ER+ expressing CD146 ^{neg} \rightarrow \uparrow tumor resistance to tamoxifen	Human tissue (Stage II & III, ER+ and/or ER-)	Immunocytochemistry (ICC)	[170]
	ER+ expressing CD146 ^{pos} \rightarrow \downarrow tumor resistance to tamoxifen	MCF-7 cell line	Gene expression	
	TNBC exhibit CAF subsets, \uparrow CAF-S1 \rightarrow \uparrow T Lymphocyte survival \rightarrow \uparrow Treg \rightarrow \emptyset T effector proliferation \rightarrow Immunosuppression	Female BC patient cohort (Luminal, HER2 and TN subtype tissues)	FC IHC	[171]
Tumor-associated Neutrophils (TANs)	N2 phenotype: pro-tumorigenic or pro-inflammatory			[172]
	N1 phenotype: anti-tumorigenic			
	Oncostatin M expressed by TANs \rightarrow \uparrow angiogenesis and metastasis	MDA-MB-231 & T47D cell lines	ICC ELISA	[112]
	\uparrow TAN in TNBC	Stage I-III breast cancer patient tumors divided into three subtypes: hormone-receptor [HR]-positive, HER2-negative (HR+, HER2-ve); HER2-positive and triple negative (TN)	Hematoxylin & eosin IHC	[115]
	\uparrow T β RIII (TGF- β receptor) in TNBC \rightarrow \uparrow mesenchymal-stem like (MSL) TNBC cells \rightarrow cell migration, invasion, and tumorigenicity	MSL cell lines SUM159, MDA-MB-231 and MDA-MB-157	Cell proliferation assay CM and invasion assay Immunoblotting FC Gene Expression	[116]
Tumor-associated Eosinophils	High presence of eosinophils at biopsy site may be linked to proliferation rate of tumor cells adjacent to wound	Female patients with primary breast cancer	Peripheral eosinophil counts	[173]
Myeloid-derived suppressor cells (MDSCs)	\uparrow Arginase 1 (ARG1) + nitric oxide synthase (iNOS) \rightarrow \uparrow superoxide and nitric oxide (NO) \rightarrow \emptyset lymphocyte responses \rightarrow \uparrow iNOS in surrounding cells \rightarrow \uparrow tumor growth and \downarrow immune cell functions			[127,128]
	stage IV patients with extensive metastasis \rightarrow \uparrow MDSC	Blood from patients with stages I-IV solid malignancies obtained prior to surgery	FC	[174]
	\uparrow MDSC correlates with worse prognosis	Peripheral blood specimens stage IV breast cancer patients	FC Proliferation assay	[175]
	MDSC \uparrow IDO \rightarrow \downarrow tryptophan \rightarrow \emptyset T-cell proliferation and induced T-cell apoptosis	Female breast cancer patients (Stages I-III)	IHC RT-PCR WB ELISA	[176]

Table 2. Immune Checkpoints and Breast Cancer.

Immune Checkpoint Factor	Mechanisms	Model	Detection	Ref
CTLA4	Expressed on the surface of activated T-cells and a subset of Tregs ↓T-cell activation → Anti-T cell response			[177]
	CTLA4 blockade → ∅ proliferation and induced apoptosis of CTLA-4+ breast cancer cells	MDA-MB-231, SKBR3, MCF-7 and T47D cell lines	FC WB	[178]
	↑CTLA-4 in lymphocytes → better prognosis	130 BC patients	IHC	[134]
	↑CTLA-4 in T cells → worse prognosis			
PD-1 (PD-L1/PD-L2)	PD-1 is expressed by activated lymphocytes → ligation PD-L1/PD-L2 → ↓T-cell activity → poor prognosis			[135]
	TNBC subtype; ↑PD-L1 → suppresses auto-immunity—T cell proliferation—Cytokine production—Cytotoxic activity			[136]
	PDI ↑TILs, but ↓PDL1 in T cells → positive TNBC prognostic factor	negative ER, PR, and HER-2 BC patients	IHC RNAscope	[137]
	↑PD-L1 in Basal & TNBC subtypes → ↑cytotoxic local immune response → ↑better survival	BC patient is Cell lines		[138]
Indoleamine 2,3-dioxygenase (IDO)	Catalyzes the oxidative break-down of tryptophan via kynurenine pathway in the presence of IFN-γ → enabling immune escape			[130]
	Correlation between expression of IDO and PD-1 in myoeptithelial, stromal, and T cells	Human BC patient tissues and healthy tissues	IHC WB RT-PCR	[179]
	↑IDO expression in TNBC and basal-like BC	200 TNBC patients	IHC RT-PCR	[180]
	IDO expression ↑ER+ as compared to ER− is	breast cancer tissue sections	IHC	[52]
Siglec-9	Siglec-9 found on neutrophils and Siglec-7 found on NK cells have been associated with anti-T immunity			[151]
	Siglec-9 → T-associated MUC1 downstream signal transduction, following T cell proliferation	Transgenic mice and murine mammary T cell	WB, Immunoperoxidase staining, IHC Gene expression Proliferation (MTT) assay	[181]
	↑Siglec-9 on DCs involved in immunoregulation through ligation with mucins in epithelial cancer	Human colon cancer cell line (LS 180 cells)	FC RT-PCR ELISA	[182]

4. Conclusions and Future Prospects

The interaction between the immune system and the TME have become a pivotal point of research. The ability of cancer cells to signal pro-tumorigenic modifications to normal anti-tumorigenic cells to aid in cell proliferation and angiogenesis has provided multiple advancements in cancer therapies and prognostic value. As a consequence, not all current therapies and treatments have been able to provide relief from the 100 or more types of cancers, with breast cancer being the most common among women. In addition, in breast cancer, each subtype and stage has risk factors for incidence, treatment response, rate of disease progression and metastasis associated with it [53]. Current studies continue to isolate the different immune cell populations and those found within the TME of breast cancers, their association with hormone receptor phenotype and immune-evasion mechanisms. The interaction between immune cells and cancer cells in the TME and immune-checkpoints could potentially be used for identification of new markers associated with progression of tumors in their early developmental stages.

Currently, breast cancer-positive hormone receptor phenotypes (ER+, PR+, HER2+) have more treatment options with favorable outcomes compared to the TNBC subtype. The immune cells function and their cytokines are key factors whose modulation strongly encourages further study and consideration as predictive markers and important therapeutic targets in different subtypes of breast cancer [183]. No systemic method has been associated with the combination of markers in early diagnosis of cancer patients. Although the current methods and biomarkers can be used to place patients within a clinical stage critical for treatment options, having a systemic method that correlates the expression of biological, immunological, and potential neurological markers could pave the way into new prognostic and clinical outcomes of cancer patients. In addition, all this combined with immune checkpoint inhibitors and cancer immunotherapeutics would yield new improved treatments for breast cancer.

Author Contributions: V.B. and N.K. wrote the article under the supervision of V.A. and K.N. and all authors reviewed and edited the article.

Funding: This research review received no external funding.

Acknowledgments: The authors would like to thank Graeme Wise and Angelina Wise, whose generous philanthropic support made possible the preparation of this paper. V.A. and K.N. would like to thank the Mechanisms and Interventions in Health and Disease Program within the Institute for Health and Sport, Victoria University Australia for their support. V.B. was supported by the College of Health and Biomedicine BSc Honors Program and N.K. was supported by the Victoria University Postgraduate Scholarship, Victoria University Australia.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Boyle, P.; Ferlay, J. Mortality and survival in breast and colorectal cancer. *Nat. Clin. Pract. Oncol.* **2005**, *2*, 424. [[CrossRef](#)] [[PubMed](#)]
- Pudkasam, S.; Tangalakis, K.; Chinlumprasert, N.; Apostolopoulos, V.; Stojanovska, L. Breast cancer and exercise: The role of adiposity and immune markers. *Maturitas* **2017**, *105*, 16–22. [[CrossRef](#)] [[PubMed](#)]
- Smith, R.A.; Duffy, S.W.; Gabe, R.; Tabar, L.; Yen, A.M.; Chen, T.H. The randomized trials of breast cancer screening: What have we learned? *Radiol. Clin. North Am.* **2004**, *42*, 793–806. [[CrossRef](#)] [[PubMed](#)]
- Smith, B.D.; Smith, G.L.; Hurria, A.; Hortobagyi, G.N.; Buchholz, T.A. Future of cancer incidence in the United States: Burdens upon an aging, changing nation. *J. Clin. Oncol.* **2009**, *27*, 2758–2765. [[CrossRef](#)] [[PubMed](#)]
- Burness, M.L.; Van Poznak, C. Metastatic breast cancer: Clinical considerations. In *Pathobiology of Human Disease*; McManus, L.M., Mitchell, R.N., Eds.; Elsevier: Amsterdam, The Netherlands, 2014; pp. 963–970.
- Lal, S.; McCart Reed, A.E.; de Luca, X.M.; Simpson, P.T. Molecular signatures in breast cancer. *Methods* **2017**, *131*, 135–146. [[CrossRef](#)] [[PubMed](#)]
- Rakha, E.A.; Reis-Filho, J.S.; Baehner, F.; Dabbs, D.J.; Decker, T.; Eusebi, V.; Fox, S.B.; Ichihara, S.; Jacquemier, J.; Lakhani, S.R.; et al. Breast cancer prognostic classification in the molecular era: The role of histological grade. *Breast Cancer Res.* **2010**, *12*, 207. [[CrossRef](#)]
- Weigelt, B.; Reis-Filho, J.S. Histological and molecular types of breast cancer: Is there a unifying taxonomy? *Nat. Rev. Clin. Oncol.* **2009**, *6*, 718–730. [[CrossRef](#)]
- Weigelt, B.; Baehner, F.L.; Reis-Filho, J.S. The contribution of gene expression profiling to breast cancer classification, prognostication and prediction: A retrospective of the last decade. *J. Pathol.* **2010**, *220*, 263–280. [[CrossRef](#)]
- Geyer, F.C.; Reis-Filho, J.S. Microarray-based gene expression profiling as a clinical tool for breast cancer management: Are we there yet? *Int. J. Surg. Pathol.* **2009**, *17*, 285–302. [[CrossRef](#)]
- Geyer, F.C.; Lopez-Garcia, M.A.; Lambros, M.B.; Reis-Filho, J.S. Genetic characterization of breast cancer and implications for clinical management. *J. Cell. Mol. Med.* **2009**, *13*, 4090–4103. [[CrossRef](#)]
- Kuol, N.; Stojanovska, L.; Apostolopoulos, V.; Nurgali, K. Role of the Nervous System in Tumor Angiogenesis. *Cancer Microenviron.* **2018**, *11*, 1–11. [[CrossRef](#)]
- Kuol, N.; Stojanovska, L.; Apostolopoulos, V.; Nurgali, K. Role of the nervous system in cancer metastasis. *J. Exp. Clin. Cancer Res.* **2018**, *37*, 5. [[CrossRef](#)]
- Kuol, N.; Stojanovska, L.; Apostolopoulos, V.; Nurgali, K. Crosstalk between cancer and the neuro-immune system. *J. Neuroimmunol.* **2018**, *315*, 15–23. [[CrossRef](#)]
- Kuol, N.; Stojanovska, L.; Nurgali, K.; Apostolopoulos, V. The mechanisms tumor cells utilize to evade the host's immune system. *Maturitas* **2017**, *105*, 8–15. [[CrossRef](#)]
- Kuol, N.; Stojanovska, L.; Nurgali, K.; Apostolopoulos, V. PD-1/PD-L1 in disease. *Immunotherapy* **2018**, *10*, 149–160. [[CrossRef](#)] [[PubMed](#)]
- Hudak, J.E.; Canham, S.M.; Bertozzi, C.R. Glycocalyx engineering reveals a Siglec-based mechanism for NK cell immunoevasion. *Nat. Chem. Biol.* **2014**, *10*, 69–75. [[CrossRef](#)]
- Beatson, R.; Tajadura-Ortega, V.; Achkova, D.; Picco, G.; Tsourouksoglou, T.D.; Klausning, S.; Hillier, M.; Maher, J.; Noll, T.; Crocker, P.R.; et al. The mucin MUC1 modulates the tumor immunological microenvironment through engagement of the lectin Siglec-9. *Nat. Immunol.* **2016**, *17*, 1273–1281. [[CrossRef](#)]

19. Solinas, G.; Germano, G.; Mantovani, A.; Allavena, P. Tumor-associated macrophages (TAM) as major players of the cancer-related inflammation. *J. Leukoc. Biol.* **2009**, *86*, 1065–1073. [\[CrossRef\]](#)
20. Apostolopoulos, V.; Stojanovska, L.; Gargosky, S.E. MUC1 (CD227): A multi-tasked molecule. *Cell. Mol. Life Sci. CMLS* **2015**, *72*, 4475–4500. [\[CrossRef\]](#)
21. Apostolopoulos, V. Peptide-based vaccines for cancer: Are we choosing the right peptides? *Expert Rev. Vaccines* **2009**, *8*, 259–260. [\[CrossRef\]](#)
22. Apostolopoulos, V.; McKenzie, I.F. Cellular mucins: Targets for immunotherapy. *Crit. Rev. Immunol.* **1994**, *14*, 293–309. [\[CrossRef\]](#)
23. Apostolopoulos, V.; McKenzie, I.F.; Pietersz, G.A. Breast cancer immunotherapy: Current status and future prospects. *Immunol. Cell Biol.* **1996**, *74*, 457–464. [\[CrossRef\]](#)
24. Apostolopoulos, V.; McKenzie, I.F.C. Cellular Mucins: Targets for Immunotherapy. *Crit. Rev. Immunol.* **2017**, *37*, 421–437. [\[CrossRef\]](#)
25. Apostolopoulos, V.; Pietersz, G.A.; McKenzie, I.F. MUC1 and breast cancer. *Curr. Opin. Mol. Ther.* **1999**, *1*, 98–103.
26. Apostolopoulos, V.; Osinski, C.; McKenzie, I.F. MUC1 cross-reactive Gal α (1,3)Gal antibodies in humans switch immune responses from cellular to humoral. *Nat. Med.* **1998**, *4*, 315–320. [\[CrossRef\]](#)
27. Apostolopoulos, V.; Pietersz, G.A.; Gordon, S.; Martinez-Pomares, L.; McKenzie, I.F. Aldehyde-mannan antigen complexes target the MHC class I antigen-presentation pathway. *Eur. J. Immunol.* **2000**, *30*, 1714–1723. [\[CrossRef\]](#)
28. Apostolopoulos, V.; Pietersz, G.A.; Loveland, B.E.; Sandrin, M.S.; McKenzie, I.F. Oxidative/reductive conjugation of mannan to antigen selects for T1 or T2 immune responses. *Proc. Natl. Acad. Sci. USA* **1995**, *92*, 10128–10132. [\[CrossRef\]](#)
29. Apostolopoulos, V.; Pietersz, G.A.; McKenzie, I.F. Cell-mediated immune responses to MUC1 fusion protein coupled to mannan. *Vaccine* **1996**, *14*, 930–938. [\[CrossRef\]](#)
30. Apostolopoulos, V.; Sandrin, M.S.; McKenzie, I.F. Mimics and cross reactions of relevance to tumour immunotherapy. *Vaccine* **1999**, *18*, 268–275. [\[CrossRef\]](#)
31. Apostolopoulos, V.; Sandrin, M.S.; McKenzie, I.F. Carbohydrate/peptide mimics: Effect on MUC1 cancer immunotherapy. *J. Mol. Med.* **1999**, *77*, 427–436. [\[CrossRef\]](#)
32. Apostolopoulos, V.; Xing, P.X.; McKenzie, I.F. Murine immune response to cells transfected with human MUC1: Immunization with cellular and synthetic antigens. *Cancer Res.* **1994**, *54*, 5186–5193.
33. Apostolopoulos, V.; Yu, M.; Corper, A.L.; Teyton, L.; Pietersz, G.A.; McKenzie, I.F.; Wilson, I.A.; Plebanski, M. Crystal structure of a non-canonical low-affinity peptide complexed with MHC class I: A new approach for vaccine design. *J. Mol. Biol.* **2002**, *318*, 1293–1305. [\[CrossRef\]](#)
34. Apostolopoulos, V.; Yuriev, E.; Ramsland, P.A.; Halton, J.; Osinski, C.; Li, W.; Plebanski, M.; Paulsen, H.; McKenzie, I.F. A glycopeptide in complex with MHC class I uses the GalNAc residue as an anchor. *Proc. Natl. Acad. Sci. USA* **2003**, *100*, 15029–15034. [\[CrossRef\]](#)
35. Lazoura, E.; Apostolopoulos, V. Rational Peptide-based vaccine design for cancer immunotherapeutic applications. *Curr. Med. Chem.* **2005**, *12*, 629–639. [\[CrossRef\]](#)
36. Lees, C.J.; Apostolopoulos, V.; Acres, B.; Ong, C.S.; Popovski, V.; McKenzie, I.F. The effect of T1 and T2 cytokines on the cytotoxic T cell response to mannan-MUC1. *Cancer Immunol. Immunother.* **2000**, *48*, 644–652. [\[CrossRef\]](#)
37. Lees, C.J.; Apostolopoulos, V.; Acres, B.; Ramshaw, I.; Ramsay, A.; Ong, C.S.; McKenzie, I.F. Immunotherapy with mannan-MUC1 and IL-12 in MUC1 transgenic mice. *Vaccine* **2000**, *19*, 158–162. [\[CrossRef\]](#)
38. Lees, C.J.; Apostolopoulos, V.; McKenzie, I.F. Cytokine production from murine CD4 and CD8 cells after mannan-MUC1 immunization. *J. Interferon Cytokine Res.* **1999**, *19*, 1373–1379. [\[CrossRef\]](#)
39. Lees, C.J.; Smorodinsky, N.; Horn, G.; Wreschner, D.H.; McKenzie, I.F.; Pietersz, G.; Stojanovska, L.; Apostolopoulos, V. MUC1 immunotherapy against a metastatic mammary adenocarcinoma model: Importance of IFN- γ . *Prilozi* **2016**, *37*, 15–25. [\[CrossRef\]](#)
40. Sheng, K.C.; Kalkanidis, M.; Pouniotis, D.S.; Esparon, S.; Tang, C.K.; Apostolopoulos, V.; Pietersz, G.A. Delivery of antigen using a novel mannosylated dendrimer potentiates immunogenicity in vitro and in vivo. *Eur. J. Immunol.* **2008**, *38*, 424–436. [\[CrossRef\]](#)

41. Son, H.Y.; Apostolopoulos, V.; Chung, J.K.; Kim, C.W.; Park, J.U. Protective efficacy of a plasmid DNA vaccine against transgene-specific tumors by Th1 cellular immune responses after intradermal injection. *Cell. Immunol.* **2018**, *329*, 17–26. [[CrossRef](#)]
42. Tang, C.K.; Apostolopoulos, V. Strategies used for MUC1 immunotherapy: Preclinical studies. *Expert Rev. Vaccines* **2008**, *7*, 951–962. [[CrossRef](#)]
43. Tang, C.K.; Sheng, K.C.; Pouniotis, D.; Esparon, S.; Son, H.Y.; Kim, C.W.; Pietersz, G.A.; Apostolopoulos, V. Oxidized and reduced mannan mediated MUC1 DNA immunization induce effective anti-tumor responses. *Vaccine* **2008**, *26*, 3827–3834. [[CrossRef](#)]
44. Karanikas, V.; Hwang, L.A.; Pearson, J.; Ong, C.S.; Apostolopoulos, V.; Vaughan, H.; Xing, P.X.; Jamieson, G.; Pietersz, G.; Tait, B.; et al. Antibody and T cell responses of patients with adenocarcinoma immunized with mannan-MUC1 fusion protein. *J. Clin. Investig.* **1997**, *100*, 2783–2792. [[CrossRef](#)]
45. Loveland, B.E.; Zhao, A.; White, S.; Gan, H.; Hamilton, K.; Xing, P.X.; Pietersz, G.A.; Apostolopoulos, V.; Vaughan, H.; Karanikas, V.; et al. Mannan-MUC1-pulsed dendritic cell immunotherapy: A phase I trial in patients with adenocarcinoma. *Clin. Cancer Res.* **2006**, *12*, 869–877. [[CrossRef](#)]
46. Xing, P.; Michael, M.; Apostolopoulos, V.; Prenzloska, J.; Marshall, C.; Bishop, J.; McKenzie, I. Phase-I study of synthetic muc1 peptides in breast-cancer. *Int. J. Oncol.* **1995**, *6*, 1283–1289. [[CrossRef](#)]
47. Apostolopoulos, V.; Pietersz, G.A.; Tsibanis, A.; Tsikkinis, A.; Drakaki, H.; Loveland, B.E.; Piddlesden, S.J.; Plebanski, M.; Pouniotis, D.S.; Alexis, M.N.; et al. Pilot phase III immunotherapy study in early-stage breast cancer patients using oxidized mannan-MUC1 [ISRCTN71711835]. *Breast Cancer Res.* **2006**, *8*, R27. [[CrossRef](#)]
48. Apostolopoulos, V.; Pietersz, G.A.; Tsibanis, A.; Tsikkinis, A.; Stojanovska, L.; McKenzie, I.F.; Vassilaros, S. Dendritic cell immunotherapy: Clinical outcomes. *Clin. Transl. Immunol.* **2014**, *3*, e21. [[CrossRef](#)]
49. Apostolopoulos, V.; Thalhammer, T.; Tzakos, A.G.; Stojanovska, L. Targeting antigens to dendritic cell receptors for vaccine development. *J. Drug Deliv.* **2013**, *2013*, 869718. [[CrossRef](#)]
50. Vassilaros, S.; Tsibanis, A.; Tsikkinis, A.; Pietersz, G.A.; McKenzie, I.F.; Apostolopoulos, V. Up to 15-year clinical follow-up of a pilot Phase III immunotherapy study in stage II breast cancer patients using oxidized mannan-MUC1. *Immunotherapy* **2013**, *5*, 1177–1182. [[CrossRef](#)]
51. Miller, S.; Senior, P.V.; Prakash, M.; Apostolopoulos, V.; Sakkal, S.; Nurgali, K. Leukocyte populations and IL-6 in the tumor microenvironment of an orthotopic colorectal cancer model. *Acta Biochim. Biophys. Sin.* **2016**, *48*, 334–341. [[CrossRef](#)]
52. Soliman, H.; Rawal, B.; Fulp, J.; Lee, J.H.; Lopez, A.; Bui, M.M.; Khalil, F.; Antonia, S.; Yfantis, H.G.; Lee, D.H.; et al. Analysis of indoleamine 2–3 dioxygenase (IDO1) expression in breast cancer tissue by immunohistochemistry. *Cancer Immunol. Immunother.* **2013**, *62*, 829–837. [[CrossRef](#)]
53. Nagarajan, D.; McArdle, S.E.B. Immune Landscape of Breast Cancers. *Biomedicines* **2018**, *6*, 20. [[CrossRef](#)]
54. Gajewski, T.F.; Schreiber, H.; Fu, Y.-X. Innate and adaptive immune cells in the tumor microenvironment. *Nat. Immunol.* **2013**, *14*, 1014–1022. [[CrossRef](#)]
55. Yuan, Y.; Jiang, Y.C.; Sun, C.K.; Chen, Q.M. Role of the tumor microenvironment in tumor progression and the clinical applications (Review). *Oncol. Rep.* **2016**, *35*, 2499–2515. [[CrossRef](#)]
56. Cirri, P.; Chiarugi, P. Cancer associated fibroblasts: The dark side of the coin. *Am. J. Cancer Res.* **2011**, *1*, 482–497.
57. Gabrilovich, D.I.; Ostrand-Rosenberg, S.; Bronte, V. Coordinated regulation of myeloid cells by tumours. *Nat. Rev. Immunol.* **2012**, *12*, 253–268. [[CrossRef](#)]
58. Cheng, L.; Huang, Z.; Zhou, W.; Wu, Q.; Donnola, S.; Liu, J.K.; Fang, X.; Sloan, A.E.; Mao, Y.; Lathia, J.D.; et al. Glioblastoma stem cells generate vascular pericytes to support vessel function and tumor growth. *Cell* **2013**, *153*, 139–152. [[CrossRef](#)]
59. Whiteside, T.L. The tumor microenvironment and its role in promoting tumor growth. *Oncogene* **2008**, *27*, 5904–5912. [[CrossRef](#)]
60. Jing, Y.; Han, Z.; Zhang, S.; Liu, Y.; Wei, L. Epithelial-Mesenchymal Transition in tumor microenvironment. *Cell Biosci.* **2011**, *1*, 29. [[CrossRef](#)]
61. Wang, Y.; Zhou, B.P. Epithelial-mesenchymal transition in breast cancer progression and metastasis. *Chin. J. Cancer* **2011**, *30*, 603–611. [[CrossRef](#)]
62. Leo, C.; Cotic, C.; Pomp, V.; Fink, D.; Varga, Z. Overexpression of Lox in triple-negative breast cancer. *Ann. Diagn. Pathol.* **2018**, *34*, 98–102. [[CrossRef](#)]

63. Onder, T.T.; Gupta, P.B.; Mani, S.A.; Yang, J.; Lander, E.S.; Weinberg, R.A. Loss of E-cadherin promotes metastasis via multiple downstream transcriptional pathways. *Cancer Res.* **2008**, *68*, 3645–3654. [\[CrossRef\]](#)
64. Kang, Y.; Massague, J. Epithelial-mesenchymal transitions: Twist in development and metastasis. *Cell* **2004**, *118*, 277–279. [\[CrossRef\]](#)
65. El-Haibi, C.P.; Bell, G.W.; Zhang, J.; Collmann, A.Y.; Wood, D.; Scherber, C.M.; Csizmadia, E.; Mariani, O.; Zhu, C.; Campagne, A.; et al. Critical role for lysyl oxidase in mesenchymal stem cell-driven breast cancer malignancy. *Proc. Natl. Acad. Sci. USA* **2012**, *109*, 17460–17465. [\[CrossRef\]](#)
66. Markiewicz, A.; Nagel, A.; Szade, J.; Majewska, H.; Skokowski, J.; Seroczynska, B.; Stokowy, T.; Welnicka-Jaskiewicz, M.; Zaczek, A.J. Aggressive Phenotype of Cells Disseminated via Hematogenous and Lymphatic Route in Breast Cancer Patients. *Transl. Oncol.* **2018**, *11*, 722–731. [\[CrossRef\]](#)
67. Qi, L.N.; Xiang, B.D.; Wu, F.X.; Ye, J.Z.; Zhong, J.H.; Wang, Y.Y.; Chen, Y.Y.; Chen, Z.S.; Ma, L.; Chen, J.; et al. Circulating Tumor Cells Undergoing EMT Provide a Metric for Diagnosis and Prognosis of Patients with Hepatocellular Carcinoma. *Cancer Res.* **2018**, *78*, 4731–4744. [\[CrossRef\]](#)
68. Ha, H.; Debnath, B.; Neamati, N. Role of the CXCL8-CXCR1/2 Axis in Cancer and Inflammatory Diseases. *Theranostics* **2017**, *7*, 1543–1588. [\[CrossRef\]](#)
69. Whiteside, T. *The Local Tumor Microenvironment*, 1st ed.; Springer: Heidelberg, Germany, 2007; p. 503.
70. Su, S.; Liao, J.; Liu, J.; Huang, D.; He, C.; Chen, F.; Yang, L.; Wu, W.; Chen, J.; Lin, L.; et al. Blocking the recruitment of naive CD4+ T cells reverses immunosuppression in breast cancer. *Cell Res.* **2017**, *27*, 461. [\[CrossRef\]](#)
71. Bense, R.D.; Sotiriou, C.; Piccart-Gebhart, M.J.; Haanen, J.B.A.G.; van Vugt, M.A.T.M.; de Vries, E.G.E.; Schröder, C.P.; Fehrmann, R.S.N. Relevance of Tumor-Infiltrating Immune Cell Composition and Functionality for Disease Outcome in Breast Cancer. *J. Natl. Cancer Inst.* **2017**, *109*. [\[CrossRef\]](#)
72. Woo, E.Y.; Chu, C.S.; Goletz, T.J.; Schlienger, K.; Yeh, H.; Coukos, G.; Rubin, S.C.; Kaiser, L.R.; June, C.H. Regulatory CD4(+)CD25(+) T cells in tumors from patients with early-stage non-small cell lung cancer and late-stage ovarian cancer. *Cancer Res.* **2001**, *61*, 4766–4772.
73. Strauss, L.; Bergmann, C.; Gooding, W.; Johnson, J.T.; Whiteside, T.L. The frequency and suppressor function of CD4+CD25highFoxp3+ T cells in the circulation of patients with squamous cell carcinoma of the head and neck. *Clin. Cancer Res.* **2007**, *13*, 6301–6311. [\[CrossRef\]](#)
74. Dushyanthen, S.; Beavis, P.A.; Savas, P.; Teo, Z.L.; Zhou, C.; Mansour, M.; Darcy, P.K.; Loi, S. Relevance of tumor-infiltrating lymphocytes in breast cancer. *BMC Med.* **2015**, *13*, 202. [\[CrossRef\]](#)
75. Hammerl, D.; Smid, M.; Timmermans, A.M.; Sleijfer, S.; Martens, J.W.M.; Debets, R. Breast cancer genomics and immuno-oncological markers to guide immune therapies. *Semin. Cancer Biol.* **2018**, *52*, 178–188. [\[CrossRef\]](#)
76. Devereaux, J.; Nurgali, K.; Kiatos, D.; Sakkal, S.; Apostolopoulos, V. Effects of platelet-rich plasma and platelet-poor plasma on human dermal fibroblasts. *Maturitas* **2018**, *117*, 34–44. [\[CrossRef\]](#)
77. Karagiannis, G.S.; Poutahidis, T.; Erdman, S.E.; Kirsch, R.; Riddell, R.H.; Diamandis, E.P. Cancer-Associated Fibroblasts Drive the Progression of Metastasis through both Paracrine and Mechanical Pressure on Cancer Tissue. *Mol. Cancer Res.* **2012**, *10*, 1403. [\[CrossRef\]](#)
78. Shiga, K.; Hara, M.; Nagasaki, T.; Sato, T.; Takahashi, H.; Takeyama, H. Cancer-Associated Fibroblasts: Their Characteristics and Their Roles in Tumor Growth. *Cancers* **2015**, *7*, 2443–2458. [\[CrossRef\]](#)
79. Buchsbaum, R.J.; Oh, S.Y. Breast Cancer-Associated Fibroblasts: Where we are and where we need to go. *Cancers* **2016**, *8*, 19. [\[CrossRef\]](#)
80. Martens, J.W.; Sieuwerts, A.M.; Bolt-deVries, J.; Bosma, P.T.; Swiggers, S.J.; Klijn, J.G.; Foekens, J.A. Aging of stromal-derived human breast fibroblasts might contribute to breast cancer progression. *Thromb. Haemost.* **2003**, *89*, 393–404.
81. Palmieri, C.; Roberts-Clark, D.; Assadi-Sabet, A.; Coope, R.C.; O'Hare, M.; Sunter, A.; Hanby, A.; Slade, M.J.; Gomm, J.J.; Lam, E.W.; et al. Fibroblast growth factor 7, secreted by breast fibroblasts, is an interleukin-1beta-induced paracrine growth factor for human breast cells. *J. Endocrinol.* **2003**, *177*, 65–81. [\[CrossRef\]](#)
82. Scherz-Shouval, R.; Santagata, S.; Mendillo, M.L.; Sholl, L.M.; Ben-Aharon, I.; Beck, A.H.; Dias-Santagata, D.; Koeva, M.; Stemmer, S.M.; Whitesell, L.; et al. The reprogramming of tumor stroma by HSF1 is a potent enabler of malignancy. *Cell* **2014**, *158*, 564–578. [\[CrossRef\]](#)

83. Pickup, M.; Novitskiy, S.; Moses, H.L. The roles of TGF β in the tumour microenvironment. *Nat. Rev. Cancer* **2013**, *13*, 788–799. [[CrossRef](#)]
84. De Wever, O.; Mareel, M. Role of tissue stroma in cancer cell invasion. *J. Pathol.* **2003**, *200*, 429–447. [[CrossRef](#)]
85. Tchou, J.; Kossenkova, A.V.; Chang, L.; Satija, C.; Herlyn, M.; Showe, L.C.; Puré, E. Human breast cancer associated fibroblasts exhibit subtype specific gene expression profiles. *BMC Med. Genomics* **2012**, *5*, 39. [[CrossRef](#)]
86. Apostolopoulos, V.; Barnes, N.; Pietersz, G.A.; McKenzie, I.F. Ex vivo targeting of the macrophage mannose receptor generates anti-tumor CTL responses. *Vaccine* **2000**, *18*, 3174–3184. [[CrossRef](#)]
87. Apostolopoulos, V.; Lofthouse, S.A.; Popovski, V.; Chelvanayagam, G.; Sandrin, M.S.; McKenzie, I.F. Peptide mimics of a tumor antigen induce functional cytotoxic T cells. *Nat. Biotechnol.* **1998**, *16*, 276–280. [[CrossRef](#)]
88. Sheng, K.C.; Day, S.; Wright, M.D.; Stojanovska, L.; Apostolopoulos, V. Enhanced Dendritic Cell-Mediated Antigen-Specific CD4⁺ T Cell Responses: IFN-Gamma Aids TLR Stimulation. *J. Drug Deliv.* **2013**, *2013*, 516749. [[CrossRef](#)]
89. Sheng, K.C.; Pouniotis, D.S.; Wright, M.D.; Tang, C.K.; Lazoura, E.; Pietersz, G.A.; Apostolopoulos, V. Mannan derivatives induce phenotypic and functional maturation of mouse dendritic cells. *Immunology* **2006**, *118*, 372–383. [[CrossRef](#)]
90. Tang, C.K.; Lodding, J.; Minigo, G.; Pouniotis, D.S.; Plebanski, M.; Scholzen, A.; McKenzie, I.F.; Pietersz, G.A.; Apostolopoulos, V. Mannan-mediated gene delivery for cancer immunotherapy. *Immunology* **2007**, *120*, 325–335. [[CrossRef](#)]
91. Mantovani, A.; Sozzani, S.; Locati, M.; Allavena, P.; Sica, A. Macrophage polarization: Tumor-associated macrophages as a paradigm for polarized M2 mononuclear phagocytes. *Trends Immunol.* **2002**, *23*, 549–555. [[CrossRef](#)]
92. Martinez, F.O.; Sica, A.; Mantovani, A.; Locati, M. Macrophage activation and polarization. *Front. Biosci.* **2008**, *13*, 453–461. [[CrossRef](#)]
93. Liou, G.Y.; Storz, P. Reactive oxygen species in cancer. *Free Radic. Res.* **2010**, *44*, 479–496. [[CrossRef](#)]
94. Alexandrow, M.G.; Moses, H.L. Transforming growth factor beta 1 inhibits mouse keratinocytes late in G1 independent of effects on gene transcription. *Cancer Res.* **1995**, *55*, 3928–3932.
95. Obeid, E.; Nanda, R.; Fu, Y.X.; Olopade, O.I. The role of tumor-associated macrophages in breast cancer progression (review). *Int. J. Oncol.* **2013**, *43*, 5–12. [[CrossRef](#)]
96. Klingen, T.A.; Chen, Y.; Aas, H.; Wik, E.; Akslen, L.A. Tumor-associated macrophages are strongly related to vascular invasion, non-luminal subtypes, and interval breast cancer. *Hum. Pathol.* **2017**, *69*, 72–80.
97. Yu, T.; Di, G. Role of tumor microenvironment in triple-negative breast cancer and its prognostic significance. *Chin. J. Cancer Res.* **2017**, *29*, 237–252. [[CrossRef](#)]
98. Pang, M.F.; Georgoudaki, A.M.; Lambut, L.; Johansson, J.; Tabor, V.; Hagikura, K.; Jin, Y.; Jansson, M.; Alexander, J.S.; Nelson, C.M.; et al. TGF-beta1-induced EMT promotes targeted migration of breast cancer cells through the lymphatic system by the activation of CCR7/CCL21-mediated chemotaxis. *Oncogene* **2016**, *35*, 748–760. [[CrossRef](#)]
99. Fuxe, J.; Karlsson, M.C. TGF-beta-induced epithelial-mesenchymal transition: A link between cancer and inflammation. *Semin. Cancer Biol.* **2012**, *22*, 455–461. [[CrossRef](#)]
100. Giampieri, S.; Manning, C.; Hooper, S.; Jones, L.; Hill, C.S.; Sahai, E. Localized and reversible TGFbeta signalling switches breast cancer cells from cohesive to single cell motility. *Nat. Cell Biol.* **2009**, *11*, 1287–1296. [[CrossRef](#)]
101. Su, S.; Liu, Q.; Chen, J.; Chen, J.; Chen, F.; He, C.; Huang, D.; Wu, W.; Lin, L.; Huang, W.; et al. A positive feedback loop between mesenchymal-like cancer cells and macrophages is essential to breast cancer metastasis. *Cancer Cell* **2014**, *25*, 605–620. [[CrossRef](#)]
102. Gan, L.; Qiu, Z.; Huang, J.; Li, Y.; Huang, H.; Xiang, T.; Wan, J.; Hui, T.; Lin, Y.; Li, H.; et al. Cyclooxygenase-2 in tumor-associated macrophages promotes metastatic potential of breast cancer cells through Akt pathway. *Int. J. Biol. Sci.* **2016**, *12*, 1533–1543.
103. Ding, J.; Jin, W.; Chen, C.; Shao, Z.; Wu, J. Tumor Associated Macrophage \times Cancer Cell Hybrids May Acquire Cancer Stem Cell Properties in Breast Cancer. *PLoS ONE* **2012**, *7*, e41942. [[CrossRef](#)]

104. Brady, N.J.; Chuntova, P.; Schwertfeger, K.L. Macrophages: Regulators of the Inflammatory Microenvironment during Mammary Gland Development and Breast Cancer. *Mediat. Inflamm.* **2016**, *2016*, 13. [\[CrossRef\]](#)
105. Laoui, D.; Movahedi, K.; Van Overmeire, E.; Van den Bossche, J.; Schouppe, E.; Mommer, C.; Nikolaou, A.; Morias, Y.; De Baetselier, P.; Van Ginderachter, J.A. Tumor-associated macrophages in breast cancer: Distinct subsets, distinct functions. *Int. J. Dev. Biol.* **2011**, *55*, 861–867. [\[CrossRef\]](#)
106. Zhang, X.; Zeng, Y.; Qu, Q.; Zhu, J.; Liu, Z.; Ning, W.; Zeng, H.; Zhang, N.; Du, W.; Chen, C.; et al. PD-L1 induced by IFN-gamma from tumor-associated macrophages via the JAK/STAT3 and PI3K/AKT signaling pathways promoted progression of lung cancer. *Int. J. Clin. Oncol.* **2017**, *22*, 1026–1033. [\[CrossRef\]](#)
107. Santarpia, M.; González-Cao, M.; Viteri, S.; Karachaliou, N.; Altavilla, G.; Rosell, R. Programmed cell death protein-1/programmed cell death ligand-1 pathway inhibition and predictive biomarkers: Understanding transforming growth factor-beta role. *Transl. Lung Cancer Res.* **2015**, *4*, 728–742.
108. Park, I.H.; Yang, H.N.; Lee, K.J.; Kim, T.-S.; Lee, E.S.; Jung, S.-Y.; Kwon, Y.; Kong, S.-Y. Tumor-derived IL-18 induces PD-1 expression on immunosuppressive NK cells in triple-negative breast cancer. *Oncotarget* **2017**, *8*, 32722–32730. [\[CrossRef\]](#)
109. Treffers, L.W.; Hiemstra, I.H.; Kuijpers, T.W.; van den Berg, T.K.; Matlung, H.L. Neutrophils in cancer. *Immunol. Rev.* **2016**, *273*, 312–328. [\[CrossRef\]](#)
110. Kerfoot, S.M.; Raharjo, E.; Ho, M.; Kaur, J.; Serirom, S.; McCafferty, D.M.; Burns, A.R.; Patel, K.D.; Kubes, P. Exclusive neutrophil recruitment with oncostatin M in a human system. *Am. J. Pathol.* **2001**, *159*, 1531–1539. [\[CrossRef\]](#)
111. Modur, V.; Feldhaus, M.J.; Weyrich, A.S.; Jicha, D.L.; Prescott, S.M.; Zimmerman, G.A.; McIntyre, T.M. Oncostatin M is a proinflammatory mediator. In vivo effects correlate with endothelial cell expression of inflammatory cytokines and adhesion molecules. *J. Clin. Invest.* **1997**, *100*, 158–168. [\[CrossRef\]](#)
112. Queen, M.M.; Ryan, R.E.; Holzer, R.G.; Keller-Peck, C.R.; Jorcyk, C.L. Breast Cancer Cells Stimulate Neutrophils to Produce Oncostatin M: Potential Implications for Tumor Progression. *Cancer Res.* **2005**, *65*, 8896. [\[CrossRef\]](#)
113. Dumitru, C.A.; Bankfalvi, A.; Gu, X.; Eberhardt, W.E.; Zeidler, R.; Lang, S.; Brandau, S. Neutrophils Activate Tumoral CORTACTIN to Enhance Progression of Oropharyngeal Carcinoma. *Front. Immunol.* **2013**, *4*, 33. [\[CrossRef\]](#)
114. Iwase, T.; Sangai, T.; Sakakibara, M.; Sakakibara, J.; Ishigami, E.; Hayama, S.; Nakagawa, A.; Masuda, T.; Tabe, S.; Nagashima, T. An increased neutrophil-to-lymphocyte ratio predicts poorer survival following recurrence for patients with breast cancer. *Mol. Clin. Oncol.* **2017**, *6*, 266–270. [\[CrossRef\]](#)
115. Soto-Perez-de-Celis, E.; Chavarri-Guerra, Y.; Leon-Rodriguez, E.; Gamboa-Dominguez, A. Tumor-Associated Neutrophils in Breast Cancer Subtypes. *Asian Pac. J. Cancer Prev. APJCP* **2017**, *18*, 2689–2694.
116. Jovanovic, B.; Beeler, J.S.; Pickup, M.W.; Chytil, A.; Gorska, A.E.; Ashby, W.J.; Lehmann, B.D.; Zijlstra, A.; Pietenpol, J.A.; Moses, H.L. Transforming growth factor beta receptor type III is a tumor promoter in mesenchymal-stem like triple negative breast cancer. *Breast Cancer Res.* **2014**, *16*, R69. [\[CrossRef\]](#)
117. Fridlender, Z.G.; Sun, J.; Kim, S.; Kapoor, V.; Cheng, G.; Ling, L.; Worthen, G.S.; Albelda, S.M. Polarization of tumor-associated neutrophil phenotype by TGF-beta: “N1” versus “N2” TAN. *Cancer Cell* **2009**, *16*, 183–194. [\[CrossRef\]](#)
118. Lagraoui, M.; Gagnon, L. Enhancement of human neutrophil survival and activation by TGF-beta 1. *Cell. Mol. Biol. (Noisy-le-grand)* **1997**, *43*, 313–318.
119. Perret, J.; McDonald, C.; Apostolopoulos, V. Elevated serum interleukin-5 levels in severe chronic obstructive pulmonary disease. *Acta Biochim. Biophys. Sin.* **2017**, *49*, 560–563. [\[CrossRef\]](#)
120. Filippone, R.T.; Sahakian, L.; Apostolopoulos, V.; Nurgali, K. Eosinophils in Inflammatory Bowel Disease. *Inflamm. Bowel Dis.* **2019**, *25*, 1140–1151. [\[CrossRef\]](#)
121. Gouon-Evans, V.; Rothenberg, M.E.; Pollard, J.W. Postnatal mammary gland development requires macrophages and eosinophils. *Development* **2000**, *127*, 2269–2282.
122. Varricchi, G.; Galdiero, M.R.; Loffredo, S.; Lucarini, V.; Marone, G.; Mattei, F.; Marone, G.; Schiavoni, G. Eosinophils: The unsung heroes in cancer? *Oncoimmunology* **2018**, *7*, e1393134. [\[CrossRef\]](#)
123. Sakkal, S.; Miller, S.; Apostolopoulos, V.; Nurgali, K. Eosinophils in Cancer: Favourable or Unfavourable? *Curr. Med. Chem.* **2016**, *23*, 650–666. [\[CrossRef\]](#)

124. Szalayova, G.; Ogronnik, A.; Spencer, B.; Wade, J.; Bunn, J.; Ambaye, A.; James, T.; Rincon, M. Human breast cancer biopsies induce eosinophil recruitment and enhance adjacent cancer cell proliferation. *Breast Cancer Res. Treat.* **2016**, *157*, 461–474. [[CrossRef](#)]
125. Amini, R.-M.; Aaltonen, K.; Nevanlinna, H.; Carvalho, R.; Salonen, L.; Heikkilä, P.; Blomqvist, C. Mast cells and eosinophils in invasive breast carcinoma. *BMC Cancer* **2007**, *7*, 165. [[CrossRef](#)]
126. Kong, Y.Y.; Fuchsberger, M.; Xiang, S.D.; Apostolopoulos, V.; Plebanski, M. Myeloid derived suppressor cells and their role in diseases. *Curr. Med. Chem.* **2013**, *20*, 1437–1444. [[CrossRef](#)]
127. Tsai, C.S.; Chen, F.H.; Wang, C.C.; Huang, H.L.; Jung, S.M.; Wu, C.J.; Lee, C.C.; McBride, W.H.; Chiang, C.S.; Hong, J.H. Macrophages from irradiated tumors express higher levels of iNOS, arginase-I and COX-2, and promote tumor growth. *Int. J. Radiat. Oncol. Biol. Phys.* **2007**, *68*, 499–507. [[CrossRef](#)]
128. Ochoa, A.C.; Zea, A.H.; Hernandez, C.; Rodriguez, P.C. Arginase, prostaglandins, and myeloid-derived suppressor cells in renal cell carcinoma. *Clin. Cancer Res.* **2007**, *13*, 721s–726s. [[CrossRef](#)]
129. Toor, S.M.; Syed Khaja, A.S.; El Salhat, H.; Faour, I.; Kanbar, J.; Quadri, A.A.; Albashir, M.; Elkord, E. Myeloid cells in circulation and tumor microenvironment of breast cancer patients. *Cancer Immunol. Immunother.* **2017**, *66*, 753–764. [[CrossRef](#)]
130. Law, A.M.K.; Lim, E.; Ormandy, C.J.; Gallego-Ortega, D. The innate and adaptive infiltrating immune systems as targets for breast cancer immunotherapy. *Endocr. Relat. Cancer* **2017**, *24*, R123–R144. [[CrossRef](#)]
131. Bu, X.; Yao, Y.; Li, X. Immune Checkpoint Blockade in Breast Cancer Therapy. *Adv. Exp. Med. Biol.* **2017**, *1026*, 383–402.
132. Walker, L.S.; Sansom, D.M. Confusing signals: Recent progress in CTLA-4 biology. *Trends Immunol.* **2015**, *36*, 63–70. [[CrossRef](#)]
133. Seidel, J.A.; Otsuka, A.; Kabashima, K. Anti-PD-1 and Anti-CTLA-4 Therapies in Cancer: Mechanisms of Action, Efficacy, and Limitations. *Front. Oncol.* **2018**, *8*, 86. [[CrossRef](#)]
134. Yu, H.; Yang, J.; Jiao, S.; Li, Y.; Zhang, W.; Wang, J. Cytotoxic T lymphocyte antigen 4 expression in human breast cancer: Implications for prognosis. *Cancer Immunol. Immunother.* **2015**, *64*, 853–860. [[CrossRef](#)]
135. Hartkopf, A.D.; Taran, F.A.; Wallwiener, M.; Walter, C.B.; Krämer, B.; Grischke, E.M.; Brucker, S.Y. PD-1 and PD-L1 Immune Checkpoint Blockade to Treat Breast Cancer. *Breast Care* **2016**, *11*, 385–390. [[CrossRef](#)]
136. Ma, W.; Gilligan, B.M.; Yuan, J.; Li, T. Current status and perspectives in translational biomarker research for PD-1/PD-L1 immune checkpoint blockade therapy. *J. Hematol. Oncol.* **2016**, *9*, 47. [[CrossRef](#)]
137. Ren, X.; Wu, H.; Lu, J.; Zhang, Y.; Luo, Y.; Xu, Q.; Shen, S.; Liang, Z. PD1 protein expression in tumor infiltrated lymphocytes rather than PDL1 in tumor cells predicts survival in triple-negative breast cancer. *Cancer Biol. Ther.* **2018**, *19*, 373–380. [[CrossRef](#)]
138. Bertucci, F.; Finetti, P.; Birnbaum, D.; Mameessier, E. The PD1/PDL1 axis, a promising therapeutic target in aggressive breast cancers. *Oncoimmunology* **2016**, *5*, e1085148. [[CrossRef](#)]
139. Mellor, A.L.; Munn, D.H. Immunology at the maternal-fetal interface: Lessons for T cell tolerance and suppression. *Annu. Rev. Immunol.* **2000**, *18*, 367–391. [[CrossRef](#)]
140. Moon, Y.W.; Hajjar, J.; Hwu, P.; Naing, A. Targeting the indoleamine 2,3-dioxygenase pathway in cancer. *J. Immunother. Cancer* **2015**, *3*, 51. [[CrossRef](#)]
141. Chuang, H.C.; Lan, J.L.; Chen, D.Y.; Yang, C.Y.; Chen, Y.M.; Li, J.P.; Huang, C.Y.; Liu, P.E.; Wang, X.; Tan, T.H. The kinase GLK controls autoimmunity and NF-kappaB signaling by activating the kinase PKC-theta in T cells. *Nat. Immunol.* **2011**, *12*, 1113–1118. [[CrossRef](#)]
142. Metz, R.; Rust, S.; Duhadaway, J.B.; Mautino, M.R.; Munn, D.H.; Vahanian, N.N.; Link, C.J.; Prendergast, G.C. IDO inhibits a tryptophan sufficiency signal that stimulates mTOR: A novel IDO effector pathway targeted by D-1-methyl-tryptophan. *Oncoimmunology* **2012**, *1*, 1460–1468. [[CrossRef](#)]
143. Harding, H.P.; Novoa, I.; Zhang, Y.; Zeng, H.; Wek, R.; Schapira, M.; Ron, D. Regulated translation initiation controls stress-induced gene expression in mammalian cells. *Mol. Cell* **2000**, *6*, 1099–1108. [[CrossRef](#)]
144. Sundrud, M.S.; Koralov, S.B.; Feuerer, M.; Calado, D.P.; Kozhaya, A.E.; Rhule-Smith, A.; Lefebvre, R.E.; Unutmaz, D.; Mazitschek, R.; Waldner, H.; et al. Halofuginone inhibits TH17 cell differentiation by activating the amino acid starvation response. *Science* **2009**, *324*, 1334–1338. [[CrossRef](#)]
145. Munn, D.H.; Sharma, M.D.; Baban, B.; Harding, H.P.; Zhang, Y.; Ron, D.; Mellor, A.L. GCN2 kinase in T cells mediates proliferative arrest and anergy induction in response to indoleamine 2,3-dioxygenase. *Immunity* **2005**, *22*, 633–642. [[CrossRef](#)]

146. Nguyen, N.T.; Kimura, A.; Nakahama, T.; Chinen, I.; Masuda, K.; Nohara, K.; Fujii-Kuriyama, Y.; Kishimoto, T. Aryl hydrocarbon receptor negatively regulates dendritic cell immunogenicity via a kynurenine-dependent mechanism. *Proc. Natl. Acad. Sci. USA* **2010**, *107*, 19961–19966. [\[CrossRef\]](#)
147. Opitz, C.A.; Litzenburger, U.M.; Sahm, F.; Ott, M.; Tritschler, I.; Trump, S.; Schumacher, T.; Jestaedt, L.; Schrenk, D.; Weller, M.; et al. An endogenous tumour-promoting ligand of the human aryl hydrocarbon receptor. *Nature* **2011**, *478*, 197–203. [\[CrossRef\]](#)
148. Isla Larrain, M.T.; Rabassa, M.E.; Lacunza, E.; Barbera, A.; Cretón, A.; Segal-Eiras, A.; Croce, M.V. IDO is highly expressed in breast cancer and breast cancer-derived circulating microvesicles and associated to aggressive types of tumors by in silico analysis. *Tumor Biol.* **2014**, *35*, 6511–6519. [\[CrossRef\]](#)
149. Asghar, K.; Loya, A.; Rana, I.A.; Tahseen, M.; Ishaq, M.; Farooq, A.; Bakar, M.A.; Masood, I. Indoleamine 2,3-dioxygenase expression and overall survival in patients diagnosed with breast cancer in Pakistan. *Cancer Manag. Res.* **2018**, *11*, 475–481. [\[CrossRef\]](#)
150. Dill, E.A.; Dillon, P.M.; Bullock, T.N.; Mills, A.M. IDO expression in breast cancer: An assessment of 281 primary and metastatic cases with comparison to PD-L1. *Mod. Pathol.* **2018**, *31*, 1513–1522. [\[CrossRef\]](#)
151. Fraschilla, I.; Pillai, S. Viewing Siglecs through the lens of tumor immunology. *Immunol. Rev.* **2017**, *276*, 178–191. [\[CrossRef\]](#)
152. Tanida, S.; Akita, K.; Ishida, A.; Mori, Y.; Toda, M.; Inoue, M.; Ohta, M.; Yashiro, M.; Sawada, T.; Hirakawa, K.; et al. Binding of the sialic acid-binding lectin, Siglec-9, to the membrane mucin, MUC1, induces recruitment of beta-catenin and subsequent cell growth. *J. Biol. Chem.* **2013**, *288*, 31842–31852. [\[CrossRef\]](#)
153. Singh, P.K.; Hollingsworth, M.A. Cell surface-associated mucins in signal transduction. *Trends Cell Biol.* **2006**, *16*, 467–476. [\[CrossRef\]](#)
154. Workman, H.C.; Sweeney, C.; Carraway, K.L., 3rd. The membrane mucin Muc4 inhibits apoptosis induced by multiple insults via ErbB2-dependent and ErbB2-independent mechanisms. *Cancer Res.* **2009**, *69*, 2845–2852. [\[CrossRef\]](#)
155. Gong, J.; Apostolopoulos, V.; Chen, D.; Chen, H.; Koido, S.; Gendler, S.J.; McKenzie, I.F.; Kufe, D. Selection and characterization of MUC1-specific CD8+ T cells from MUC1 transgenic mice immunized with dendritic-carcinoma fusion cells. *Immunology* **2000**, *101*, 316–324. [\[CrossRef\]](#)
156. Koido, S.; Enomoto, Y.; Apostolopoulos, V.; Gong, J. Tumor regression by CD4 T-cells primed with dendritic/tumor fusion cell vaccines. *Anticancer Res.* **2014**, *34*, 3917–3924.
157. Pietersz, G.A.; Li, W.; Popovski, V.; Caruana, J.A.; Apostolopoulos, V.; McKenzie, I.F. Parameters for using mannan-MUC1 fusion protein to induce cellular immunity. *Cancer Immunol. Immunother. CII* **1998**, *45*, 321–326. [\[CrossRef\]](#)
158. Pietersz, G.A.; Pouniotis, D.S.; Apostolopoulos, V. Design of peptide-based vaccines for cancer. *Curr. Med. Chem.* **2006**, *13*, 1591–1607. [\[CrossRef\]](#)
159. Son, H.Y.; Apostolopoulos, V.; Kim, C.W. T/Tn immunotherapy avoiding immune deviation. *Int. J. Immunopathol. Pharmacol.* **2016**, *29*, 812–817. [\[CrossRef\]](#)
160. Son, H.Y.; Apostolopoulos, V.; Kim, C.W. Mannosylated T/Tn with Freund's adjuvant induces cellular immunity. *Int. J. Immunopathol. Pharmacol.* **2018**, *31*, 394632017742504. [\[CrossRef\]](#)
161. Tang, C.K.; Katsara, M.; Apostolopoulos, V. Strategies used for MUC1 immunotherapy: Human clinical studies. *Expert Rev. Vaccines* **2008**, *7*, 963–975. [\[CrossRef\]](#)
162. Macao, B.; Johansson, D.G.; Hansson, G.C.; Hard, T. Autoproteolysis coupled to protein folding in the SEA domain of the membrane-bound MUC1 mucin. *Nat. Struct. Mol. Biol.* **2006**, *13*, 71–76. [\[CrossRef\]](#)
163. Kufe, D.W. MUC1-C oncoprotein as a target in breast cancer: Activation of signaling pathways and therapeutic approaches. *Oncogene* **2013**, *32*, 1073–1081. [\[CrossRef\]](#)
164. Huang, L.; Chen, D.; Liu, D.; Yin, L.; Kharbanda, S.; Kufe, D. MUC1 oncoprotein blocks glycogen synthase kinase 3beta-mediated phosphorylation and degradation of beta-catenin. *Cancer Res.* **2005**, *65*, 10413–10422. [\[CrossRef\]](#)
165. Jing, X.; Liang, H.; Hao, C.; Yang, X.; Cui, X. Overexpression of MUC1 predicts poor prognosis in patients with breast cancer. *Oncol. Rep.* **2019**, *41*, 801–810. [\[CrossRef\]](#)
166. Dong, R.; Wang, Q.; He, X.L.; Chu, Y.K.; Lu, J.G.; Ma, Q.J. Role of nuclear factor kappa B and reactive oxygen species in the tumor necrosis factor- α -induced epithelial-mesenchymal transition of MCF-7 cells. *Braz. J. Med. Biol. Res.* **2007**, *40*, 1071–1078. [\[CrossRef\]](#)

167. Martin, F.T.; Dwyer, R.M.; Kelly, J.; Khan, S.; Murphy, J.M.; Curran, C.; Miller, N.; Hennessy, E.; Dockery, P.; Barry, F.P.; et al. Potential role of mesenchymal stem cells (MSCs) in the breast tumour microenvironment: Stimulation of epithelial to mesenchymal transition (EMT). *Breast Cancer Res. Treat.* **2010**, *124*, 317–326. [\[CrossRef\]](#)
168. Sousa, S.; Brion, R.; Lintunen, M.; Kronqvist, P.; Sandholm, J.; Mönkkönen, J.; Kellokumpu-Lehtinen, P.-L.; Lanttia, S.; Tynnenen, O.; Joensuu, H.; et al. Human breast cancer cells educate macrophages toward the M2 activation status. *Breast Cancer Res. BCR* **2015**, *17*, 101. [\[CrossRef\]](#)
169. Medrek, C.; Pontén, F.; Jirstrom, K.; Leandersson, K. The presence of tumor associated macrophages in tumor stroma as a prognostic marker for breast cancer patients. *BMC Cancer* **2012**, *12*, 306. [\[CrossRef\]](#)
170. Brechbuhl, H.M.; Finlay-Schultz, J.; Yamamoto, T.M.; Gillen, A.E.; Cittelly, D.M.; Tan, A.-C.; Sams, S.B.; Pillai, M.M.; Elias, A.D.; Robinson, W.A.; et al. Fibroblast Subtypes Regulate Responsiveness of Luminal Breast Cancer to Estrogen. *Clin. Cancer Res.* **2017**, *23*, 1710–1721. [\[CrossRef\]](#)
171. Costa, A.; Kieffer, Y.; Scholer-Dahirel, A.; Pelon, F.; Bourachot, B.; Cardon, M.; Sirven, P.; Magagna, I.; Fuhrmann, L.; Bernard, C.; et al. Fibroblast Heterogeneity and Immunosuppressive Environment in Human Breast Cancer. *Cancer Cell* **2018**, *33*, 463–479. [\[CrossRef\]](#)
172. Liu, Q.; Li, A.; Tian, Y.; Wu, J.D.; Liu, Y.; Li, T.; Chen, Y.; Han, X.; Wu, K. The CXCL8-CXCR1/2 pathways in cancer. *Cytokine Growth Factor Rev.* **2016**, *31*, 61–71. [\[CrossRef\]](#)
173. Ownby, H.E.; Roi, L.D.; Isenberg, R.R.; Brennan, M.J. Peripheral lymphocyte and eosinophil counts as indicators of prognosis in primary breast cancer. *Cancer* **1983**, *52*, 126–130. [\[CrossRef\]](#)
174. Diaz-Montero, C.M.; Salem, M.L.; Nishimura, M.I.; Garrett-Mayer, E.; Cole, D.J.; Montero, A.J. Increased circulating myeloid-derived suppressor cells correlate with clinical cancer stage, metastatic tumor burden, and doxorubicin-cyclophosphamide chemotherapy. *Cancer Immunol. Immunother. CII* **2009**, *58*, 49–59. [\[CrossRef\]](#)
175. Solito, S.; Falisi, E.; Diaz-Montero, C.M.; Doni, A.; Pinton, L.; Rosato, A.; Francescato, S.; Basso, G.; Zanovello, P.; Onicescu, G.; et al. A human promyelocytic-like population is responsible for the immune suppression mediated by myeloid-derived suppressor cells. *Blood* **2011**, *118*, 2254–2265. [\[CrossRef\]](#)
176. Yu, J.; Du, W.; Yan, F.; Wang, Y.; Li, H.; Cao, S.; Yu, W.; Shen, C.; Liu, J.; Ren, X. Myeloid-Derived Suppressor Cells Suppress Antitumor Immune Responses through IDO Expression and Correlate with Lymph Node Metastasis in Patients with Breast Cancer. *J. Immunol.* **2013**, *190*, 3783. [\[CrossRef\]](#)
177. Holmgaard, R.B.; Zamarin, D.; Munn, D.H.; Wolchok, J.D.; Allison, J.P. Indoleamine 2,3-dioxygenase is a critical resistance mechanism in antitumor T cell immunotherapy targeting CTLA-4. *J. Exp. Med.* **2013**, *210*, 1389. [\[CrossRef\]](#)
178. Chen, X.; Shao, Q.; Hao, S.; Zhao, Z.; Wang, Y.; Guo, X.; He, Y.; Gao, W.; Mao, H. CTLA-4 positive breast cancer cells suppress dendritic cells maturation and function. *Oncotarget* **2017**, *8*, 13703–13715. [\[CrossRef\]](#)
179. Ye, Q.; Wang, C.; Xian, J.; Zhang, M.; Cao, Y.; Cao, Y. Expression of programmed cell death protein 1 (PD-1) and indoleamine 2,3-dioxygenase (IDO) in the tumor microenvironment and in tumor-draining lymph nodes of breast cancer. *Hum. Pathol.* **2018**, *75*, 81–90. [\[CrossRef\]](#)
180. Kim, S.; Park, S.; Cho, M.S.; Lim, W.; Moon, B.-I.; Sung, S.H. Strong Correlation of Indoleamine 2,3-Dioxygenase 1 Expression with Basal-Like Phenotype and Increased Lymphocytic Infiltration in Triple-Negative Breast Cancer. *J. Cancer* **2017**, *8*, 124–130. [\[CrossRef\]](#)
181. Tomioka, Y.; Morimatsu, M.; Nishijima, K.-I.; Usui, T.; Yamamoto, S.; Suyama, H.; Ozaki, K.; Ito, T.; Ono, E. A soluble form of Siglec-9 provides an antitumor benefit against mammary tumor cells expressing MUC1 in transgenic mice. *Biochem. Biophys. Res. Commun.* **2014**, *450*, 532–537. [\[CrossRef\]](#)
182. Ohta, M.; Ishida, A.; Toda, M.; Akita, K.; Inoue, M.; Yamashita, K.; Watanabe, M.; Murata, T.; Usui, T.; Nakada, H. Immunomodulation of monocyte-derived dendritic cells through ligation of tumor-produced mucins to Siglec-9. *Biochem. Biophys. Res. Commun.* **2010**, *402*, 663–669. [\[CrossRef\]](#)
183. Segovia-Mendoza, M.; Morales-Montor, J. Immune Tumor Microenvironment in Breast Cancer and the Participation of Estrogens and Its Receptors into Cancer Physiopathology. *Front. Immunol.* **2019**, *10*, 348. [\[CrossRef\]](#)

