



**VICTORIA UNIVERSITY**  
MELBOURNE AUSTRALIA

*Further inequalities for log-convex functions related to Hermite-Hadamard result*

This is the Published version of the following publication

Dragomir, Sever S (2019) Further inequalities for log-convex functions related to Hermite-Hadamard result. *Proyecciones*, 38 (2). pp. 267-293. ISSN 0716-0917

The publisher's official version can be found at  
<https://scielo.conicyt.cl/pdf/proy/v38n2/0717-6279-proy-38-02-00267.pdf>  
Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/39887/>

Proyecciones Journal of Mathematics  
Vol. 38, N° 2, pp. 267-293, June 2019.  
Universidad Católica del Norte  
Antofagasta - Chile

## Further inequalities for log-convex functions related to Hermite-Hadamard result

*S. S. Dragomir*

*Victoria University, Australia*

*Received : July 2017. Accepted : December 2018*

### Abstract

*Some unweighted and weighted inequalities of Hermite-Hadamard type for log-convex functions defined on real intervals are given.*

**Subjclass:** *26D15; 25D10.*

**Keywords:** *Convex functions, Integral inequalities, Log-Convex functions.*

## 1. Introduction

A function  $f : I \rightarrow (0, \infty)$  is said to be *log-convex* or *multiplicatively convex* if  $\log f$  is convex, or, equivalently, if for all  $x, y \in I$  and  $t \in [0, 1]$  one has the inequality:

$$(1.1) \quad f(tx + (1-t)y) \leq [f(x)]^t [f(y)]^{1-t}.$$

We note that if  $f$  and  $g$  are convex and  $g$  is increasing, then  $g \circ f$  is convex; moreover, since  $f = \exp(\log f)$ , it follows that a log-convex function is convex, but the converse may not necessarily be true. This follows directly from (1.1) because, by the *arithmetic-geometric mean inequality*, we have

$$[f(x)]^t [f(y)]^{1-t} \leq tf(x) + (1-t)f(y)$$

for all  $x, y \in I$  and  $t \in [0, 1]$ .

Let us recall the *Hermite-Hadamard inequality*

$$(1.2) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2},$$

where  $f : I \subseteq \mathbf{R} \rightarrow \mathbf{R}$  is a convex function on the interval  $I$ ,  $a, b \in I$  and  $a < b$ .

For related results, see [1]-[22], [25]-[29], [30]-[40] and [41]-[53].

Note that if we apply the above inequality for the log-convex functions  $f : I \rightarrow (0, \infty)$ , we have that

$$(1.3) \quad \ln \left[ f\left(\frac{a+b}{2}\right) \right] \leq \frac{1}{b-a} \int_a^b \ln f(x) dx \leq \frac{\ln f(a) + \ln f(b)}{2},$$

from which we get

$$(1.4) \quad f\left(\frac{a+b}{2}\right) \leq \exp \left[ \frac{1}{b-a} \int_a^b \ln f(x) dx \right] \leq \sqrt{f(a)f(b)}$$

that is an inequality of Hermite-Hadamard's type for log-convex functions.

By using simple properties of log-convex functions Dragomir and Mond proved in 1998 the following result [32].

**Theorem 1.** *Let  $f : I \rightarrow (0, \infty)$  be a log-convex mapping on  $I$  and  $a, b \in I$  with  $a < b$ . Then one has the inequality:*

$$(1.5) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b \sqrt{f(x)f(a+b-x)} dx \leq \sqrt{f(a)f(b)}.$$

The inequality between the first and second term in (1.5) may be improved as follows [32]. A different upper bound for the middle term in (1.5) can be also provided.

**Theorem 2.** *Let  $f : I \rightarrow (0, \infty)$  be a log-convex mapping on  $I$  and  $a, b \in I$  with  $a < b$ . Then one has the inequalities:*

$$(1.6) \quad \begin{aligned} f\left(\frac{a+b}{2}\right) &\leq \exp\left[\frac{1}{b-a} \int_a^b \ln f(x) dx\right] \\ &\leq \frac{1}{b-a} \int_a^b \sqrt{f(x)f(a+b-x)} dx \\ &\leq \frac{1}{b-a} \int_a^b f(x) dx \leq L(f(a), f(b)), \end{aligned}$$

where  $L(p, q)$  is the logarithmic mean of the strictly positive real numbers  $p, q$ , i.e.,

$$L(p, q) := \frac{p - q}{\ln p - \ln q} \text{ if } p \neq q \text{ and } L(p, p) := p.$$

The last inequality in (1.6) was obtained in a different context in [43]. As shown in [59], the following result also holds:

**Theorem 3.** *Let  $f : I \rightarrow (0, \infty)$  be a log-convex mapping on  $I$  and  $a, b \in I$  with  $a < b$ . Then one has the inequalities:*

$$(1.7) \quad f\left(\frac{a+b}{2}\right) \leq \left(\frac{1}{b-a} \int_a^b \sqrt{f(x)} dx\right)^2 \leq \frac{1}{b-a} \int_a^b f(x) dx.$$

We define the  $p$ -logarithmic mean as

$$L_p(a, b) := \begin{cases} \left[\frac{b^{p+1} - a^{p+1}}{(p+1)(b-a)}\right]^{\frac{1}{p}}, & \text{with } a \neq b \\ a, & \text{if } a = b \end{cases}$$

for  $p \neq 0, -1$  and  $a, b > 0$ .

In the recent work [28] we generalized the inequality (1.6) as follows:

**Theorem 4.** Let  $f : [a, b] \rightarrow (0, \infty)$  be a log-convex function on  $[a, b]$ . Then for any  $p > 0$  we have the inequality

$$\begin{aligned}
 (1.8) \quad f\left(\frac{a+b}{2}\right) &\leq \exp\left[\frac{1}{b-a} \int_a^b \ln f(x) dx\right] \\
 &\leq \left(\frac{1}{b-a} \int_a^b f^p(x) f^p(a+b-x) dx\right)^{\frac{1}{2p}} \\
 &\leq \left(\frac{1}{b-a} \int_a^b f^{2p}(x) dx\right)^{\frac{1}{2p}} \\
 &\leq \begin{cases} [L_{2p-1}(f(a), f(b))]^{1-\frac{1}{2p}} [L(f(a), f(b))]^{\frac{1}{2p}}, & p \neq \frac{1}{2}; \\ L(f(a), f(b)), & p = \frac{1}{2}. \end{cases}
 \end{aligned}$$

If  $p \in \left(0, \frac{1}{2}\right)$ , then we have

$$\begin{aligned}
 (1.9) \quad f\left(\frac{a+b}{2}\right) &\leq \exp\left[\frac{1}{b-a} \int_a^b \ln f(x) dx\right] \\
 &\leq \left(\frac{1}{b-a} \int_a^b f^p(x) f^p(a+b-x) dx\right)^{\frac{1}{2p}} \\
 &\leq \left(\frac{1}{b-a} \int_a^b f^{2p}(x) dx\right)^{\frac{1}{2p}} \leq \frac{1}{b-a} \int_a^b f(x) dx.
 \end{aligned}$$

**Remark 1.** If we take in (1.8)  $p = 1$ , then we get

$$\begin{aligned}
 (1.10) \quad f\left(\frac{a+b}{2}\right) &\leq \exp\left[\frac{1}{b-a} \int_a^b \ln f(x) dx\right] \\
 &\leq \left(\frac{1}{b-a} \int_a^b f(x) f(a+b-x) dx\right)^{\frac{1}{2}} \\
 &\leq \left(\frac{1}{b-a} \int_a^b f^2(x) dx\right)^{\frac{1}{2}} \\
 &\leq [A(f(a), f(b))]^{\frac{1}{2}} [L(f(a), f(b))]^{\frac{1}{2}}.
 \end{aligned}$$

If we take  $p = \frac{1}{4}$  in (1.9), then we get

$$\begin{aligned}
 (1.11) \quad f\left(\frac{a+b}{2}\right) &\leq \exp\left[\frac{1}{b-a} \int_a^b \ln f(x) dx\right] \\
 &\leq \left(\frac{1}{b-a} \int_a^b \sqrt[4]{f(x)f(a+b-x)} dx\right)^2 \\
 &\leq \left(\frac{1}{b-a} \int_a^b \sqrt{f(x)} dx\right)^2 \leq \frac{1}{b-a} \int_a^b f(x) dx.
 \end{aligned}$$

This improves the inequality (1.7).

Motivated by the above results, we establish in this paper some new un-weighted and weighted inequalities for log-convex functions, some of them improving earlier results. Applications for special means are also provided.

### 2. New Results

The following result holds.

**Theorem 1.** *Let  $f : [a, b] \rightarrow (0, \infty)$  be a log-convex function. Then for every  $t \in [0, 1]$  we have*

$$\begin{aligned}
 (2.1) \quad \int_a^b f(x) dx &\geq \int_a^b [f(x)]^{1-t} [f(a+b-x)]^t dx \\
 &\geq \begin{cases} \frac{1}{1-2t} \int_{(1-t)a+tb}^{ta+(1-t)b} f(u) du, & \text{if } t \neq \frac{1}{2} \\ (b-a) f\left(\frac{a+b}{2}\right), & \text{if } t = \frac{1}{2}. \end{cases}
 \end{aligned}$$

**Proof.** The cases  $t = 0, \frac{1}{2}, 1$  are obvious.

Assume that  $t \in (0, 1) \setminus \left\{\frac{1}{2}\right\}$ . By the log-convexity of  $f$  we have

$$\begin{aligned}
 (2.2) \quad [f(x)]^{1-t} [f(a+b-x)]^t &\geq f((1-t)x + t(a+b-x)) \\
 &= f((1-2t)x + t(a+b))
 \end{aligned}$$

for any  $x \in [a, b]$ .

Integrating the inequality (2.2) over  $x$  on  $[a, b]$  we have

$$\int_a^b [f(x)]^{1-t} [f(a+b-x)]^t dx \geq \int_a^b f((1-2t)x + t(a+b)) dx.$$

Since  $t \neq \frac{1}{2}$ , then  $u := (1-2t)x + t(a+b)$  is a change of variable with  $du = (1-2t) dx$ .

For  $x = a$  we get  $u = (1-t)a + tb$  and for  $x = b$  we get  $u = ta + (1-t)b$ . Therefore

$$\int_a^b f((1-2t)x + t(a+b)) dx = \frac{1}{1-2t} \int_{(1-t)a+tb}^{ta+(1-t)b} f(u) du$$

and the second inequality in (2.1) is proved.

By the Hölder integral inequality for  $p = \frac{1}{1-t}$ ,  $q = \frac{1}{t}$  we have

$$\begin{aligned} & \int_a^b [f(x)]^{1-t} [f(a+b-x)]^t dx \\ & \leq \left( \int_a^b ([f(x)]^{1-t})^{\frac{1}{1-t}} dx \right)^{1-t} \left( \int_a^b ([f(a+b-x)]^t)^{\frac{1}{t}} dx \right)^t \\ & = \left( \int_a^b f(x) dx \right)^{1-t} \left( \int_a^b f(a+b-x) dx \right)^t \\ & = \left( \int_a^b f(x) dx \right)^{1-t} \left( \int_a^b f(x) dx \right)^t = \int_a^b f(x) dx, \end{aligned}$$

that proves the first inequality in (2.1).  $\square$

**Corollary 1.** *Let  $f : [a, b] \rightarrow (0, \infty)$  be a log-convex function. Then for every  $t \in [0, 1] \setminus \left\{ \frac{1}{2} \right\}$  we have for  $p > 0$  that*

$$(2.3) \quad \left( \frac{1}{b-a} \int_a^b f^{2p}(x) dx \right)^{\frac{1}{2p}}$$

$$\begin{aligned}
 &\geq \left( \frac{1}{b-a} \int_a^b [f^{2p}(x)]^{1-t} [f^{2p}(a+b-x)]^t dx \right)^{\frac{1}{2p}} \\
 &\geq \left( \frac{1}{(1-2t)(b-a)} \int_{(1-t)a+tb}^{ta+(1-t)b} f^{2p}(u) du \right)^{\frac{1}{2p}} \\
 &\geq \left( \frac{1}{(1-2t)(b-a)} \int_{(1-t)a+tb}^{ta+(1-t)b} f^p(u) f^p(a+b-u) dx \right)^{\frac{1}{2p}} \\
 &\geq \exp \left[ \frac{1}{(1-2t)(b-a)} \int_{(1-t)a+tb}^{ta+(1-t)b} \ln f(u) du \right] \geq f\left(\frac{a+b}{2}\right).
 \end{aligned}$$

**Proof.** Follows from Theorem 5 applied for the log-convex function  $f^{2p}$  with  $p > 0$  and by Theorem 4 applied for the interval  $[(1-t)a+tb, ta+(1-t)b]$  when  $t \in (0, \frac{1}{2})$  or  $[ta+(1-t)b, (1-t)a+tb]$  when  $t \in (\frac{1}{2}, 1)$ .  $\square$

If we take  $p = 1$  in (2.3), then we get

$$\left( \frac{1}{b-a} \int_a^b f^2(x) dx \right)^{\frac{1}{2}} \geq \left( \frac{1}{b-a} \int_a^b [f^2(x)]^{1-t} [f^2(a+b-x)]^t dx \right)^{\frac{1}{2}} \tag{2.4}$$

$$\begin{aligned}
 &\geq \left( \frac{1}{(1-2t)(b-a)} \int_{(1-t)a+tb}^{ta+(1-t)b} f^2(u) du \right)^{\frac{1}{2}} \\
 &\geq \left( \frac{1}{(1-2t)(b-a)} \int_{(1-t)a+tb}^{ta+(1-t)b} f(u) f(a+b-u) dx \right)^{\frac{1}{2}} \\
 &\geq \exp \left[ \frac{1}{(1-2t)(b-a)} \int_{(1-t)a+tb}^{ta+(1-t)b} \ln f(u) du \right] \geq f\left(\frac{a+b}{2}\right).
 \end{aligned}$$

If we take  $p = \frac{1}{2}$  in (2.3), then we get

$$\left( \frac{1}{b-a} \int_a^b f(x) dx \right) \tag{2.5}$$



$$\begin{aligned}
&\geq \frac{1}{b-a} \int_a^b [f(x)]^{1-t} [f(a+b-x)]^t dx \\
&\geq \frac{1}{(1-2t)(b-a)} \int_{(1-t)a+tb}^{ta+(1-t)b} f(u) du \\
&\geq \frac{1}{(1-2t)(b-a)} \int_{(1-t)a+tb}^{ta+(1-t)b} \sqrt{f(u) f(a+b-u)} dx \\
&\geq \exp \left[ \frac{1}{(1-2t)(b-a)} \int_{(1-t)a+tb}^{ta+(1-t)b} \ln f(u) du \right] \geq f\left(\frac{a+b}{2}\right).
\end{aligned}$$

**Corollary 2.** Let  $f : [a, b] \rightarrow (0, \infty)$  be a log-convex function. Then for every  $t \in [0, 1] \setminus \left\{\frac{1}{2}\right\}$  we have for  $p \in \left(0, \frac{1}{2}\right)$  that

$$\begin{aligned}
(2.6) \quad &\frac{1}{b-a} \int_a^b f(x) dx \\
&\geq \frac{1}{b-a} \int_a^b [f(x)]^{1-t} [f(a+b-x)]^t dx \\
&\geq \frac{1}{(1-2t)(b-a)} \int_{(1-t)a+tb}^{ta+(1-t)b} f(u) du \\
&\geq \left( \frac{1}{(1-2t)(b-a)} \int_{(1-t)a+tb}^{ta+(1-t)b} f^{2p}(u) du \right)^{\frac{1}{2p}} \\
&\geq \left( \frac{1}{(1-2t)(b-a)} \int_{(1-t)a+tb}^{ta+(1-t)b} f^p(u) f^p(a+b-u) du \right)^{\frac{1}{2p}} \\
&\geq \exp \left[ \frac{1}{(1-2t)(b-a)} \int_{(1-t)a+tb}^{ta+(1-t)b} \ln f(u) du \right] \geq f\left(\frac{a+b}{2}\right).
\end{aligned}$$

Follows by Theorem 5 and Theorem 4 for  $p \in \left(0, \frac{1}{2}\right)$ .

If we take  $p = \frac{1}{4}$  in (2.6), then we get

$$(2.7) \quad \frac{1}{b-a} \int_a^b f(x) dx$$

$$\begin{aligned}
 &\geq \frac{1}{b-a} \int_a^b [f(x)]^{1-t} [f(a+b-x)]^t dx \\
 &\geq \frac{1}{(1-2t)(b-a)} \int_{(1-t)a+tb}^{ta+(1-t)b} f(u) du \\
 &\geq \left( \frac{1}{(1-2t)(b-a)} \int_{(1-t)a+tb}^{ta+(1-t)b} \sqrt{f(u)} du \right)^2 \\
 &\geq \left( \frac{1}{(1-2t)(b-a)} \int_{(1-t)a+tb}^{ta+(1-t)b} \sqrt[4]{f(u) f(a+b-u)} du \right)^2 \\
 &\geq \exp \left[ \frac{1}{(1-2t)(b-a)} \int_{(1-t)a+tb}^{ta+(1-t)b} \ln f(u) du \right] \geq f\left(\frac{a+b}{2}\right).
 \end{aligned}$$

If we use the inequality

$$(2.8) \quad \frac{1}{b-a} \int_a^b f(x) dx \geq \frac{1}{(1-2t)(b-a)} \int_{(1-t)a+tb}^{ta+(1-t)b} f(u) du,$$

where  $t \neq \frac{1}{2}$ , for the log-convex function  $f : [a, b] \rightarrow (0, \infty)$ ,  $f(x) = \frac{1}{x}$ , then we get

$$\frac{\ln b - \ln a}{b-a} \geq \frac{\ln(ta + (1-t)b) - \ln((1-t)a + tb)}{(ta + (1-t)b) - ((1-t)a + tb)}$$

which, in terms of logarithmic mean, is

$$(2.9) \quad L(ta + (1-t)b, (1-t)a + tb) \geq L(a, b)$$

for any  $t \in [0, 1] \setminus \left\{\frac{1}{2}\right\}$ .

When  $t = \frac{1}{2}$  the inequality (2.9) becomes  $A(a, b) \geq L(a, b)$  that is also true.

For  $q \neq 0, -1$  define the  $L_q$ -logarithmic mean as

$$L_q(a, b) := \begin{cases} \left(\frac{b^{q+1}-a^{q+1}}{(q+1)(b-a)}\right)^{\frac{1}{q}} & \text{if } b \neq a, \\ a & \text{if } b = a. \end{cases}$$

If we use the inequality (2.8) for the log-convex function  $f : [a, b] \rightarrow (0, \infty)$ ,  $f(x) = \frac{1}{x^p}$ , with  $p > 0, p \neq 1$ , then we have for any  $t \in [0, 1] \setminus \left\{\frac{1}{2}\right\}$  that

$$\frac{1}{b-a} \int_a^b x^{-p} dx \geq \frac{1}{(1-2t)(b-a)} \int_{(1-t)a+tb}^{ta+(1-t)b} u^{-p} du,$$

i.e.

$$\frac{b^{-p+1} - a^{-p+1}}{(1-p)(b-a)} \geq \frac{(ta + (1-t)b)^{-p+1} - ((1-t)a + tb)^{-p+1}}{((1-p)) [(ta + (1-t)b) - ((1-t)a + tb)]},$$

which can be written as

$$L_{-p}^{-p}(a, b) \geq L_{-p}^{-p}(ta + (1-t)b, (1-t)a + tb).$$

Therefore we have

$$(2.10) \quad L_{-p}(ta + (1-t)b, (1-t)a + tb) \geq L_{-p}(a, b)$$

for any  $p > 0$ ,  $p \neq 1$  and for any  $t \in [0, 1] \setminus \left\{\frac{1}{2}\right\}$ . The case  $t = \frac{1}{2}$  reduces to the known inequality  $A(a, b) \geq L_{-p}(a, b)$  for any  $p > 0$ .

### 3. Weighted Inequalities

We have the following generalized weighted version of the inequality (1.5).

**Theorem 1.** Let  $f : [a, b] \rightarrow (0, \infty)$  be a log-convex function. If  $w : [a, b] \rightarrow [0, \infty)$  is integrable and  $\int_a^b w(x) dx > 0$ , then

$$(3.1) \quad f\left(\frac{a+b}{2}\right) \leq \left(\frac{\int_a^b w(x) f^p(x) f^p(a+b-x) dx}{\int_a^b w(x) dx}\right)^{\frac{1}{2p}} \leq \sqrt{f(a) f(b)}$$

for any  $p > 0$ .

In particular, we have

$$(3.2) \quad f\left(\frac{a+b}{2}\right) \leq \left(\frac{\int_a^b w(x) f(x) f(a+b-x) dx}{\int_a^b w(x) dx}\right)^{\frac{1}{2}} \leq \sqrt{f(a) f(b)}.$$

**Proof.** We know that, see [32] or [35, p. 198], if  $g$  is log-convex, then

$$(3.3) \quad g\left(\frac{a+b}{2}\right) \leq \sqrt{g(x)g(a+b-x)} \leq \sqrt{g(a)g(b)}$$

for any  $x \in [a, b]$ .

For any  $p > 0$  the function  $f^{2p}$  is log-convex and by (3.3) we have

$$(3.4) \quad f^{2p}\left(\frac{a+b}{2}\right) \leq f^p(x) f^p(a+b-x) \leq f^p(a) f^p(b)$$

for any  $x \in [a, b]$ .

If we multiply (3.4) by  $w(x) \geq 0$  and integrate, then we get

$$\begin{aligned} f^{2p}\left(\frac{a+b}{2}\right) \int_a^b w(x) dx &\leq \int_a^b w(x) f^p(x) f^p(a+b-x) dx \\ &\leq f^p(a) f^p(b) \int_a^b w(x) dx \end{aligned}$$

namely

$$(3.5) \quad f^{2p}\left(\frac{a+b}{2}\right) \leq \frac{\int_a^b w(x) f^p(x) f^p(a+b-x) dx}{\int_a^b w(x) dx} \leq f^p(a) f^p(b).$$

Taking the power  $\frac{1}{2p}$  in (3.5) we get the desired result (3.1).  $\square$

We also have the inequality

$$(3.6) \quad f\left(\frac{a+b}{2}\right) \leq \frac{\int_a^b w(x) \sqrt{f(x)f(a+b-x)} dx}{\int_a^b w(x) dx} \leq \sqrt{f(a)f(b)},$$

that is a weighted version of (1.5).

If we take  $p = \frac{1}{4}$  in (3.1), then we get

$$(3.7) \quad f\left(\frac{a+b}{2}\right) \leq \left(\frac{\int_a^b w(x) \sqrt[4]{f(x)f(a+b-x)} dx}{\int_a^b w(x) dx}\right)^2 \leq \sqrt{f(a)f(b)}.$$

Using Jensen's inequality for the power  $p \geq 1$  ( $p \in (0, 1)$ ), namely

$$\left(\frac{\int_a^b w(x) g(x) dx}{\int_a^b w(x) dx}\right)^p \leq (\geq) \frac{\int_a^b w(x) g^p(x) dx}{\int_a^b w(x) dx},$$

we can start the following more precise result:

**Corollary 3.** Let  $f : [a, b] \rightarrow (0, \infty)$  be a log-convex function and  $w : [a, b] \rightarrow [0, \infty)$  be integrable and  $\int_a^b w(x) dx > 0$ .

If  $p \geq 1$ , then

$$(3.8) \quad f\left(\frac{a+b}{2}\right) \leq \left(\frac{\int_a^b w(x) f(x) f(a+b-x) dx}{\int_a^b w(x) dx}\right)^{\frac{1}{2}} \\ \leq \left(\frac{\int_a^b w(x) f^p(x) f^p(a+b-x) dx}{\int_a^b w(x) dx}\right)^{\frac{1}{2p}} \leq \sqrt{f(a) f(b)}.$$

If  $p \in (0, 1)$ , then

$$(3.9) \quad f\left(\frac{a+b}{2}\right) \leq \left(\frac{\int_a^b w(x) f^p(x) f^p(a+b-x) dx}{\int_a^b w(x) dx}\right)^{\frac{1}{2p}} \\ \leq \left(\frac{\int_a^b w(x) f(x) f(a+b-x) dx}{\int_a^b w(x) dx}\right)^{\frac{1}{2}} \leq \sqrt{f(a) f(b)}.$$

**Remark 2.** Let  $f : [a, b] \rightarrow (0, \infty)$  be a log-convex function. We observe that if we take in (3.1)  $w(x) = f^{-p}(a+b-x)$ ,  $p > 0$ , then we get

$$(3.10) \quad f\left(\frac{a+b}{2}\right) \leq \left(\frac{\int_a^b f^p(x) dx}{\int_a^b f^{-p}(x) dx}\right)^{\frac{1}{2p}} \leq \sqrt{f(a) f(b)}$$

for any  $p > 0$ .

In particular, we have the inequalities

$$(3.11) \quad f\left(\frac{a+b}{2}\right) \leq \left(\frac{\int_a^b f(x) dx}{\int_a^b \frac{1}{f(x)} dx}\right)^{\frac{1}{2}} \leq \sqrt{f(a) f(b)},$$

and

$$(3.12) \quad f\left(\frac{a+b}{2}\right) \leq \frac{\int_a^b \sqrt{f(x)} dx}{\int_a^b \frac{1}{\sqrt{f(x)}} dx} \leq \sqrt{f(a) f(b)}.$$

If we take in (3.10)  $f(x) = x^{-1}$ ,  $x \in [a, b] \subset (0, \infty)$ , which is log-convex, then we have

$$(3.13) \quad \left(\frac{a+b}{2}\right)^{-1} \leq \left(\frac{\int_a^b x^{-p} dx}{\int_a^b x^p dx}\right)^{\frac{1}{2p}} \leq \left(\sqrt{f(a)f(b)}\right)^{-1}$$

Observe that

$$\begin{aligned} \frac{\int_a^b x^{-p} dx}{\int_a^b x^p dx} &= \frac{\frac{1}{b-a} \int_a^b x^{-p} dx}{\frac{1}{b-a} \int_a^b x^p dx} \\ &= \frac{[L_{-p}(a,b)]^{-p}}{[L_p(a,b)]^p} \end{aligned}$$

and by (3.13) we have

$$\left(\frac{a+b}{2}\right)^{-1} \leq \left(\frac{1}{[L_p(a,b)][L_{-p}(a,b)]}\right)^{\frac{1}{2}} \leq \left(\sqrt{f(a)f(b)}\right)^{-1}$$

that is equivalent to

$$G(a,b) \leq \sqrt{[L_p(a,b)][L_{-p}(a,b)]} \leq A(a,b).$$

This can be also written as

$$(3.14) \quad G(a,b) \leq G(L_p(a,b), L_{-p}(a,b)) \leq A(a,b).$$

If we take in the first inequality (3.14)  $p = 1$ , then we get

$$(3.15) \quad G^2(a,b) \leq A(a,b)L(a,b).$$

We have the following weighted version of (1.4).

**Theorem 2.** Let  $f : [a, b] \rightarrow (0, \infty)$  be a log-convex function. If  $w : [a, b] \rightarrow [0, \infty)$  is integrable and  $\int_a^b w(x) dx > 0$ , then

$$(3.16) \quad \begin{aligned} &f\left(\frac{\int_a^b w(x) x dx}{\int_a^b w(x) dx}\right) \\ &\leq \exp\left(\frac{\int_a^b w(x) \ln f(x) dx}{\int_a^b w(x) dx}\right) \\ &\leq [f(b)]^{\frac{1}{b-a}\left(\frac{\int_a^b xw(x) dx}{\int_a^b w(x) dx} - a\right)} [f(a)]^{\frac{1}{b-a}\left(b - \frac{\int_a^b xw(x) dx}{\int_a^b w(x) dx}\right)} \\ &\leq \frac{1}{b-a} \left[ \left(\frac{\int_a^b xw(x) dx}{\int_a^b w(x) dx} - a\right) f(b) + \left(b - \frac{\int_a^b xw(x) dx}{\int_a^b w(x) dx}\right) f(a) \right]. \end{aligned}$$

**Proof.** Since  $\ln f$  is convex, then by Jensen's inequality we have

$$(3.17) \quad \ln f \left( \frac{\int_a^b w(x) x dx}{\int_a^b w(x) dx} \right) \leq \frac{\int_a^b w(x) \ln f(x) dx}{\int_a^b w(x) dx}.$$

Taking the exponential in (3.17) we get the first inequality in (3.16). Since  $\ln f$  is convex, then

$$(3.18) \quad \ln f(x) = \ln f \left( \frac{x-a}{b-a}b + \frac{b-x}{b-a}a \right) \leq \frac{x-a}{b-a} \ln f(b) + \frac{b-x}{b-a} \ln f(a)$$

for any  $x \in [a, b]$ .

By taking the weighted integral mean in (3.18) we get

$$(3.19) \quad \frac{\int_a^b w(x) \ln f(x) dx}{\int_a^b w(x) dx} \leq \frac{1}{b-a} \left[ \left( \frac{\int_a^b xw(x) dx}{\int_a^b w(x) dx} - a \right) \ln f(b) + \left( b - \frac{\int_a^b xw(x) dx}{\int_a^b w(x) dx} \right) \ln f(a) \right] \\ = \ln \left( [f(b)]^{\frac{1}{b-a} \left( \frac{\int_a^b xw(x) dx}{\int_a^b w(x) dx} - a \right)} [f(a)]^{\frac{1}{b-a} \left( b - \frac{\int_a^b xw(x) dx}{\int_a^b w(x) dx} \right)} \right).$$

By taking the exponential in (3.19), we get the second inequality in (3.16).

The last part of (3.16) follows by the weighted geometric mean-arithmetic mean inequality.  $\square$

**Remark 3.** If we take  $w(x) = 1$ ,  $x \in [a, b]$  in the first two inequalities (3.16), we recapture (1.4).

If we take  $w(x) = \frac{1}{x}$ ,  $x \in [a, b] \subset (0, \infty)$  in (3.16), then we get

$$(3.20) \quad f(L(a, b)) \leq \exp \left( \frac{1}{\ln b - \ln a} \int_a^b \frac{\ln f(x)}{x} dx \right) \\ \leq [f(b)]^{\frac{L(a,b)-a}{b-a}} [f(a)]^{\frac{b-L(a,b)}{b-a}} \\ \leq \frac{(L(a,b)-a)f(b) + (b-L(a,b))f(a)}{b-a},$$

where  $L(a, b)$  is the logarithmic mean, i.e.

$$L(a, b) = \begin{cases} \frac{b-a}{\ln b - \ln a} & \text{if } a \neq b \\ a & \text{if } a = b. \end{cases}$$

If we take  $w(x) = \frac{1}{x^2}$ ,  $x \in [a, b] \subset (0, \infty)$  in (3.16), then we get

$$\begin{aligned} (3.21) \quad f\left(\frac{G^2(a, b)}{L(a, b)}\right) &\leq \exp\left(\frac{ab}{b-a} \int_a^b \frac{\ln f(x)}{x^2} dx\right) \\ &\leq [f(b)]^{\frac{1}{b-a}\left(\frac{G^2(a, b)}{L(a, b)} - a\right)} [f(a)]^{\frac{1}{b-a}\left(b - \frac{G^2(a, b)}{L(a, b)}\right)} \\ &\leq \frac{1}{b-a} \left[ \left(\frac{G^2(a, b)}{L(a, b)} - a\right) f(b) + \left(b - \frac{G^2(a, b)}{L(a, b)}\right) f(a) \right]. \end{aligned}$$

We also have the alternative result:

**Theorem 3.** Let  $f : [a, b] \rightarrow (0, \infty)$  be a log-convex function. If  $w : [a, b] \rightarrow [0, \infty)$  is integrable and  $\int_a^b w(x) dx > 0$ , then

$$\begin{aligned} (3.22) \quad &f\left(\frac{\int_a^b w(x) x dx}{\int_a^b w(x) dx}\right) \\ &\leq \exp\left(\frac{\int_a^b w(x) \ln f(x) dx}{\int_a^b w(x) dx}\right) \\ &\leq \frac{\int_a^b w(x) f(x) dx}{\int_a^b w(x) dx} \leq \left(\frac{[f(a)]^b}{[f(b)]^a}\right)^{\frac{1}{b-a}} \frac{\int_a^b w(x) \left(\frac{f(b)}{f(a)}\right)^{\frac{x}{b-a}} dx}{\int_a^b w(x) dx} \\ &\leq \frac{1}{b-a} \left[ \left(\frac{\int_a^b x w(x) dx}{\int_a^b w(x) dx} - a\right) f(b) + \left(b - \frac{\int_a^b x w(x) dx}{\int_a^b w(x) dx}\right) f(a) \right]. \end{aligned}$$

**Proof.** Using Jensen's inequality for the exponential function we have

$$\begin{aligned} \exp\left(\frac{\int_a^b w(x) \ln f(x) dx}{\int_a^b w(x) dx}\right) &\leq \frac{\int_a^b w(x) \exp(\ln f(x)) dx}{\int_a^b w(x) dx} \\ &= \frac{\int_a^b w(x) f(x) dx}{\int_a^b w(x) dx} \end{aligned}$$

and the second inequality in (3.22) is proved.

From (3.18) and the arithmetic mean - geometric mean inequality we have



$$\begin{aligned}
 (3.23) \quad f(x) &\leq [f(b)]^{\frac{x-a}{b-a}} [f(a)]^{\frac{b-x}{b-a}} = \left( \frac{[f(a)]^b}{[f(b)]^a} \right)^{\frac{1}{b-a}} \left( \frac{f(b)}{f(a)} \right)^{\frac{x}{b-a}} \\
 &\leq \frac{x-a}{b-a} f(b) + \frac{b-x}{b-a} f(a)
 \end{aligned}$$

for any  $x \in [a, b]$ .

By taking the weighted integral mean in (3.23) we get

$$\begin{aligned}
 &\frac{\int_a^b w(x) f(x) dx}{\int_a^b w(x) dx} \\
 &\leq \left( \frac{[f(a)]^b}{[f(b)]^a} \right)^{\frac{1}{b-a}} \frac{\int_a^b w(x) \left( \frac{f(b)}{f(a)} \right)^{\frac{x}{b-a}} dx}{\int_a^b w(x) dx} \\
 &\leq \frac{1}{b-a} \left[ \left( \frac{\int_a^b x w(x) dx}{\int_a^b w(x) dx} - a \right) f(b) + \left( b - \frac{\int_a^b x w(x) dx}{\int_a^b w(x) dx} \right) f(a) \right]
 \end{aligned}$$

and the last part of (3.22) is proved.  $\square$

**Remark 4.** If we take  $w(x) = 1$ ,  $x \in [a, b]$  in (3.22), then we have

$$\begin{aligned}
 (3.24) \quad f\left(\frac{a+b}{2}\right) &\leq \exp\left(\frac{1}{b-a} \int_a^b \ln f(x) dx\right) \\
 &\leq \frac{1}{b-a} \int_a^b f(x) dx \leq L(f(a), f(b)) \leq \frac{f(a)+f(b)}{2}.
 \end{aligned}$$

If we take  $w(x) = \frac{1}{x}$ ,  $x \in [a, b] \subset (0, \infty)$  in (3.22), then we get

$$\begin{aligned}
 (3.25) \quad f(L(a, b)) &\leq \exp\left(\frac{1}{\ln b - \ln a} \int_a^b \frac{\ln f(x)}{x} dx\right) \\
 &\leq \frac{1}{\ln b - \ln a} \int_a^b \frac{f(x)}{x} dx \\
 &\leq \left( \frac{[f(a)]^b}{[f(b)]^a} \right)^{\frac{1}{b-a}} \frac{\int_a^b \frac{1}{x} \left( \frac{f(b)}{f(a)} \right)^{\frac{x}{b-a}} dx}{\ln b - \ln a} \\
 &\leq \frac{(L(a, b) - a) f(b) + (b - L(a, b)) f(a)}{b - a}.
 \end{aligned}$$

#### 4. Inequalities for Symmetric Weights

We say that the weight  $w : [a, b] \rightarrow [0, \infty)$  is *symmetric* on  $[a, b]$  if

$$w(a + b - x) = w(x) \text{ for all } x \in [a, b].$$

It is well known that if  $f : [a, b] \rightarrow \mathbf{R}$  is convex and  $w : [a, b] \rightarrow [0, \infty)$  is integrable and symmetric on  $[a, b]$ , then the *Fejér inequality* holds

$$(4.1) \quad f\left(\frac{a+b}{2}\right) \leq \frac{\int_a^b w(x) f(x) dx}{\int_a^b w(x) dx} \leq \frac{f(a) + f(b)}{2}.$$

If  $f : [a, b] \rightarrow (0, \infty)$  is a log-convex function on  $[a, b]$  and  $w : [a, b] \rightarrow [0, \infty)$  is integrable and symmetric on  $[a, b]$ , then by (4.1) we have

$$\ln f\left(\frac{a+b}{2}\right) \leq \frac{\int_a^b w(x) \ln f(x) dx}{\int_a^b w(x) dx} \leq \frac{\ln f(a) + \ln f(b)}{2},$$

which is equivalent to

$$(4.2) \quad f\left(\frac{a+b}{2}\right) \leq \exp\left(\frac{\int_a^b w(x) \ln f(x) dx}{\int_a^b w(x) dx}\right) \leq \sqrt{f(a) f(b)}.$$

**Theorem 1.** Let  $f : [a, b] \rightarrow (0, \infty)$  be a log-convex function on  $[a, b]$  and  $w : [a, b] \rightarrow [0, \infty)$  be integrable and symmetric on  $[a, b]$ . Then

$$(4.3) \quad f\left(\frac{a+b}{2}\right) \leq \exp\left(\frac{\int_a^b w(x) \ln f(x) dx}{\int_a^b w(x) dx}\right) \\ \leq \frac{\int_a^b w(x) \sqrt{f(x)f(a+b-x)} dx}{\int_a^b w(x) dx} \leq \frac{\int_a^b w(x) f(x) dx}{\int_a^b w(x) dx}.$$

**Proof.** By Jensen’s integral inequality for the exponential we have

$$(4.4) \quad \exp\left(\frac{\int_a^b w(x) \ln \sqrt{f(x) f(a+b-x)} dx}{\int_a^b w(x) dx}\right) \\ \leq \frac{\int_a^b w(x) \exp(\ln \sqrt{f(x) f(a+b-x)}) dx}{\int_a^b w(x) dx} \\ = \frac{\int_a^b w(x) \sqrt{f(x) f(a+b-x)} dx}{\int_a^b w(x) dx}.$$

Observe, by the symmetry of  $w$ , that

$$\begin{aligned}
& \int_a^b w(x) \ln \sqrt{f(x) f(a+b-x)} dx \\
&= \frac{1}{2} \left[ \int_a^b w(x) \ln f(x) dx + \int_a^b w(x) \ln f(a+b-x) dx \right] \\
&= \frac{1}{2} \left[ \int_a^b w(x) \ln f(x) dx + \int_a^b w(a+b-x) \ln f(a+b-x) dx \right] \\
&= \int_a^b w(x) \ln f(x) dx
\end{aligned}$$

since, obviously

$$\int_a^b w(a+b-x) \ln f(a+b-x) dx = \int_a^b w(x) \ln f(x) dx.$$

By (4.4) we then get the second inequality in (4.3).

By Cauchy-Buniakovski-Schwarz integral inequality we also have

$$\begin{aligned}
& \int_a^b w(x) \sqrt{f(x) f(a+b-x)} dx \\
&\leq \sqrt{\int_a^b w(x) f(x) dx} \sqrt{\int_a^b w(x) f(a+b-x) dx} \\
&= \sqrt{\int_a^b w(x) f(x) dx} \sqrt{\int_a^b w(x) f(x) dx} \\
&= \int_a^b w(x) f(x) dx,
\end{aligned}$$

which proves the third inequality in (4.3).  $\square$

The above inequality (4.2) may be generalized as follows by replacing  $f$  with  $f^{2p}$  for  $p > 0$ .

**Corollary 4.** *Let  $f : [a, b] \rightarrow (0, \infty)$  be a log-convex function on  $[a, b]$  and  $w : [a, b] \rightarrow [0, \infty)$  be integrable and symmetric on  $[a, b]$ . Then for any  $p > 0$  we have*

$$(4.5) \quad f\left(\frac{a+b}{2}\right) \leq \exp\left(\frac{\int_a^b w(x) \ln f(x) dx}{\int_a^b w(x) dx}\right)$$

$$\begin{aligned} &\leq \left( \frac{\int_a^b w(x) f^p(x) f^{p(a+b-x)} dx}{\int_a^b w(x) dx} \right)^{\frac{1}{2p}} \\ &\leq \left( \frac{\int_a^b w(x) f^{2p}(x) dx}{\int_a^b w(x) dx} \right)^{\frac{1}{2p}}. \end{aligned}$$

**Remark 5.** We observe that for  $p \geq 1$  we have

$$\begin{aligned} (4.6) \quad f\left(\frac{a+b}{2}\right) &\leq \exp\left(\frac{\int_a^b w(x) \ln f(x) dx}{\int_a^b w(x) dx}\right) \\ &\leq \left(\frac{\int_a^b w(x) f(x) f(a+b-x) dx}{\int_a^b w(x) dx}\right)^{\frac{1}{2}} \\ &\leq \left(\frac{\int_a^b w(x) f^p(x) f^{p(a+b-x)} dx}{\int_a^b w(x) dx}\right)^{\frac{1}{2p}} \\ &\leq \left(\frac{\int_a^b w(x) f^{2p}(x) dx}{\int_a^b w(x) dx}\right)^{\frac{1}{2p}} \end{aligned}$$

and for  $p \in (0, 1)$

$$\begin{aligned} (4.7) \quad f\left(\frac{a+b}{2}\right) &\leq \exp\left(\frac{\int_a^b w(x) \ln f(x) dx}{\int_a^b w(x) dx}\right) \\ &\leq \left(\frac{\int_a^b w(x) f^p(x) f^{p(a+b-x)} dx}{\int_a^b w(x) dx}\right)^{\frac{1}{2p}} \\ &\leq \left(\frac{\int_a^b w(x) f(x) f(a+b-x) dx}{\int_a^b w(x) dx}\right)^{\frac{1}{2}} \\ &\leq \left(\frac{\int_a^b w(x) f^2(x) dx}{\int_a^b w(x) dx}\right)^{\frac{1}{2}}. \end{aligned}$$

Finally, we have:

**Theorem 2.** Let  $f : [a, b] \rightarrow (0, \infty)$  be a log-convex function on  $[a, b]$  and  $w : [a, b] \rightarrow [0, \infty)$  be integrable and symmetric on  $[a, b]$ . Then for any  $p > 0$  we have

$$(4.8) \quad f\left(\frac{a+b}{2}\right) \leq \left(\frac{\int_a^b f^p(x)w(x)dx}{\int_a^b \frac{w(x)dx}{f^p(x)}}\right)^{\frac{1}{2p}} \leq \sqrt{f(a)f(b)}.$$

**Proof.** From (3.4) we have

$$(4.9) \quad f^{2p}\left(\frac{a+b}{2}\right) \frac{1}{f^p(a+b-x)} \leq f^p(x) \leq f^p(a)f^p(b) \frac{1}{f^p(a+b-x)}$$

for any  $x \in [a, b]$ .

If we multiply by  $w(x) \geq 0$  and integrate on  $[a, b]$ , then we get

$$\begin{aligned} & f^{2p}\left(\frac{a+b}{2}\right) \int_a^b \frac{w(x)dx}{f^p(a+b-x)} \\ & \leq \int_a^b f^p(x)w(x)dx \\ & \leq f^p(a)f^p(b) \int_a^b \frac{w(x)dx}{f^p(a+b-x)}. \end{aligned}$$

Since, by symmetry of  $w$  we have

$$\int_a^b \frac{w(x)dx}{f^p(a+b-x)} = \int_a^b \frac{w(a+b-x)dx}{f^p(a+b-x)} = \int_a^b \frac{w(x)dx}{f^p(x)},$$

which implies that

$$\begin{aligned} & f^{2p}\left(\frac{a+b}{2}\right) \int_a^b \frac{w(x)dx}{f^p(x)} \\ & \leq \int_a^b f^p(x)w(x)dx \\ & \leq f^p(a)f^p(b) \int_a^b \frac{w(x)dx}{f^p(x)}. \end{aligned}$$

and the inequality (4.8) is proved.  $\square$

**Remark 6.** If we write the inequality (4.8) for the log-convex function  $f : [a, b] \subset (0, \infty) \rightarrow (0, \infty)$ ,  $f(x) = \frac{1}{x}$ , then we have for  $p > 0$

$$\left(\frac{a+b}{2}\right)^{-1} \leq \left(\frac{\int_a^b x^{-p}w(x)dx}{\int_a^b x^p w(x)dx}\right)^{\frac{1}{2p}} \leq (\sqrt{ab})^{-1},$$

that is equivalent to

$$(4.10) \quad G(a, b) \leq \left( \frac{\int_a^b x^p w(x) dx}{\int_a^b x^{-p} w(x) dx} \right)^{\frac{1}{2p}} \leq A(a, b),$$

for any symmetric integrable weight  $w : [a, b] \rightarrow [0, \infty)$ .

## References

- [1] M. Alomari and M. Darus, The Hadamard's inequality for s-convex function. *Int. J. Math. Anal.* (Ruse) **2**, No. 13-16, pp. 639–646, (2008).
- [2] M. Alomari and M. Darus, Hadamard-type inequalities for s-convex functions. *Int. Math. Forum* **3**, No. 37-40, pp. 1965–1975, (2008).
- [3] G. A. Anastassiou, Univariate Ostrowski inequalities, revisited. *Monatsh. Math.*, **135**, No. 3, pp. 175–189, (2002).
- [4] G. D. Anderson, M. K. Vamanamurthy and M. Vuorinen, Generalized convexity and inequalities, *J. Math. Anal. Appl.* **335**, pp. 1294–1308, (2007).
- [5] N. S. Barnett, P. Cerone, S. S. Dragomir, M. R. Pinheiro, and A. Sofo, Ostrowski type inequalities for functions whose modulus of the derivatives are convex and applications. *Inequality Theory and Applications*, **Vol. 2** (Chinju/Masan, 2001), 19–32, Nova Sci. Publ., Hauppauge, NY, 2003. Preprint: *RGMA Res. Rep. Coll.* **5**, (2002), No. 2, Art. 1 [Online <http://rgmia.org/papers/v5n2/Paperwapp2q.pdf>].
- [6] E. F. Beckenbach, Convex functions, *Bull. Amer. Math. Soc.* **54**, pp. 439–460, (1948).
- [7] M. Bombardelli and S. Varošanec, Properties of h-convex functions related to the Hermite-Hadamard-Fejér inequalities. *Comput. Math. Appl.* **58**, No. 9, pp. 1869–1877, (2009).
- [8] W. W. Breckner, Stetigkeitsaussagen für eine Klasse verallgemeinerter konvexer Funktionen in topologischen linearen Räumen. (German) *Publ. Inst. Math. (Beograd) (N.S.)* **23 (37)**, pp. 13–20, (1978).

- [9] W. W. Breckner and G. Orbán, Continuity properties of rationally  $s$ -convex mappings with values in an ordered topological linear space. Universitatea "Babeş-Bolyai, Facultatea de Matematica, Cluj-Napoca, viii+92, (1978).
- [10] P. Cerone and S. S. Dragomir, Midpoint-type rules from an inequalities point of view, Ed. G. A. Anastassiou, *Handbook of Analytic-Computational Methods in Applied Mathematics*, CRC Press, New York., pp. 135-200, (2000).
- [11] P. Cerone and S. S. Dragomir, New bounds for the three-point rule involving the Riemann-Stieltjes integrals, in *Advances in Statistics Combinatorics and Related Areas*, C. Gulati, et al. (Eds.), World Science Publishing, pp. 53-62, (2002).
- [12] P. Cerone, S. S. Dragomir and J. Roumeliotis, Some Ostrowski type inequalities for  $n$ -time differentiable mappings and applications, *Demonstratio Mathematica*, **32** (2), pp. 697—712, (1999).
- [13] G. Cristescu, Hadamard type inequalities for convolution of  $h$ -convex functions. *Ann. Tiberiu Popoviciu Semin. Funct. Equ. Approx. Convexity* **8**, pp. 3–11, (2010).
- [14] S. S. Dragomir, Some remarks on Hadamard's inequalities for convex functions, *Extracta Math.*, **9** (2), pp. 88-94, (1994).
- [15] S. S. Dragomir, Refinements of the Hermite-Hadamard integral inequality for log-convex functions, *Austral. Math. Soc. Gaz.* **28**, No. 3, pp. 129–134, (2001).
- [16] S. S. Dragomir, Ostrowski's inequality for monotonous mappings and applications, *J. KSIAM*, **3** (1), pp. 127-135, (1999).
- [17] S. S. Dragomir, The Ostrowski's integral inequality for Lipschitzian mappings and applications, *Comp. Math. Appl.*, **38**, pp. 33-37, (1999).
- [18] S. S. Dragomir, On the Ostrowski's inequality for Riemann-Stieltjes integral, *Korean J. Appl. Math.*, **7**, pp. 477-485, (2000).
- [19] S. S. Dragomir, On the Ostrowski's inequality for mappings of bounded variation and applications, *Math. Ineq. & Appl.*, **4** (1), pp. 33-40, (2001).

- [20] S. S. Dragomir, On the Ostrowski inequality for Riemann-Stieltjes integral  $\int_a^b f(t) du(t)$  where  $f$  is of Hölder type and  $u$  is of bounded variation and applications, *J. KSIAM*, **5** (1), pp. 35-45, (2001).
- [21] S. S. Dragomir, Ostrowski type inequalities for isotonic linear functionals, *J. Inequal. Pure & Appl. Math.*, **3**(5), Art. 68, (2002).
- [22] S. S. Dragomir, An inequality improving the first Hermite-Hadamard inequality for convex functions defined on linear spaces and applications for semi-inner products. *J. Inequal. Pure Appl. Math.* **3**, No. 2, Article 31, 8, (2002).
- [23] S. S. Dragomir, An inequality improving the first Hermite-Hadamard inequality for convex functions defined on linear spaces and applications for semi-inner products, *J. Inequal. Pure and Appl. Math.*, **3** (2), Art. 31, (2002).
- [24] S. S. Dragomir, An inequality improving the second Hermite-Hadamard inequality for convex functions defined on linear spaces and applications for semi-inner products, *J. Inequal. Pure and Appl. Math.*, **3** (3), Art. 35, (2002).
- [25] S. S. Dragomir, An Ostrowski like inequality for convex functions and applications, *Revista Math. Complutense*, **16** (2), pp. 373-382, (2003).
- [26] S. S. Dragomir, *Operator Inequalities of Ostrowski and Trapezoidal Type*. Springer Briefs in Mathematics. Springer, New York, (2012). x+112 pp. ISBN: 978-1-4614-1778-1
- [27] S. S. Dragomir, Some new inequalities of Hermite-Hadamard type for GA-convex functions, Preprint *RGMA Res. Rep. Coll.* 18 (2015), Art 33. [<http://rgmia.org/papers/v18/v18a33.pdf>].
- [28] S. S. Dragomir, New inequalities of Hermite-Hadamard type for log-convex functions, Preprint *RGMA Res. Rep. Coll.* 18, (2015), Art 42. [<http://rgmia.org/papers/v18/v18a42.pdf>].
- [29] S. S. Dragomir, P. Cerone, J. Roumeliotis and S. Wang, A weighted version of Ostrowski inequality for mappings of Hölder type and applications in numerical analysis, *Bull. Math. Soc. Sci. Math. Romania*, **42(90)** (4), pp. 301-314, (1999).



- [30] S. S. Dragomir and S. Fitzpatrick, The Hadamard inequalities for  $s$ -convex functions in the second sense. *Demonstratio Math.* **32**, No. 4, pp. 687–696, (1999).
- [31] S. S. Dragomir and S. Fitzpatrick, The Jensen inequality for  $s$ -Breckner convex functions in linear spaces. *Demonstratio Math.* **33**, No. 1, pp. 43–49, (2000).
- [32] S. S. Dragomir and B. Mond, Integral inequalities of Hadamard's type for log-convex functions, *Demonstratio Math.*, **31** (2), pp. 354–364, (1998).
- [33] S. S. Dragomir and C. E. M. Pearce, On Jensen's inequality for a class of functions of Godunova and Levin. *Period. Math. Hungar.*, **33**, No. 2, pp. 93–100, (1996).
- [34] S. S. Dragomir and C. E. M. Pearce, Quasi-convex functions and Hadamard's inequality, *Bull. Austral. Math. Soc.* **57**, pp. 377–385, (1998).
- [35] S. S. Dragomir and C. E. M. Pearce, *Selected Topics on Hermite-Hadamard Inequalities and Applications*, RGMIA Monographs, Victoria University, (2000).
- [36] S. S. Dragomir, J. Pečarić and L. Persson, Some inequalities of Hadamard type. *Soochow J. Math.* **21**, No. 3, pp. 335–341, (1995).
- [37] S. S. Dragomir and Th. M. Rassias (Eds), *Ostrowski Type Inequalities and Applications in Numerical Integration*, Kluwer Academic Publisher, (2002).
- [38] S. S. Dragomir and S. Wang, A new inequality of Ostrowski's type in  $L_1$ -norm and applications to some special means and to some numerical quadrature rules, *Tamkang J. of Math.*, **28**, pp. 239–244, (1997).
- [39] S. S. Dragomir and S. Wang, Applications of Ostrowski's inequality to the estimation of error bounds for some special means and some numerical quadrature rules, *Appl. Math. Lett.*, **11**, pp. 105–109, (1998).
- [40] S. S. Dragomir and S. Wang, A new inequality of Ostrowski's type in  $L_p$ -norm and applications to some special means and to some numerical quadrature rules, *Indian J. of Math.*, **40** (3), pp. 245–304, (1998).

- [41] A. El Farissi, Simple proof and refinement of Hermite-Hadamard inequality, *J. Math. Ineq.* **4**, No. 3, pp. 365–369, (2010).
- [42] L. Fejér, Über die Fourierreihen, II, *Math. Naturwiss. Anz Ungar. Akad. Wiss.* **24**, pp. 369-390, (1906). (In Hungarian).
- [43] P. M. Gill, C. E. M. Pearce and J. Pečarić, Hadamard's inequality for r-convex functions, *Journal of Mathematical Analysis and Applications.* **215**, No. 2, pp. 461-470, (1997).
- [44] E. K. Godunova and V. I. Levin, Inequalities for functions of a broad class that contains convex, monotone and some other forms of functions. (Russian) *Numerical mathematics and mathematical physics* (Russian), 138–142, 166, Moskov. Gos. Ped. Inst., Moscow, (1985).
- [45] H. Hudzik and L. Maligranda, Some remarks on s-convex functions. *Aequationes Math.* **48**, No. 1, pp. 100–111, (1994).
- [46] E. Kikianty and S. S. Dragomir, Hermite-Hadamard's inequality and the p-HH-norm on the Cartesian product of two copies of a normed space, *Math. Inequal. Appl. Mathematical Inequalities Applications*, Volume 13, Number 1, pp. 1-32, (2010).
- [47] U. S. Kirmaci, M. Klaričić Bakula, M. E. Özdemir and J. Pečarić, Hadamard-type inequalities for s-convex functions. *Appl. Math. Comput.* **193**, No. 1, pp. 26–35, (2007).
- [48] M. A. Latif, On some inequalities for h-convex functions. *Int. J. Math. Anal.* (Ruse) **4**, No. 29-32, pp. 1473–1482, (2010).
- [49] D. S. Mitrinović and I. B. Lacković, Hermite and convexity, *Aequationes Math.* **28**, pp. 229–232, (1985).
- [50] D. S. Mitrinović and J. E. Pečarić, Note on a class of functions of Godunova and Levin. *C. R. Math. Rep. Acad. Sci. Canada* **12**, No. 1, pp. 33–36, (1990).
- [51] M. A. Noor, K. I. Noor and M. U. Awan, Some inequalities for geometrically-arithmetically  $h$ -convex functions, *Creat. Math. Inform.* **23**, No. 1, 91-98, (2014).
- [52] C. E. M. Pearce and A. M. Rubinov, P-functions, quasi-convex functions, and Hadamard-type inequalities. *J. Math. Anal. Appl.* **240**, No. 1, pp. 92–104, (1999).

- [53] J. E. Pečarić and S. S. Dragomir, On an inequality of Godunova-Levin and some refinements of Jensen integral inequality. *Itinerant Seminar on Functional Equations, Approximation and Convexity* (Cluj-Napoca, 263–268, Preprint, 89-6, Univ. "Babeş-Bolyai, Cluj-Napoca, (1989).
- [54] J. Pečarić and S. S. Dragomir, A generalization of Hadamard's inequality for isotonic linear functionals, *Radovi Mat.* (Sarajevo) **7**, pp. 103–107, (1991).
- [55] M. Radulescu, S. Radulescu and P. Alexandrescu, On the Godunova-Levin-Schur class of functions. *Math. Inequal. Appl.* **12**, No. 4, pp. 853–862, (2009).
- [56] M. Z. Sarikaya, A. Sağlam, and H. Yildirim, On some Hadamard-type inequalities for h-convex functions. *J. Math. Inequal.* **2**, No. 3, pp. 335–341, (2008).
- [57] E. Set, M. E. Özdemir and M. Z. Sarikaya, New inequalities of Ostrowski's type for s-convex functions in the second sense with applications. *Facta Univ. Ser. Math. Inform.* **27**, No. 1, pp. 67–82, (2012).
- [58] M. Z. Sarikaya, E. Set and M. E. Özdemir, On some new inequalities of Hadamard type involving h-convex functions. *Acta Math. Univ. Comenian.* (N.S.) **79**, No. 2, pp. 265–272, (2010).
- [59] W. T. Sulaiman, Refinements to Hadamard's inequality for log-convex functions. *Applied Mathematics*, **2**, pp. 899-903, (2011).
- [60] M. Tunç, Ostrowski-type inequalities via h-convex functions with applications to special means. *J. Inequal. Appl.*, **326**, (2013).
- [61] S. Varošanec, On h-convexity. *J. Math. Anal. Appl.* **326**, No. 1, pp. 303–311, (2007).
- [62] X.-M. Zhang, Y.-M. Chu and X.-H. Zhang, The Hermite-Hadamard type inequality of GA-convex functions and its application, *Journal of Inequalities and Applications*, Volume, Article ID 507560, 11 pages, (2010).

**S. S. Dragomir**

Mathematics,  
College of Engineering & Science  
Victoria University,  
P. O. Box 14428  
Melbourne City, MC 8001,  
Australia  
e-mail : sever.dragomir@vu.edu.au