



VICTORIA UNIVERSITY
MELBOURNE AUSTRALIA

A new integral version of generalized Ostrowski-Grüss type inequality with applications

This is the Published version of the following publication

Dragomir, Sever S, Khan, Asif R, Khan, Maria, Mehmood, Faraz and Shaikh, Muhammad Awais (2022) A new integral version of generalized Ostrowski-Grüss type inequality with applications. *Journal of King Saud University - Science*, 34 (5). ISSN 1018-3647

The publisher's official version can be found at
<https://www.sciencedirect.com/science/article/pii/S1018364722002385>
Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/45660/>



Contents lists available at ScienceDirect

Journal of King Saud University – Science

journal homepage: www.sciencedirect.com

Original article

A new integral version of generalized Ostrowski-Grüss type inequality with applications

Sever S. Dragomir^a, Asif R. Khan^b, Maria Khan^{b,c}, Faraz Mehmood^{c,*}, Muhammad Awais Shaikh^d^a Department of Mathematics, College of Engineering & Science, Victoria University, PO Box 14428, Melbourne City, MC 8001, Australia^b Department of Mathematics, University of Karachi, University Road, Karachi 75270, Pakistan^c Department of Mathematics, Dawood University of Engineering and Technology, New M. A Jinnah Road, Karachi 74800, Pakistan^d Department of Mathematics, Nabi Bagh Z. M. Govt. Science College, Saddar, Karachi 74400, Pakistan

ARTICLE INFO

Article history:

Received 25 July 2021

Revised 10 April 2022

Accepted 22 April 2022

Available online 28 April 2022

2010 Mathematics Subject Classifications:

26D15

26D20

26D99

Keywords:

Ostrowski-Grüss inequality

Čebyšev functional

Korkine's identity

Cauchy-Schwartz inequality

numerical integration

Special means

ABSTRACT

Our aim is to improve and further generalize the result of integral Ostrowski-Grüss type inequalities involving differentiable functions and then apply these obtained inequalities to probability theory, special means and numerical integration.

© 2022 Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

In Ostrowski (1938), Ostrowski presented an inequality which is now known as “Ostrowski's inequality” stated below:

$$\left| \zeta(z) - \frac{1}{n-m} \int_m^n \zeta(\tau) d\tau \right| \leq \left[\frac{1}{4} + \frac{(z - \frac{m+n}{2})^2}{(n-m)^2} \right] (n-m)M, \quad z \in [m, n] \quad (1.1)$$

* Corresponding author.

E-mail addresses: sever.dragomir@vu.edu.au (S.S. Dragomir), asifrk@uok.edu.pk (A.R. Khan), maria.khan@duet.edu.pk (M. Khan), faraz.mehmood@duet.edu.pk (F. Mehmood).

Peer review under responsibility of King Saud University.



where $\zeta : [m, n] \rightarrow \mathbb{R}$ is a differentiable function such that $|\zeta'(z)| \leq M$, for every $z \in [m, n]$.

In present era, a large number of papers has been written about generalizations of Ostrowski's inequality see for example (Anastassiou, 1997; Cheng, 2001; Dragomir and Wang, 1997; Irshad and Khan, 2017; Liu, 2008; Matić et al., 2000; Milovanovic and Pecaric, 1976; Shaikh et al., 2021; Zafar and Mir, 2010). Ostrowski's inequality has proven to be an important tool for improvement of various branches of mathematical sciences. Very well said (Zafar, 2010) “Inequalities involving integrals that create bounds in the physical quantities are of great significance in the sense that these kinds of inequalities are not only used in approximation theory, operator theory, nonlinear analysis, numerical integration, stochastic analysis, information theory, statistics and probability theory but we may also see their uses in the various fields of biological sciences, engineering and physics”.

In the history, an important inequality that “estimate for the difference between the product of the integral of two functionals and the integral of their product” is known as “Grüss inequality”.

<https://doi.org/10.1016/j.jksus.2022.102057>

1018-3647/© 2022 Published by Elsevier B.V. on behalf of King Saud University.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

This celebrated integral inequality was proved by Grüss (1935) in 1935, is stated below (see also Mitrinović et al. (1994) [p. 296]),

$$\left| \frac{1}{n-m} \int_m^n \zeta(z)\eta(z) dz - \left(\frac{1}{n-m} \int_m^n \zeta(z) dz \right) \left(\frac{1}{n-m} \int_m^n \eta(z) dz \right) \right| \leq \frac{1}{4}(M_1 - m_1)(N_1 - n_1) \tag{1.2}$$

provided that ζ and η are integrable functions on $[m, n]$ such that $m_1 \leq \zeta(z) \leq M_1, \quad n_1 \leq \eta(z) \leq N_1,$

$\forall z \in [m, n]$, where m_1, M_1, n_1, N_1 are real constants.

By using Grüss inequality, Dragomir and Wang proved an inequality, in the year 1997, which we would refer as “Ostrowski-Grüss inequality” (Dragomir and Wang, 1997) which is stated as follows:

Proposition 1.1. Suppose $\zeta : I \rightarrow \mathbb{R}$ be a function differentiable in the interior I° of I , where $I \subseteq \mathbb{R}$, and let $m, n \in I^\circ$ and $n > m$. If $\gamma \leq \zeta'(z) \leq \Gamma, z \in [m, n]$ for real constants γ, Γ , then

$$\left| \zeta(z) - \frac{1}{n-m} \int_m^n \zeta(\tau) d\tau - \frac{\zeta(n) - \zeta(m)}{n-m} \left(z - \frac{m+n}{2} \right) \right| \leq \frac{1}{4}(n-m)(\Gamma - \gamma) \tag{1.3}$$

holds, $\forall z \in [m, n]$.

Above inequality gives a relationship between Ostrowski inequality (1.1) and Grüss inequality (1.2).

If ζ and g belong to $L_2[m, n]$, then the Čebyšev functional $T(\zeta, \eta)$ is defined as

$$T(\zeta, \eta) = \frac{1}{n-m} \int_m^n \zeta(z)\eta(z) dz - \left(\frac{1}{n-m} \int_m^n \zeta(z) dz \right) \left(\frac{1}{n-m} \int_m^n \eta(z) dz \right).$$

From Matić et al. (2000) pre-Grüss inequality is given below.

Proposition 1.2. Let $\zeta, \eta : [m, n] \rightarrow \mathbb{R}$ be integrable such that $\zeta\eta \in L(m, n)$. If

$$\gamma \leq \eta(z) \leq \Gamma \quad \text{for } z \in [m, n],$$

then

$$|T(\zeta, \eta)| \leq \frac{1}{2}(\Gamma - \gamma)\sqrt{T(\zeta, \zeta)}.$$

In the article (Matić et al., 2000) of year 2000, Matić, Pecarić and Ujević improved inequality (1.1), by using pre-Grüss inequality, which is as follows:

Proposition 1.3. Suppose $\zeta : I \rightarrow \mathbb{R}$ be a function differentiable in the interior I° of I , where $I \subseteq \mathbb{R}$, and let $m, n \in I^\circ$ and $n > m$. If $\gamma \leq \zeta'(z) \leq \Gamma, z \in [m, n]$ for real constants γ, Γ , then

$$\left| \zeta(z) - \frac{1}{n-m} \int_m^n \zeta(\tau) d\tau - \frac{\zeta(n) - \zeta(m)}{n-m} \left(z - \frac{m+n}{2} \right) \right| \leq \frac{1}{4\sqrt{3}}(n-m)(\Gamma - \gamma)$$

holds, $\forall z \in [m, n]$.

In the article (Barnett et al., 2000), by using Čebyšev functional, improved the Matić-Pecarić-Ujević result (1.3) in terms of “Euclidean norm” as under:

Proposition 1.4. Let function $\zeta : [m, n] \rightarrow \mathbb{R}$ be an absolutely continuous and derivative $\zeta' \in L_2[m, n]$. If $\gamma \leq \zeta'(\tau) \leq \Gamma$ almost everywhere for $\tau \in [m, n]$, then $\forall z \in [m, n]$

$$\begin{aligned} & \left| \zeta(z) - \frac{1}{n-m} \int_m^n \zeta(\tau) d\tau - \frac{\zeta(n) - \zeta(m)}{n-m} \left(z - \frac{m+n}{2} \right) \right| \\ & \leq \frac{n-m}{2\sqrt{3}} \left[\frac{1}{n-m} \|\zeta'\|_2^2 - \left(\frac{\zeta(n) - \zeta(m)}{n-m} \right)^2 \right]^{\frac{1}{2}} \\ & \leq \frac{1}{4\sqrt{3}}(n-m)(\Gamma - \gamma) \end{aligned} \tag{1.4}$$

holds.

This article is divided into six sections: the 1st section totally based on introduction and preliminaries. In the 2nd section, we would give our main result about generalization of integral Ostrowski-Grüss type inequalities and would discuss its different special cases. In the 3rd, 4th and 5th sections, using the obtained result we would give some applications to probability theory, special means and numerical integration respectively and the 6th concludes the article.

2. New generalization of integral Ostrowski-Grüss type inequality

Our main theorem of this section is given in the following:

Theorem 2.1. Let $\zeta : [m, n] \rightarrow \mathbb{R}$ be a differentiable function whose 1st derivative belongs to $L_2(m, n)$. If $\gamma \leq \zeta'(\tau) \leq \Gamma$ almost everywhere for $\tau \in [m, n]$, then $\forall z \in [m + \lambda \frac{n-m}{2}, \frac{m+n}{2}]$ and $\lambda \in [0, 1]$

$$\begin{aligned} & \left| (1-\lambda) \frac{\zeta(z) + \zeta(m+n-z)}{2} + \lambda \frac{\zeta(m) + \zeta(n)}{2} - \frac{1}{n-m} \int_m^n \zeta(\tau) d\tau \right| \\ & \leq \left[\frac{(n-m)^2}{12} (3\lambda^2 - 3\lambda + 1) + \left(z - \frac{m+n}{2} \right)^2 (1-\lambda) \right. \\ & \quad \left. + \frac{(n-m)(1-\lambda)^2}{2} \left(z - \frac{m+n}{2} \right) \right]^{\frac{1}{2}} \left[\frac{1}{n-m} \|\zeta'\|_2^2 - \left(\frac{\zeta(n) - \zeta(m)}{n-m} \right)^2 \right]^{\frac{1}{2}} \\ & \leq \frac{1}{2}(\Gamma - \gamma) \left[\frac{(n-m)^2}{12} (3\lambda^2 - 3\lambda + 1) + \left(z - \frac{m+n}{2} \right)^2 (1-\lambda) \right. \\ & \quad \left. + \frac{(n-m)(1-\lambda)^2}{2} \left(z - \frac{m+n}{2} \right) \right]^{\frac{1}{2}} \end{aligned} \tag{2.1}$$

holds.

Proof. We begin the proof of this theorem by defining the piecewise continuous function $K : [m, n]^2 \rightarrow \mathbb{R}$ for $\lambda \in [0, 1]$ as:

$$K(z, \tau; \lambda) = \begin{cases} \tau - m - \lambda \frac{(n-m)}{2}, & \text{if } \tau \in [m, z], \\ \tau - \frac{m+n}{2}, & \text{if } \tau \in (z, m+n-z), \\ \tau - n + \lambda \frac{(n-m)}{2}, & \text{if } \tau \in (m+n-z, n), \end{cases}$$

by Korkine's identity

$$T(\zeta, g) := \frac{1}{2(n-m)^2} \int_m^n \int_m^n (\zeta(\tau) - \zeta(s))(g(\tau) - g(s)) d\tau ds, \tag{2.2}$$

we obtain

$$\begin{aligned} & \frac{1}{n-m} \int_m^n K(z, \tau; \lambda) \zeta'(\tau) d\tau - \frac{1}{n-m} \int_m^n K(z, \tau; \lambda) dt \int_m^n \zeta'(\tau) d\tau \\ & = \frac{1}{2(n-m)^2} \int_m^n \int_m^n (K(z, \tau; \lambda) - K(z, s; \lambda)) (\zeta'(\tau) - \zeta'(s)) d\tau ds. \end{aligned} \tag{2.3}$$

Since

$$\begin{aligned} & \frac{1}{n-m} \int_m^n K(z, \tau; \lambda) \zeta'(\tau) d\tau = (1-\lambda) \frac{\zeta(z) + \zeta(m+n-z)}{2} + \lambda \frac{\zeta(m) + \zeta(n)}{2} \\ & - \frac{1}{n-m} \int_m^n \zeta(\tau) d\tau, \\ & \int_m^n K(z, \tau; \lambda) d\tau = 0, \end{aligned}$$

then by (2.3) we get the following identity

$$\begin{aligned} & (1-\lambda) \frac{\zeta(z) + \zeta(m+n-z)}{2} + \lambda \frac{\zeta(m) + \zeta(n)}{2} - \frac{1}{n-m} \int_m^n \zeta(\tau) d\tau \\ & = \frac{1}{2(n-m)^2} \int_m^n \int_m^n (K(z, \tau; \lambda) - K(z, s; \lambda)) (\zeta'(\tau) - \zeta'(s)) d\tau ds, \end{aligned} \tag{2.4}$$

$$\forall z \in [m + \lambda \frac{n-m}{2}, \frac{m+n}{2}] \text{ and } \lambda \in [0, 1].$$

By applying Cauchy-Schwartz inequality for double integrals, we can write

$$\begin{aligned} & \frac{1}{2(n-m)^2} \left| \int_m^n \int_m^n (K(z, \tau; \lambda) - K(z, s; \lambda)) (\zeta'(\tau) - \zeta'(s)) d\tau ds \right| \\ & \leq \left(\frac{1}{2(n-m)^2} \int_m^n \int_m^n (K(z, \tau; \lambda) - K(z, s; \lambda))^2 d\tau ds \right)^{\frac{1}{2}} \\ & \times \left(\frac{1}{2(n-m)^2} \int_m^n \int_m^n (\zeta'(\tau) - \zeta'(s))^2 d\tau ds \right)^{\frac{1}{2}}. \end{aligned} \tag{2.5}$$

However

$$\begin{aligned} & \frac{1}{2(n-m)^2} \int_m^n \int_m^n (K(z, \tau; \lambda) - K(z, s; \lambda))^2 d\tau ds \\ & = \frac{1}{(n-m)} \int_m^n K^2(z, \tau; \lambda) d\tau - \left(\frac{1}{n-m} \int_m^n K(z, \tau; \lambda) d\tau \right)^2 \\ & = \frac{1}{(n-m)} \left[\frac{2}{3} \left((z-m-\lambda \frac{n-m}{2})^3 - (z-\frac{m+n}{2})^3 \right) + \frac{\lambda^3 (n-m)^3}{12} \right]. \end{aligned} \tag{2.6}$$

Consider above terms in the following and simplifying:

$$\begin{aligned} & (z-m-\lambda \frac{n-m}{2})^3 - (z-\frac{m+n}{2})^3 \\ & = \frac{(n-m)^3}{8} (1-\lambda)^3 + \frac{3}{2} (z-\frac{m+n}{2})^2 (n-m)(1-\lambda) \\ & + \frac{3}{4} (n-m)^2 (1-\lambda)^2 (z-\frac{m+n}{2}), \end{aligned} \tag{2.7}$$

and

$$\frac{1}{2(n-m)^2} \int_m^n \int_m^n (\zeta'(\tau) - \zeta'(s))^2 d\tau ds = \frac{1}{(n-m)} \|\zeta'\|_2^2 - \left(\frac{\zeta(n) - \zeta(m)}{n-m} \right)^2. \tag{2.8}$$

Using (2.4), (2.6), (2.7) and (2.8), we get the 1st inequality of (2.1). Since $\gamma \leq \zeta'(\tau) \leq \Gamma$ almost everywhere for $\tau \in [m, n]$, by applying Grüss inequality (1.2) we get

$$0 \leq \frac{1}{n-m} \int_m^n (\zeta'(\tau))^2 d\tau - \left(\frac{1}{n-m} \int_m^n \zeta'(\tau) d\tau \right)^2 \leq \frac{1}{4} (\Gamma - \gamma)^2, \tag{2.9}$$

which completes the proof of last inequality of (2.1). □

Following remark (Remark 1 of Barnett et al. (2000)) is also valid for our main result.

Remark 2.2. Since $L_\infty[m, n] \subset L_2[m, n]$ (and the inclusion is strict), then we remark that the inequality (2.1) can be applied also for the mappings ζ whose derivatives are unbounded on (m, n) , but $\zeta' \in L_2[m, n]$.

Remark 2.3. Since $3\lambda^2 - 3\lambda + 1 \leq 1, \forall \lambda \in [0, 1]$ and this is minimum when $\lambda = \frac{1}{2}$. Therefore, (2.1) captures various special cases of main result which is obtained by authors of article (Barnett et al., 2000) as can be seen in remark given below.

Remark 2.4. We can get different special cases of (2.1) by using several values of λ by fixing $z = \frac{m+n}{2}$. Under the assumptions of Theorem 2.1 following results (special cases) are valid:

Special Case I: For $\lambda = 1$ (2.1) gives trapezoid inequality

$$\begin{aligned} & \left| \frac{\zeta(m) + \zeta(n)}{2} - \frac{1}{n-m} \int_m^n \zeta(\tau) d\tau \right| \\ & \leq \frac{1}{2\sqrt{3}} \left[(n-m) \|\zeta'\|_2^2 - (\zeta(n) - \zeta(m))^2 \right]^{\frac{1}{2}} \\ & \leq \frac{1}{4\sqrt{3}} (\Gamma - \gamma)(n-m), \end{aligned}$$

which is Remark 3.2 (i) of Zafar (2010).

Special Case II: For $\lambda = 0$ (2.1) gives mid-point inequality

$$\begin{aligned} & \left| \zeta\left(\frac{m+n}{2}\right) - \frac{1}{n-m} \int_m^n \zeta(\tau) d\tau \right| \\ & \leq \frac{1}{2\sqrt{3}} \left[(n-m) \|\zeta'\|_2^2 - (\zeta(n) - \zeta(m))^2 \right]^{\frac{1}{2}} \\ & \leq \frac{1}{4\sqrt{3}} (\Gamma - \gamma)(n-m). \end{aligned}$$

which is Corollary 1 of Barnett et al. (2000) and Remark 3.2 (ii) of Zafar (2010).

Special Case III: For $\lambda = \frac{1}{2}$ (2.1) gives averaged mid-point and trapezoid inequality

$$\begin{aligned} & \left| \frac{\zeta(m) + 2\zeta\left(\frac{m+n}{2}\right) + \zeta(n)}{4} - \frac{1}{n-m} \int_m^n \zeta(\tau) d\tau \right| \\ & \leq \frac{1}{4\sqrt{3}} \left[(n-m) \|\zeta'\|_2^2 - (\zeta(n) - \zeta(m))^2 \right]^{\frac{1}{2}} \\ & \leq \frac{1}{8\sqrt{3}} (\Gamma - \gamma)(n-m). \end{aligned}$$

which is Remark 3.2 (iii) of Zafar (2010).

Special Case IV: For $\lambda = \frac{1}{3}$ (2.1) gives a variant of Simpson's inequality for differentiable function ζ

$$\begin{aligned} & \left| \frac{\zeta(m) + 4\zeta\left(\frac{m+n}{2}\right) + \zeta(n)}{6} - \frac{1}{n-m} \int_m^n \zeta(\tau) d\tau \right| \\ & \leq \frac{1}{6} \left[(n-m) \|\zeta'\|_2^2 - (\zeta(n) - \zeta(m))^2 \right]^{\frac{1}{2}} \\ & \leq \frac{1}{12} (\Gamma - \gamma)(n-m). \end{aligned}$$

which is Remark 3.2 (iv) of Zafar (2010).

3. Application to probability theory

Suppose random variable 'Z' be continuous with PDF $\zeta : [m, n] \rightarrow \mathbb{R}_+$ and CDF $\Phi : [m, n] \rightarrow [0, 1]$ is defined as

$$\Phi(z) = \int_m^z \zeta(\tau) d\tau, \quad z \in \left[m + \lambda \frac{n-m}{2}, \frac{m+n}{2} \right],$$

and

$$E(Z) = \int_m^n \tau \zeta(\tau) d\tau,$$

is expectation of random variable 'Z' on $[m, n]$. Then we have following result:

Theorem 3.1. Let the suppositions of Theorem 2.1 be valid and if PDF $\zeta \in L_2[m, n]$, then

$$\begin{aligned}
 & \left| (1-\lambda) \frac{\Phi(z) + \Phi(m+n-z)}{2} + \frac{\lambda}{2} - \frac{n-E(Z)}{n-m} \right| \\
 & \leq \frac{1}{n-m} \left[\frac{(n-m)^2}{12} (3\lambda^2 - 3\lambda + 1) + \left(z - \frac{m+n}{2} \right)^2 (1-\lambda) \right. \\
 & \quad \left. + \frac{(n-m)(1-\lambda)^2}{2} \left(z - \frac{m+n}{2} \right) \right]^{\frac{1}{2}} \left[(n-m) \|\Phi'\|_2^2 - 1 \right]^{\frac{1}{2}} \\
 & \leq \frac{(H-h)}{2} \left[\frac{(n-m)^2}{12} (3\lambda^2 - 3\lambda + 1) + \left(z - \frac{m+n}{2} \right)^2 (1-\lambda) \right. \\
 & \quad \left. + \frac{(n-m)(1-\lambda)^2}{2} \left(z - \frac{m+n}{2} \right) \right]^{\frac{1}{2}},
 \end{aligned} \tag{3.1}$$

where $h \leq \Phi'(\tau) \leq H, \forall \tau \in [m, n]$.

Proof. Put $\zeta = \Phi$ in (2.1) we obtain (3.2), by applying the identity

$$\int_m^n \Phi(\tau) d\tau = n - E(Z) \quad \text{where} \quad \Phi(m) = 0, \Phi(n) = 1.$$

□

Corollary 3.2. Under the assumptions as stated in Theorem 3.1, if we put $z = \frac{m+n}{2}$, then

$$\begin{aligned}
 & \left| (1-\lambda) \Phi\left(\frac{m+n}{2}\right) + \frac{\lambda}{2} - \frac{n-E(Z)}{n-m} \right| \\
 & \leq \frac{1}{2\sqrt{3}} (3\lambda^2 - 3\lambda + 1)^{\frac{1}{2}} \left[(n-m) \|\Phi'\|_2^2 - 1 \right]^{\frac{1}{2}} \\
 & \leq \frac{(n-m)}{4\sqrt{3}} (3\lambda^2 - 3\lambda + 1)^{\frac{1}{2}} (H-h)
 \end{aligned}$$

hold for $h \leq \Phi'(\tau) \leq H \forall \tau \in [m, n]$.

Remark 3.3. The Corollary 3.2 is in fact Corollary 3.1 of Zafar (2010).

4. Application to special means

Before we proceed further we need here some definitions of special means.

Special Means: These means can be found in Zafar (2010).

(a) Arithmetic Mean

$$A = \frac{m+n}{2}; \quad m, n \geq 0.$$

(b) Geometric Mean

$$G = G(m, n) = \sqrt{mn}; \quad m, n \geq 0.$$

(c) Harmonic Mean

$$H = H(m, n) = \frac{2}{\frac{1}{m} + \frac{1}{n}}; \quad m, n > 0.$$

(d) Logarithmic Mean

$$L = L(m, n) = \begin{cases} m, & \text{if } m = n \\ \frac{n-m}{\ln n - \ln m}, & \text{if } m \neq n; m, n > 0. \end{cases}$$

(e) Identric Mean

$$I = I(m, n) = \begin{cases} m, & \text{if } m = n \\ \ln \left(\frac{\left(\frac{m}{n}\right)^{\frac{1}{n-m}}}{e} \right), & \text{if } m \neq n; m, n > 0. \end{cases}$$

(f) p-Logarithmic Mean

$$L_p = L_p(m, n) = \begin{cases} m, & \text{if } m = n \\ \left(\frac{p^{p+1} - m^{p+1}}{(p+1)(n-m)} \right)^{\frac{1}{p}}, & \text{if } m \neq n, \end{cases}$$

where $p \in \mathbb{R} \setminus \{-1, 0\}, m, n > 0$. It is known that “ L_p is monotonically increasing over $p \in \mathbb{R}$ ”, “ $L_0 = I$ ” and “ $L_{-1} = L$ ”.

Example 4.1. Consider $\zeta(z) = z^p, p \in \mathbb{R} \setminus \{-1, 0\}$, then for $n > m$

$$\begin{aligned}
 & \frac{1}{n-m} \int_m^n \zeta(\tau) d\tau = L_p^p(m, n), \\
 & \frac{\zeta(n) - \zeta(m)}{n-m} = p L_{p-1}^{p-1}(m, n), \\
 & \frac{\zeta(m) + \zeta(n)}{2} = A(m^p, n^p), \\
 & \frac{m+n}{2} = A
 \end{aligned}$$

$$\text{and} \quad \frac{1}{n-m} \|\zeta'\|_2^2 = \frac{1}{n-m} \int_m^n |\zeta'(\tau)|^2 d\tau = p^2 L_{2(p-1)}^{2(p-1)},$$

where $z \in [m + \lambda \frac{n-m}{2}, \frac{m+n}{2}]$.

Therefore, (2.1) becomes

$$\begin{aligned}
 & \left| (1-\lambda) \frac{z^{p+(m+n-z)^p} + \lambda A(m^p, n^p) - L_p^p}{2} \right| \\
 & \leq |p| \left[\frac{(n-m)^2}{12} (3\lambda^2 - 3\lambda + 1) + (z-A)^2 (1-\lambda) + \frac{(n-m)(1-\lambda)^2}{2} (z-A) \right]^{\frac{1}{2}} \left[L_{2(p-1)}^{2(p-1)} - L_{(p-1)}^{2(p-1)} \right]^{\frac{1}{2}}.
 \end{aligned} \tag{4.1}$$

Choose $z = A$ in (4.1), get

$$\left| (1-\lambda) A^p + \lambda A(m^p, n^p) - L_p^p \right| \leq |p| \frac{n-m}{2\sqrt{3}} (3\lambda^2 - 3\lambda + 1)^{\frac{1}{2}} \left[L_{2(p-1)}^{2(p-1)} - L_{(p-1)}^{2(p-1)} \right]^{\frac{1}{2}},$$

which is minimum for $\lambda = \frac{1}{2}$. Moreover for $\lambda = 1$

$$\left| A(m^p, n^p) - L_p^p \right| \leq \frac{n-m}{2\sqrt{3}} |p| \left[L_{2(p-1)}^{2(p-1)} - L_{(p-1)}^{2(p-1)} \right]^{\frac{1}{2}}.$$

Example 4.2. Consider $\zeta(z) = \frac{1}{z}, z \neq 0$, then

$$\begin{aligned}
 & \frac{1}{n-m} \int_m^n \zeta(\tau) d\tau = L^{-1}(m, n), \\
 & \frac{\zeta(n) - \zeta(m)}{n-m} = -\frac{1}{G^2}, \\
 & \frac{\zeta(m) + \zeta(n)}{2} = \frac{A}{G^2}, \\
 & \frac{1}{n-m} \int_m^n |\zeta'(\tau)|^2 d\tau = \frac{m^2 + mn + n^2}{3m^2 n^2}
 \end{aligned}$$

$$\text{and} \quad \frac{1}{n-m} \int_m^n |\zeta'(\tau)|^2 d\tau - \left(\frac{\zeta(n) - \zeta(m)}{n-m} \right)^2 = \frac{(n-m)^2}{3m^2 n^2} = \frac{(n-m)^2}{3G^6},$$

where $z \in [m + \lambda \frac{n-m}{2}, \frac{m+n}{2}] \subset (0, \infty)$.

Therefore, (2.1) becomes

$$\begin{aligned} & \left| \frac{(1-\lambda)}{2} \left(\frac{1}{z} + \frac{1}{(m+n-z)} \right) + \lambda \frac{A}{G^2} - \frac{1}{L} \right| \\ \leq & \left[\frac{(n-m)^2}{12} (3\lambda^2 - 3\lambda + 1) + (z-A)^2 (1-\lambda) + \frac{(n-m)(1-\lambda)^2}{2} (z-A) \right]^{\frac{1}{2}} \frac{(n-m)}{\sqrt{3}G^3}. \end{aligned} \tag{4.2}$$

If we choose $z = A$ in (4.2), we get

$$\left| (1-\lambda) \frac{1}{A} + \lambda \frac{A}{G^2} - \frac{1}{L} \right| \leq \frac{(n-m)^2}{6G^3} (3\lambda^2 - 3\lambda + 1)^{\frac{1}{2}}.$$

For $\lambda = 1$

$$\left| \frac{A}{G^2} - \frac{1}{L} \right| \leq \frac{(n-m)^2}{6G^3}.$$

Example 4.3. Consider $\zeta(z) = \ln z, z > 0$, then

$$\begin{aligned} \frac{1}{n-m} \int_m^n \zeta(\tau) d\tau &= \ln(I(m, n)), \\ \frac{\zeta(n) - \zeta(m)}{n-m} &= \frac{1}{L}, \\ \frac{\zeta(m) + \zeta(n)}{2} &= \ln G, \end{aligned}$$

$$\frac{1}{n-m} \int_m^n |\zeta'(\tau)|^2 dt = \frac{1}{G^2} \text{ and}$$

$$\frac{1}{n-m} \int_m^n |\zeta'(\tau)|^2 dt - \left(\frac{\zeta(n) - \zeta(m)}{n-m} \right)^2 = \frac{L^2 - G^2}{L^2 G^2},$$

where $z \in [m + \lambda \frac{n-m}{2}, \frac{m+n}{2}] \subset (0, \infty)$.

Therefore, (2.1) becomes

$$\begin{aligned} & \left| \ln \left(\frac{(z(m+n-z))^{\frac{(1-\lambda)}{2}} G^2}{I} \right) \right| \\ \leq & \left[\frac{(n-m)^2}{12} (3\lambda^2 - 3\lambda + 1) + (z-A)^2 (1-\lambda) + \frac{(n-m)(1-\lambda)^2}{2} (z-A) \right]^{\frac{1}{2}} \frac{(L^2 - G^2)^{\frac{1}{2}}}{LG}. \end{aligned}$$

For $z = A$

$$\left| \ln \left(\frac{A^{(1-\lambda)} G^2}{I} \right) \right| \leq \frac{(n-m)}{2\sqrt{3}LG} \left((3\lambda^2 - 3\lambda + 1)(L^2 - G^2) \right)^{\frac{1}{2}}.$$

For $\lambda = 1$

$$\left| \ln \left(\frac{G}{I} \right) \right| \leq \frac{(n-m)}{2\sqrt{3}LG} (L^2 - G^2)^{\frac{1}{2}}.$$

5. Application to numerical integration

To get the composite quadrature rules, we have to let $I_j : m = z_0 < z_1 < \dots < z_{j-1} < z_j = n$ be the partition of the interval $[m, n]$, $h_j = z_{j+1} - z_j, \lambda \in [0, 1], z_j + \lambda \frac{h_j}{2} \leq \eta_j \leq \frac{z_j + z_{j+1}}{2}, j \in \{0, \dots, i-1\}$, then the following results hold:

Theorem 5.1. If $\gamma \leq \zeta'(\tau) \leq \Gamma$ almost everywhere for $\tau \in [z_j + \lambda \frac{h_j}{2}, z_{j+1}] (j \in \{0, \dots, i-1\})$, then under the assumptions of Theorem 2.1 the following quadrature formula holds

$$\int_m^n \zeta(\tau) d\tau = Q(\zeta, \zeta', I_j, \eta, \lambda) + R(\zeta, \zeta', I_j, \eta, \lambda), \tag{5.1}$$

where

$$Q(\zeta, \zeta', I_j, \eta, \lambda) = \sum_{j=0}^{i-1} h_j \left[(1-\lambda) \frac{\zeta(\eta_j) + \zeta(z_j + z_{j+1} - \eta_j)}{2} + \lambda \frac{\zeta(z_j) + \zeta(z_{j+1})}{2} \right] \tag{5.2}$$

and remainder R satisfies the estimate

$$\begin{aligned} |R(\zeta, \zeta', I_j, \eta, \lambda)| &\leq \sum_{j=0}^{i-1} \left[\frac{h_j^2}{12} (3\lambda^2 - 3\lambda + 1) + \left(\eta_j - \frac{z_j + z_{j+1}}{2} \right)^2 (1-\lambda) \right. \\ &\quad \left. + \frac{h_j(1-\lambda)^2}{2} \left(\eta_j - \frac{z_j + z_{j+1}}{2} \right) \right]^{\frac{1}{2}} \left[h_j \|\zeta'\|_2^2 - (\zeta(z_{j+1}) - \zeta(z_j))^2 \right]^{\frac{1}{2}} \\ &\leq \frac{1}{2} (\Gamma - \gamma) \sum_{j=0}^{i-1} h_j \left[\frac{h_j^2}{12} (3\lambda^2 - 3\lambda + 1) + \left(\eta_j - \frac{z_j + z_{j+1}}{2} \right)^2 (1-\lambda) + \frac{h_j(1-\lambda)^2}{2} \left(\eta_j - \frac{z_j + z_{j+1}}{2} \right) \right]^{\frac{1}{2}}. \end{aligned} \tag{5.3}$$

Proof. By using inequalities (2.4), (2.5) and (2.9) on $z_j + \lambda \frac{h_j}{2} \leq \eta_j \leq \frac{z_j + z_{j+1}}{2}$ and summing over j from 0 to $i-1$, then we get required result. \square

By putting several values of λ and by fixing $\eta_j = \frac{z_j + z_{j+1}}{2}$, under the assumptions of Theorem 5.1 following results (special cases) are valid.

Special Case I: Put $\lambda = 1$ in (5.2) and (5.3), we have

$$Q\left(\zeta, \zeta', I_j, \frac{z_j + z_{j+1}}{2}, 1\right) = \frac{1}{2} \sum_{j=0}^{i-1} h_j (\zeta(z_j) + \zeta(z_{j+1}))$$

and

$$\begin{aligned} & |R\left(\zeta, \zeta', I_j, \frac{z_j + z_{j+1}}{2}, 1\right)| \\ \leq & \frac{1}{2\sqrt{3}} \sum_{j=0}^{i-1} h_j \left[h_j \|\zeta'\|_2^2 - (\zeta(z_{j+1}) - \zeta(z_j))^2 \right]^{\frac{1}{2}} \leq \frac{1}{4\sqrt{3}} (\Gamma - \gamma) \sum_{j=0}^{i-1} h_j^2. \end{aligned}$$

Special Case II: Put $\lambda = 0$ in (5.2) and (5.3), we have

$$Q\left(\zeta, \zeta', I_j, \frac{z_j + z_{j+1}}{2}\right) = \sum_{j=0}^{i-1} h_j \zeta\left(\frac{z_j + z_{j+1}}{2}\right)$$

and

$$\begin{aligned} & |R\left(\zeta, \zeta', I_j, \frac{z_j + z_{j+1}}{2}\right)| \\ \leq & \frac{1}{2\sqrt{3}} \sum_{j=0}^{i-1} h_j \left[h_j \|\zeta'\|_2^2 - (\zeta(z_{j+1}) - \zeta(z_j))^2 \right]^{\frac{1}{2}} \leq \frac{1}{4\sqrt{3}} (\Gamma - \gamma) \sum_{j=0}^{i-1} h_j^2. \end{aligned}$$

Special Case III: Put $\lambda = \frac{1}{2}$ in (5.2) and (5.3), we have

$$Q\left(\zeta, \zeta', I_j, \frac{z_j + z_{j+1}}{2}, \frac{1}{2}\right) = \frac{1}{4} \sum_{j=0}^{i-1} h_j \left(\zeta(z_j) + 2\zeta\left(\frac{z_j + z_{j+1}}{2}\right) + \zeta(z_{j+1}) \right)$$

and

$$\begin{aligned} & |R\left(\zeta, \zeta', I_j, \frac{z_j + z_{j+1}}{2}, \frac{1}{2}\right)| \\ \leq & \frac{1}{4\sqrt{3}} \sum_{j=0}^{i-1} h_j \left[h_j \|\zeta'\|_2^2 - (\zeta(z_{j+1}) - \zeta(z_j))^2 \right]^{\frac{1}{2}} \leq \frac{1}{8\sqrt{3}} (\Gamma - \gamma) \sum_{j=0}^{i-1} h_j^2. \end{aligned}$$

Special Case IV: Put $\lambda = \frac{1}{3}$ in (5.2) and (5.3), we have

$$Q\left(\zeta, \zeta', I_j, \frac{z_j + z_{j+1}}{2}, \frac{1}{3}\right) = \frac{1}{6} \sum_{j=0}^{i-1} h_j \left(\zeta(z_j) + 4\zeta\left(\frac{z_j + z_{j+1}}{2}\right) + \zeta(z_{j+1}) \right)$$

and

$$\begin{aligned} & |R\left(\zeta, \zeta', I_j, \frac{z_j + z_{j+1}}{2}, \frac{1}{3}\right)| \\ \leq & \frac{1}{6} \sum_{j=0}^{i-1} h_j \left[h_j \|\zeta'\|_2^2 - (\zeta(z_{j+1}) - \zeta(z_j))^2 \right]^{\frac{1}{2}} \leq \frac{1}{12} (\Gamma - \gamma) \sum_{j=0}^{i-1} h_j^2. \end{aligned}$$

6. Conclusion

Using three step kernel, we have obtained new generalized Ostrowski-Grüss type inequalities (2.1) which is a variant of (1.4) which was obtained in article (Barnett et al., 2000). By fixing

$z = \frac{m+n}{2}$ and by choosing different values of parameter λ we captured many results stated in Barnett et al. (2000) and Zafar (2010). We also got different important results from our main results as special cases such as trapezoidal inequality, mid-point inequality, averaged mid-point and trapezoidal inequality and Simpson's inequality. Moreover, applications are deduced for probability theory, special means and numerical integration.

Conflict of Interest: Authors declared: No conflict of interest.

Acknowledgment

The research of the Second Author is financially supported by the Dean's Science Research Grant 2021-2022, University of Karachi, with grant number, KURP Award Letter No. 161/2021-22.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <https://doi.org/10.1016/j.jksus.2022.102057>.

References

- Anastassiou, G.A., 1997. Multivariate Ostrowski type inequalities. *Acta. Math. Hungar.* 76, 267–278.
- Barnett, N.S., Dragomir, S.S., Sofo, A., 2001. Better bounds for an inequality of Ostrowski type with applications, Preprint RGMIA Res. Rep. Coll. 3 no. 1, (2000), Article 11, *Demonstratio Math.* 34(3), 533–542.
- Cheng, X.L., 2001. Improvement of some Ostrowski-Grüss type inequalities. *Comput. Math. Appl.* 42 (1/2), 109–114.
- Dragomir, S.S., Wang, S., 1997. An inequality of Ostrowski-Grüss type and its applications to the estimation of error bounds for some special means and for some numerical quadrature rules. *Comput. Math. Appl.* 33 (11), 16–20.
- Grüss, G., 1935. Über das Maximum des Absoluten Betrages von $[1/(n-m)] \int_m^n f(x)g(x) dx - [1/(n-m)^2] \int_m^n f(x) dx \int_m^n g(x) dx$. *Math. Z.* 39 (1), 215–226.
- Irshad, Nazia, Khan, Asif R., 2017. Some Applications of Quadrature Rules for Mappings on $L_p[u, v]$ Space via Ostrowski-type Inequality. *J. Num. Anal. Approx. Theory* 46(2), 141–149.
- Liu, Zheng, 2008. Some Ostrowski type inequalities. *Math. Comput. Model.* 48, 949–960.
- Matić, M., Pecarić, J.E., Ujević, N., 2000. Improvement and further generalization of some inequalities of Ostrowski-Grüss type. *Comput. Math. Appl.* 39 (3/4), 161–175.
- Matić, M., Pečarić, J.E., Ujević, N., 2000. Improvement and further generalization of inequalities of Ostrowski-Grüss type. *Comput. Math. Appl.* 39, 161–175.
- Milovanovic, G.V., Pecarić, J.E., 1976. On generalization of the inequality of A. Ostrowski and some related applications, *Univ. Beograd Publ., Elektrotehn. Fak. Ser. Mat. Fiz.* No 544–576, 155–158.
- Mitrinović, D.S., Pečarić, J.E., Fink, A.M., 1994. *Inequalities for Functions and Their Integrals and Derivatives*. Kluwer Academic Publishers, Dordrecht.
- Shaikh, Muhammad Awais, Khan, Asif R., Mehmood, Faraz. Estimates for Weighted Ostrowski-Grüss Type Inequalities with Applications, *Analysis-De Gruyter*, to appear.
- Ostrowski, A., 1938. Über die Absolutabweichung einer differentienbaren Funktionen von ihren Integralmittelwert. *Comment. Math. Helv.* 10, 226–227.
- Zafar, F., 2010. *Some Generalizations of Ostrowski Inequalities and Their Applications to Numerical Integration and Special means* (PhD Dissertation). Bahauddin Zakariya University Multan, Pakistan.
- Zafar, Fiza, Mir, Nazir Ahmad, 2010. A generalization of Ostrowski-Grüss type inequality for first differentiable mappings. *Tamsui Oxford J. Math. Sci.* 26(1), 61–76.