



**VICTORIA UNIVERSITY**  
MELBOURNE AUSTRALIA

*Reverses of the first Hermite-Hadamard type inequality for the square operator modulus in Hilbert spaces*

This is the Published version of the following publication

Dragomir, Sever S (2022) Reverses of the first Hermite-Hadamard type inequality for the square operator modulus in Hilbert spaces. *Journal of Linear and Topological Algebra*, 11 (1). pp. 1-13. ISSN 2252-0201

The publisher's official version can be found at  
[https://jlta.ctb.iau.ir/article\\_688390.html](https://jlta.ctb.iau.ir/article_688390.html)

Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/46590/>

## Reverses of the first Hermite-Hadamard type inequality for the square operator modulus in Hilbert spaces

S. S. Dragomir<sup>a,b</sup>

<sup>a</sup>Mathematics, College of Engineering & Science, Victoria University, PO Box 14428  
Melbourne City, MC 8001, Australia

<sup>b</sup>DST-NRF Centre of Excellence in the Mathematical and Statistical Sciences, School of Computer  
Science & Applied Mathematics, University of the Witwatersrand, Private Bag 3  
Johannesburg 2050, South Africa.

Received 30 November 2021; Revised 10 January 2022; Accepted 16 January 2022.

Communicated by Ghasem Soleimani Rad

---

**Abstract.** Let  $(H; \langle \cdot, \cdot \rangle)$  be a complex Hilbert space. Denote by  $\mathcal{B}(H)$  the Banach  $C^*$ -algebra of bounded linear operators on  $H$ . For  $A \in \mathcal{B}(H)$  we define the modulus of  $A$  by  $|A| := (A^*A)^{1/2}$  and  $\operatorname{Re} A := \frac{1}{2}(A^* + A)$ . In this paper we show among other that, if  $A, B \in \mathcal{B}(H)$  with  $0 \leq m \leq |(1-t)A + tB|^2 \leq M$  for all  $t \in [0, 1]$ , then

$$\begin{aligned} 0 &\leq \int_0^1 f(|(1-t)A + tB|^2) dt - f\left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right) \\ &\leq 2 \left[ \frac{f(m) + f(M)}{2} - f\left(\frac{m+M}{2}\right) \right] 1_H \end{aligned}$$

for operator convex functions  $f : [0, \infty) \rightarrow \mathbb{R}$ . Applications for power and logarithmic functions are also provided.

---

**Keywords:** Operator convex functions, Hermite-Hadamard inequality, midpoint inequality, operator power and logarithmic functions.

**2010 AMS Subject Classification:** 47A63, 26D15, 46C05.

---

E-mail address: sever.dragomir@vu.edu.au (S. S. Dragomir).

## 1. Introduction

Let  $\mathbb{R}$  be the set of real numbers and  $I \subseteq \mathbb{R}$  be an interval. A function  $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$  is said to be *convex* in the classical sense if it satisfies the following inequality

$$f(tx + (1-t)y) \leq tf(x) + (1-t)f(y)$$

for all  $x, y \in I$  and  $t \in [0, 1]$ .

Let  $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$  and  $a, b \in I$  with  $a < b$ . Then the inequality

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq \frac{f(a)+f(b)}{2} \quad (1)$$

holds if  $f$  is convex, and is known in the literature as *Hermite-Hadamard inequality*, after the name of C. Hermite and J. Hadamard (see [17]). The inequalities in (1) hold in reversed direction if  $f$  is a concave function.

A vast literature related to (1) have been produced by a large number of mathematicians [13] since it is considered to be one of the most famous inequality for convex functions due to its usefulness and many applications in various branches of Pure and Applied Mathematics, such as Numerical Analysis [3], Information Theory [2], Operator Theory [9, 10] and others.

Let  $X$  be a vector space over the real or complex number field  $\mathbb{K}$  and  $x, y \in X$ ,  $x \neq y$ . Define the segment  $[x, y] := \{(1-t)x + ty, t \in [0, 1]\}$ . We consider the function  $f : [x, y] \rightarrow \mathbb{R}$  and the associated function  $g(x, y) : [0, 1] \rightarrow \mathbb{R}$  with  $g(x, y)(t) := f[(1-t)x + ty]$  for  $t \in [0, 1]$ . Note that  $f$  is convex on  $[x, y]$  if and only if  $g(x, y)$  is convex on  $[0, 1]$ .

For any convex function defined on a segment  $[x, y] \subset X$ , we have the *Hermite-Hadamard integral inequality* (see [4, p. 2], [5, p. 2])

$$f\left(\frac{x+y}{2}\right) \leq \int_0^1 f[(1-t)x + ty]dt \leq \frac{f(x)+f(y)}{2}, \quad (2)$$

which can be derived from the classical Hermite-Hadamard inequality (1) for the convex function  $g(x, y) : [0, 1] \rightarrow \mathbb{R}$ .

Since  $f(x) = \|x\|^p$  ( $x \in X$  and  $1 \leq p < \infty$ ) is a convex function, then for any  $x, y \in X$  we have the following norm inequality from (2) (see [16, p. 106])

$$\left\|\frac{x+y}{2}\right\|^p \leq \int_0^1 \|(1-t)x + ty\|^p dt \leq \frac{\|x\|^p + \|y\|^p}{2}. \quad (3)$$

A real valued continuous function  $f$  on an interval  $I$  is said to be *operator convex* (*operator concave*) on  $I$  if

$$f((1-\lambda)A + \lambda B) \leq (\geq) (1-\lambda)f(A) + \lambda f(B) \quad (4)$$

in the operator order, for all  $\lambda \in [0, 1]$  and for every selfadjoint operator  $A$  and  $B$  on a Hilbert space  $H$  whose spectra are contained in  $I$ . Notice that a function  $f$  is operator concave if  $-f$  is operator convex.

A real valued continuous function  $f$  on an interval  $I$  is said to be *operator monotone* if it is monotone with respect to the operator order, i.e.,  $A \leq B$  with  $\text{Sp}(A), \text{Sp}(B) \subset I$  imply  $f(A) \leq f(B)$ . As examples of such functions, we note that  $f(t) = t^r$  is operator

monotone on  $[0, \infty)$  if and only if  $0 \leq r \leq 1$ . The function  $f(t) = t^r$  is operator convex on  $(0, \infty)$  if either  $1 \leq r \leq 2$  or  $-1 \leq r \leq 0$  and is operator concave on  $(0, \infty)$  if  $0 \leq r \leq 1$ . The logarithmic function  $f(t) = \ln t$  is operator monotone and operator concave on  $(0, \infty)$ . The entropy function  $f(t) = -t \ln t$  is operator concave on  $(0, \infty)$ . The exponential function  $f(t) = e^t$  is neither operator convex nor operator monotone.

In the recent paper [12], we obtained among others the following Hermite-Hadamard type inequalities for operator convex functions:

**Theorem 1.1** Let  $f : I \rightarrow \mathbb{R}$  be an operator convex function on the interval  $I$ . Then for any selfadjoint operators  $A$  and  $B$  with spectra in  $I$  and for any  $\lambda \in [0, 1]$  we have the inequalities

$$\begin{aligned} f\left(\frac{A+B}{2}\right) &\leq (1-\lambda)f\left[\frac{(1-\lambda)A+(1+\lambda)B}{2}\right] + \lambda f\left[\frac{(2-\lambda)A+\lambda B}{2}\right] \quad (5) \\ &\leq \int_0^1 f((1-s)A+sB) ds \\ &\leq \frac{1}{2}[f((1-\lambda)A+\lambda B) + (1-\lambda)f(B) + \lambda f(A)] \\ &\leq \frac{f(A)+f(B)}{2}. \end{aligned}$$

A similar result and with a different proof was obtained by B. Li in [15]. For  $\lambda = \frac{1}{2}$  in (5) we recapture the result obtained in the earlier paper [11] by the author. For other similar inequalities for operator convex functions see [1, 18–22].

Denote by  $\mathcal{B}(H)$  the Banach  $C^*$ -algebra of bounded linear operators on Hilbert space  $H$ . For  $A \in \mathcal{B}(H)$  we define the modulus of  $A$  by  $|A| := (A^*A)^{1/2}$ . It is well known that the modulus of operators does not satisfy, in general, the triangle inequality  $|A+B| \leq |A|+|B|$ , so the classical arguments using this inequality can not be used.

In the recent paper [14], we obtained among others the following Hermite-Hadamard type inequalities for operator convex (concave) functions  $f : [0, \infty) \rightarrow \mathbb{R}$  and  $A, B \in \mathcal{B}(H)$  with  $\operatorname{Re}(B^*A) \geq 0$ ,

$$\begin{aligned} f\left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right) &\leq (\geq) \int_0^1 f(|(1-t)A+tB|^2) dt \\ &\leq (\geq) \frac{1}{3} \left[ f(|A|^2) + f[\operatorname{Re}(B^*A)] + f(|B|^2) \right]. \end{aligned}$$

Some examples for power functions and logarithm were also provided.

In this paper, we show among other that, if  $A, B \in \mathcal{B}(H)$  with  $0 \leq m \leq |(1-t)A+tB|^2 \leq M$  for all  $t \in [0, 1]$ , then

$$\begin{aligned} 0 &\leq \int_0^1 f(|(1-t)A+tB|^2) dt - f\left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right) \\ &\leq 2 \left[ \frac{f(m)+f(M)}{2} - f\left(\frac{m+M}{2}\right) \right] 1_H \end{aligned}$$

for operator convex functions  $f : [0, \infty) \rightarrow \mathbb{R}$ . Applications for power and logarithmic

functions are also provided.

## 2. Main Results

The first main result is as follows:

**Theorem 2.1** Let  $f : [0, \infty) \rightarrow \mathbb{R}$  be operator convex on  $[0, \infty)$ . If  $A, B \in \mathcal{B}(H)$  with

$$0 \leq m \leq |(1-t)A + tB|^2 \leq M$$

for all  $t \in [0, 1]$ , then

$$\begin{aligned} 0 &\leq \int_0^1 f\left(|(1-t)A + tB|^2\right) dt - f\left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right) \\ &\leq f(m) \frac{M - \frac{1}{3}\left[|A|^2 + \operatorname{Re}(B^*A) + |B|^2\right]}{M - m} \\ &\quad + f(M) \frac{\frac{1}{3}\left[|A|^2 + \operatorname{Re}(B^*A) + |B|^2\right] - m}{M - m} - f\left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right) \\ &\leq 2 \left[ \frac{f(m) + f(M)}{2} - f\left(\frac{m + M}{2}\right) \right]. \end{aligned} \tag{6}$$

**Proof.** We use the double inequality, see [6],

$$\begin{aligned} &2 \min\{\alpha, 1 - \alpha\} \left[ \frac{f(m) + f(M)}{2} - f\left(\frac{m + M}{2}\right) \right] \\ &\leq (1 - \alpha) f(m) + \alpha f(M) - f((1 - \alpha)m + \alpha M) \\ &\leq 2 \max\{\alpha, 1 - \alpha\} \left[ \frac{f(m) + f(M)}{2} - f\left(\frac{m + M}{2}\right) \right] \end{aligned} \tag{7}$$

for  $\alpha \in [0, 1]$ .

Let  $x \in [m, M]$  and take  $\alpha = \frac{x-m}{M-m} \in [0, 1]$  to get

$$\begin{aligned} &2 \min\left\{ \frac{x-m}{M-m}, \frac{M-x}{M-m} \right\} \left[ \frac{f(m) + f(M)}{2} - f\left(\frac{m + M}{2}\right) \right] \\ &\leq \frac{M-x}{M-m} f(m) + \frac{x-m}{M-m} f(M) - f\left(\frac{M-x}{M-m}m + \frac{x-m}{M-m}M\right) \\ &\leq 2 \max\left\{ \frac{x-m}{M-m}, \frac{M-x}{M-m} \right\} \left[ \frac{f(m) + f(M)}{2} - f\left(\frac{m + M}{2}\right) \right], \end{aligned}$$

namely

$$\begin{aligned}
 0 &\leq \frac{2}{M-m} \left\{ \frac{1}{2} (M-m) - \left| x - \frac{m+M}{2} \right| \right\} \times \left[ \frac{f(m)+f(M)}{2} - f\left(\frac{m+M}{2}\right) \right] \quad (8) \\
 &\leq \frac{M-x}{M-m} f(m) + \frac{x-m}{M-m} f(M) - f(x) \\
 &\leq \frac{2}{M-m} \left\{ \frac{1}{2} (M-m) + \left| x - \frac{m+M}{2} \right| \right\} \times \left[ \frac{f(m)+f(M)}{2} - f\left(\frac{m+M}{2}\right) \right].
 \end{aligned}$$

If we use the continuous functional calculus for selfadjoint operators and the inequality (8) we get for  $m1_H \leq T \leq M1_H$

$$\begin{aligned}
 0 &\leq \frac{2}{M-m} \left\{ \frac{1}{2} (M-m) 1_H - \left| T - \frac{m+M}{2} \right| \right\} \quad (9) \\
 &\times \left[ \frac{f(m)+f(M)}{2} - f\left(\frac{m+M}{2}\right) \right] \\
 &\leq f(m) \frac{M-T}{M-m} + f(M) \frac{T-m}{M-m} - f(T) \\
 &\leq \frac{2}{M-m} \left\{ \frac{1}{2} (M-m) 1_H + \left| T - \frac{m+M}{2} \right| \right\} \\
 &\times \left[ \frac{f(m)+f(M)}{2} - f\left(\frac{m+M}{2}\right) \right].
 \end{aligned}$$

So, if we take  $T = \frac{|A|^2 + \text{Re}(B^*A) + |B|^2}{3}$  in (9), then we get the inequality of interest

$$\begin{aligned}
 0 &\leq \frac{2}{M-m} \left[ \frac{f(m)+f(M)}{2} - f\left(\frac{m+M}{2}\right) \right] \quad (10) \\
 &\times \left\{ \frac{1}{2} (M-m) 1_H - \left| \frac{|A|^2 + \text{Re}(B^*A) + |B|^2}{3} - \frac{m+M}{2} \right| \right\} \\
 &\leq f(m) \frac{M - \frac{1}{3} [ |A|^2 + \text{Re}(B^*A) + |B|^2 ]}{M-m} \\
 &+ f(M) \frac{\frac{1}{3} [ |A|^2 + \text{Re}(B^*A) + |B|^2 ] - m}{M-m} - f\left(\frac{|A|^2 + \text{Re}(B^*A) + |B|^2}{3}\right) \\
 &\leq \frac{2}{M-m} \left[ \frac{f(m)+f(M)}{2} - f\left(\frac{m+M}{2}\right) \right] \\
 &\times \left\{ \frac{1}{2} (M-m) 1_H + \left| \frac{|A|^2 + \text{Re}(B^*A) + |B|^2}{3} - \frac{m+M}{2} \right| \right\} \\
 &\leq 2 \left[ \frac{f(m)+f(M)}{2} - f\left(\frac{m+M}{2}\right) \right].
 \end{aligned}$$

From (9) we also get

$$f(T) \leq f(m) \frac{M-T}{M-m} + f(M) \frac{T-m}{M-m},$$

which gives for  $T = |(1-t)A + tB|^2$ ,  $t \in [0, 1]$  that

$$\begin{aligned} & f\left(|(1-t)A + tB|^2\right) \\ & \leq f(m) \frac{M - |(1-t)A + tB|^2}{M-m} + f(M) \frac{|(1-t)A + tB|^2 - m}{M-m}, \end{aligned}$$

$t \in [0, 1]$ .

By taking the integral and by observing also that

$$\begin{aligned} & \int_0^1 |(1-t)A + tB|^2 dt \\ & = \int_0^1 \left[ (1-t)^2 |A|^2 + 2t(1-t) \operatorname{Re}(B^*A) + t^2 |B|^2 \right] dt \\ & = \left( \int_0^1 (1-t)^2 dt \right) |A|^2 + 2 \left( \int_0^1 t(1-t) dt \right) \operatorname{Re}(B^*A) + \left( \int_0^1 t^2 dt \right) |B|^2 \\ & = \frac{1}{3} \left[ |A|^2 + \operatorname{Re}(B^*A) + |B|^2 \right], \end{aligned}$$

we get

$$\begin{aligned} \int_0^1 f\left(|(1-t)A + tB|^2\right) dt & \leq f(m) \frac{M - \frac{1}{3} \left[ |A|^2 + \operatorname{Re}(B^*A) + |B|^2 \right]}{M-m} \\ & \quad + f(M) \frac{\frac{1}{3} \left[ |A|^2 + \operatorname{Re}(B^*A) + |B|^2 \right] - m}{M-m}. \end{aligned} \quad (11)$$

By subtracting in (11)  $f\left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right)$ , then we get

$$\begin{aligned} & \int_0^1 f\left(|(1-t)A + tB|^2\right) dt - f\left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right) \\ & \leq f(m) \frac{M - \frac{1}{3} \left[ |A|^2 + \operatorname{Re}(B^*A) + |B|^2 \right]}{M-m} \\ & \quad + f(M) \frac{\frac{1}{3} \left[ |A|^2 + \operatorname{Re}(B^*A) + |B|^2 \right] - m}{M-m} - f\left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right). \end{aligned} \quad (12)$$

By (12) and (10) we derive the desired result (6). ■

**Corollary 2.2** Let  $f : [0, \infty) \rightarrow \mathbb{R}$  be operator convex on  $[0, \infty)$  with  $f(0) = 0$ . If  $A, B \in \mathcal{B}(H)$  with  $|A|^2, |B|^2 \leq M$ , where  $M > 0$ , then

$$\begin{aligned} 0 &\leq \int_0^1 f\left(|(1-t)A + tB|^2\right) dt - f\left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right) \\ &\leq \frac{f(M)}{M} \left[\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right] - f\left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right) \\ &\leq 2 \left[\frac{1}{2}f(M) - f\left(\frac{1}{2}M\right)\right]. \end{aligned} \tag{13}$$

**Proof.** The following Cauchy-Bunyakowsky-Schwarz inequality holds

$$\sum_{k=1}^n w_k |z_k|^2 \sum_{k=1}^n w_k |A_k|^2 \geq \left| \sum_{k=1}^n w_k z_k A_k \right|^2, \tag{14}$$

where  $z_k \in \mathbb{C}, A_k \in \mathcal{B}(H), w_k \geq 0$  for  $k \in \{1, \dots, n\}$  and  $\sum_{k=1}^n w_k = 1$ .

Let  $A, B \in \mathcal{B}(H)$  and  $t \in [0, 1]$ . Then by (14), we get

$$\begin{aligned} |(1-t)A + tB|^2 &= \left| (1-t)^{1/2} (1-t)^{1/2} A + t^{1/2} t^{1/2} B \right|^2 \\ &\leq \left[ \left( (1-t)^{1/2} \right)^2 + \left( t^{1/2} \right)^2 \right] \left[ \left| (1-t)^{1/2} A \right|^2 + \left| t^{1/2} B \right|^2 \right] \\ &= (1-t+t) \left[ (1-t)|A|^2 + t|B|^2 \right] \\ &= (1-t)|A|^2 + t|B|^2. \end{aligned}$$

So, if  $|A|^2 \leq M$  and  $|B|^2 \leq M$ , then  $|(1-t)A + tB|^2 \leq (1-t)M + tM = M$  and by writing the inequality (6) for  $m = 0$  we derive (13). ■

We also have:

**Theorem 2.3** Let  $f : [0, \infty) \rightarrow \mathbb{R}$  be operator convex on  $[0, \infty)$ . If  $A, B \in \mathcal{B}(H)$  with  $0 \leq m \leq |(1-t)A + tB|^2 \leq M$  for all  $t \in [0, 1]$ , then

$$\begin{aligned} 0 &\leq \int_0^1 f\left(|(1-t)A + tB|^2\right) dt - f\left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right) \\ &\leq f(m) \frac{M1_H - \frac{1}{3} \left[ |A|^2 + \operatorname{Re}(B^*A) + |B|^2 \right]}{M - m} \\ &\quad + f(M) \frac{\frac{1}{3} \left[ |A|^2 + \operatorname{Re}(B^*A) + |B|^2 \right] - m1_H}{M - m} - f\left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right) \\ &\leq \frac{f'_-(M) - f'_+(m)}{M - m} \left( \frac{1}{3} \left[ |A|^2 + \operatorname{Re}(B^*A) + |B|^2 \right] - m1_H \right) \end{aligned} \tag{15}$$



$$\begin{aligned} & \times \left( M1_H - \frac{1}{3} \left[ |A|^2 + \operatorname{Re}(B^*A) + |B|^2 \right] \right) \\ & \leq \frac{1}{4} (M - m) [f'_-(M) - f'_+(m)] 1_H, \end{aligned}$$

provided that the lateral derivatives

$$f'_-(M) := \lim_{t \rightarrow M, t < M} \frac{f(t) - f(M)}{t - M}, \quad f'_+(m) := \lim_{t \rightarrow m, t > m} \frac{f(t) - f(m)}{t - m}$$

are finite.

**Proof.** We use the inequality, see [8]

$$(1 - \alpha) f(m) + \alpha f(M) - f((1 - \alpha)m + \alpha M) \leq \alpha(1 - \alpha)(M - m) [f'_-(M) - f'_+(m)]$$

for  $\alpha \in [0, 1]$ . Let  $x \in [m, M]$  and take  $\alpha = \frac{x-m}{M-m} \in [0, 1]$  to get

$$\frac{M-x}{M-m} f(m) + \frac{x-m}{M-m} f(M) - f(x) \leq \frac{f'_-(M) - f'_+(m)}{M-m} (x-m)(M-x). \quad (16)$$

If we use the continuous functional calculus for selfadjoint operators and the inequality (16) we get

$$\begin{aligned} & f(m) \frac{M1_H - T}{M-m} + f(M) \frac{T - m1_H}{M-m} - f(T) \\ & \leq \frac{f'_-(M) - f'_+(m)}{M-m} (T - m1_H)(M1_H - T). \end{aligned} \quad (17)$$

So, if we take  $T = \frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}$  in (17), then we get the inequality of interest

$$\begin{aligned} & f(m) \frac{M1_H - \frac{1}{3} \left[ |A|^2 + \operatorname{Re}(B^*A) + |B|^2 \right]}{M-m} \\ & + f(M) \frac{\frac{1}{3} \left[ |A|^2 + \operatorname{Re}(B^*A) + |B|^2 \right] - m1_H}{M-m} \\ & - f \left( \frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3} \right) \\ & \leq \frac{f'_-(M) - f'_+(m)}{M-m} \left( \frac{1}{3} \left[ |A|^2 + \operatorname{Re}(B^*A) + |B|^2 \right] - m1_H \right) \\ & \times \left( M1_H - \frac{1}{3} \left[ |A|^2 + \operatorname{Re}(B^*A) + |B|^2 \right] \right) \\ & \leq \frac{1}{4} (M - m) [f'_-(M) - f'_+(m)] 1_H \end{aligned} \quad (18)$$

From (12) and (18) we derive (15). ■

**Corollary 2.4** Let  $f : [0, \infty) \rightarrow \mathbb{R}$  be operator convex on  $[0, \infty)$  with  $f(0) = 0$ . If  $A, B \in \mathcal{B}(H)$  with  $|A|^2, |B|^2 \leq M$ , where  $M > 0$ , then

$$\begin{aligned} 0 &\leq \int_0^1 f(|(1-t)A + tB|^2) dt - f\left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right) \\ &\leq \frac{f(M)}{M} \left[\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right] - f\left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right) \\ &\leq \frac{f'_-(M) - f'_+(0)}{M} \left(\frac{1}{3} [|A|^2 + \operatorname{Re}(B^*A) + |B|^2]\right) \\ &\quad \times \left(M1_H - \frac{1}{3} [|A|^2 + \operatorname{Re}(B^*A) + |B|^2]\right) \\ &\leq \frac{1}{4} M [f'_-(M) - f'_+(0)] 1_H, \end{aligned} \tag{19}$$

provided that  $f'_+(0)$  is finite.

**Theorem 2.5** Let  $f : [0, \infty) \rightarrow \mathbb{R}$  be twice differentiable and so that there exists the constants  $0 \leq d, D$  such that  $d \leq f''(t) \leq D$  for any  $t \in (m, M) \subset [0, \infty)$ . If  $A, B \in \mathcal{B}(H)$  with  $0 \leq m \leq |(1-t)A + tB|^2 \leq M$  for all  $t \in [0, 1]$ , then

$$\begin{aligned} 0 &\leq \int_0^1 f(|(1-t)A + tB|^2) dt - f\left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right) \\ &\leq f(m) \frac{M1_H - \frac{1}{3} [|A|^2 + \operatorname{Re}(B^*A) + |B|^2]}{M - m} \\ &\quad + f(M) \frac{\frac{1}{3} [|A|^2 + \operatorname{Re}(B^*A) + |B|^2] - m1_H}{M - m} - f\left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right) \\ &\leq \frac{1}{2} D \left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3} - m1_H\right) \left(M1_H - \frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right) \\ &\leq \frac{1}{8} (M - m)^2 D. \end{aligned} \tag{20}$$

**Proof.** If there exists the constants  $0 \leq d, D$  such that

$$d \leq f''(t) \leq D \text{ for any } t \in (m, M),$$

then, see for instance [7],

$$\begin{aligned} \frac{1}{2} \nu (1 - \nu) d (M - m)^2 &\leq (1 - \nu) f(m) + \nu f(M) - f((1 - \nu)m + \nu M) \\ &\leq \frac{1}{2} \nu (1 - \nu) D (M - m)^2 \end{aligned} \tag{21}$$

for any  $\nu \in [0, 1]$ .

Let  $x \in [m, M]$  and take  $\alpha = \frac{x-m}{M-m} \in [0, 1]$  in (21) to get

$$\begin{aligned} \frac{1}{2}d(x-m)(M-x) &\leq \frac{M-x}{M-m}f(m) + \frac{x-m}{M-m}f(M) - f(x) \\ &\leq \frac{1}{2}D(x-m)(M-x). \end{aligned} \quad (22)$$

If we use the continuous functional calculus for selfadjoint operators and the inequality (22) we get

$$\begin{aligned} &\frac{1}{2}d(T - m1_H)(M1_H - T) \\ &\leq f(m) \frac{M1_H - T}{M - m} + f(M) \frac{T - m1_H}{M - m} - f(T) \\ &\leq \frac{1}{2}D(T - m1_H)(M1_H - T). \end{aligned} \quad (23)$$

So, if we take  $T = \frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}$  in (17), then we get the inequality of interest

$$\begin{aligned} &\frac{1}{2}d\left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3} - m1_H\right) \\ &\times \left(M1_H - \frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right) \\ &\leq f(m) \frac{M1_H - \frac{1}{3} \left[|A|^2 + \operatorname{Re}(B^*A) + |B|^2\right]}{M - m} \\ &+ f(M) \frac{\frac{1}{3} \left[|A|^2 + \operatorname{Re}(B^*A) + |B|^2\right] - m1_H}{M - m} \\ &- f\left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right) \\ &\leq \frac{1}{2}D\left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3} - m1_H\right) \\ &\times \left(M1_H - \frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right) \\ &\leq \frac{1}{8}(M - m)^2 D \end{aligned} \quad (24)$$

Now, if we use (12) and (24) we derive (20). ■

**Corollary 2.6** Let  $f : [0, \infty) \rightarrow \mathbb{R}$  be operator convex on  $[0, \infty)$  with  $f(0) = 0$ . If  $A$ ,

$B \in \mathcal{B}(H)$  with  $|A|^2, |B|^2 \leq M$ , where  $M > 0$ , then

$$\begin{aligned}
 0 &\leq \int_0^1 f(|(1-t)A + tB|^2) dt - f\left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right) \\
 &\leq \frac{f(M)}{M} \left[ \frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3} \right] - f\left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right) \\
 &\leq \frac{1}{2}D \left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right) \left(M1_H - \frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right) \\
 &\leq \frac{1}{8}M^2D,
 \end{aligned} \tag{25}$$

provided that  $f''(t) \leq D$  for any  $t \in (0, M)$ .

### 3. Some Examples

The function  $f(t) = t^p, p \in [1, 2]$  is operator convex on  $[0, \infty)$  with  $f(0) = 0$ . From (13) and (19) we derive

$$\begin{aligned}
 0 &\leq \int_0^1 |(1-t)A + tB|^{2p} dt - \left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right)^p \\
 &\leq M^{p-1} \left[ \frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3} \right] - \left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right)^p \\
 &\leq M^p \left(\frac{2^{p-1} - 1}{2^{p-1}}\right)
 \end{aligned} \tag{26}$$

and

$$\begin{aligned}
 0 &\leq \int_0^1 |(1-t)A + tB|^{2p} dt - \left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right)^p \\
 &\leq M^{p-1} \left[ \frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3} \right] - \left(\frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3}\right)^p \\
 &\leq pM^{p-2} \left(\frac{1}{3} [ |A|^2 + \operatorname{Re}(B^*A) + |B|^2 ]\right) \\
 &\quad \times \left( M1_H - \frac{1}{3} [ |A|^2 + \operatorname{Re}(B^*A) + |B|^2 ]\right) \\
 &\leq \frac{1}{4^p} pM^p 1_H,
 \end{aligned} \tag{27}$$

provided that  $A, B \in \mathcal{B}(H)$  with  $|A|^2, |B|^2 \leq M$ , where  $M > 0$ .

The function  $f(t) = t^r, r \in (0, 1)$  is operator concave on  $[0, \infty)$  with  $f(0) = 0$ . From

(13) we derive

$$\begin{aligned} 0 &\leq \left( \frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3} \right)^r - \int_0^1 |(1-t)A + tB|^{2r} dt \\ &\leq \left( \frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3} \right)^r - \frac{1}{M^{1-r}} \left[ \frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3} \right] \\ &\leq M^r (2^{1-r} - 1) \end{aligned} \quad (28)$$

and, in particular

$$\begin{aligned} 0 &\leq \left( \frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3} \right)^{1/2} - \int_0^1 |(1-t)A + tB| dt \\ &\leq \left( \frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3} \right)^{1/2} - \frac{1}{\sqrt{M}} \left[ \frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3} \right] \\ &\leq \sqrt{M} (\sqrt{2} - 1) \end{aligned} \quad (29)$$

provided that  $A, B \in \mathcal{B}(H)$  with  $|A|^2, |B|^2 \leq M$ , where  $M > 0$ .

Consider the operator concave function  $f(t) = \ln(t+1)$ . By (13) we get

$$\begin{aligned} 0 &\leq \ln \left( \frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3} + 1 \right) - \int_0^1 \ln \left( |(1-t)A + tB|^2 + 1 \right) dt \\ &\leq \ln \left( \frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3} + 1 \right) \\ &\quad - \frac{\ln(M+1)}{M} \left[ \frac{|A|^2 + \operatorname{Re}(B^*A) + |B|^2}{3} \right] \\ &\leq 2 \ln \left( \frac{1}{2}M + 1 \right) - \ln(M+1) \end{aligned} \quad (30)$$

provided that  $A, B \in \mathcal{B}(H)$  with  $|A|^2, |B|^2 \leq M$ , where  $M > 0$ .

## References

- [1] R. P. Agarwal, S. S. Dragomir, A survey of Jensen type inequalities for functions of selfadjoint operators in Hilbert spaces, *Comput. Math. Appl.* 59 (12) (2010), 3785-3812.
- [2] N. S. Barnett, P. Cerone, S. S. Dragomir, Some new inequalities for Hermite-Hadamard divergence in information theory, *Stochastic Analysis and Applications*, 3, Nova Sci. Publ., Hauppauge, NY, 2003.
- [3] P. Cerone, S. S. Dragomir, *Mathematical Inequalities. A Perspective*, CRC Press, Boca Raton, 2011.
- [4] S. S. Dragomir, An inequality improving the first Hermite-Hadamard inequality for convex functions defined on linear spaces and applications for semi-inner products, *J. Inequal. Pure Appl. Math.* 3 (2002), 2:31.
- [5] S. S. Dragomir, An inequality improving the second Hermite-Hadamard inequality for convex functions defined on linear spaces and applications for semi-inner products, *J. Inequal. Pure Appl. Math.* 3 (2002), 3:35.
- [6] S. S. Dragomir, Superadditivity of some functionals associated with Jensen's inequality for convex functions on linear spaces with applications, *Bull. Aust. Math. Soc.* 82 (1) (2010), 44-61.

- [7] S. S. Dragomir, A note on new refinements and reverses of Young's inequality, *Transylv. J. Math. Mech.* 8 (1) (2016), 45-49.
- [8] S. S. Dragomir, A note on Young's inequality, *RACSAM.* 111 (2) (2017), 349-354.
- [9] S. S. Dragomir, *Operator Inequalities of Ostrowski and Trapezoidal Type*, Springer Briefs in Mathematics, Springer, New York, 2012.
- [10] S. S. Dragomir, *Operator Inequalities of the Jensen, Čebyšev and Grüss Type*, Springer Briefs in Mathematics, Springer, New York, 2012.
- [11] S. S. Dragomir, Hermite-Hadamard's type inequalities for operator convex functions, *Appl. Math. Comput.* 218 (3) (2011), 766-772.
- [12] S. S. Dragomir, Some Hermite-Hadamard type inequalities for operator convex functions and positive maps, *Spec. Matrices.* 7 (2019), 38-51.
- [13] S. S. Dragomir, C. E. M. Pearce, *Selected Topics on Hermite-Hadamard Inequalities and Applications*, RGMIA Monographs, 2000.
- [14] S. S. Dragomir, Some Hermite-Hadamard type inequalities for the square operator modulus in Hilbert spaces, *Preprint RGMIA Res. Rep. Coll.* 24 (2021), 1-16.
- [15] B. Li, Refinements of Hermite-Hadamard's type inequalities for operator convex functions, *Int. J. Contemp. Math. Sci.* 8 (9-12) (2013), 463-467.
- [16] J. Pečarić, S. S. Dragomir, A generalization of Hadamard's inequality for isotonic linear functionals, *Radovi Mat. (Sarajevo).* 7 (1991), 103-107.
- [17] J. E. Pečarić, F. Proschan, Y. L. Tong, *Convex Functions, Partial Orderings and Statistical Applications (Mathematics in Science and Engineering)*, 187, Academic Press Inc., 1992.
- [18] A. Taghavi, V. Darvish, H. M. Nazari, S. S. Dragomir, Hermite-Hadamard type inequalities for operator geometrically convex functions, *Monatsh. Math.* 181 (1) (2016), 187-203.
- [19] M. Vivas Cortez, E. J. H. Hernández, Refinements for Hermite-Hadamard type inequalities for operator  $h$ -convex function, *Appl. Math. Inf. Sci.* 11 (5) (2017), 1299-1307.
- [20] M. Vivas Cortez, E. J. H. Hernández, On some new generalized Hermite-Hadamard-Fejér inequalities for product of two operator  $h$ -convex functions, *Appl. Math. Inf. Sci.* 11 (4) (2017), 983-992.
- [21] S.-H. Wang, Hermite-Hadamard type inequalities for operator convex functions on the co-ordinates, *J. Non-linear Sci. Appl.* 10 (3) (2017), 1116-1125.
- [22] S.-H. Wang, New integral inequalities of Hermite-Hadamard type for operator  $m$ -convex and  $(\alpha, m)$ -convex functions, *J. Comput. Anal. Appl.* 22 (4) (2017), 744-753.