

VICTORIA UNIVERSITY
MELBOURNE AUSTRALIA

Assessing the effects of foot strike patterns and shoe types on the control of leg length and orientation in running

This is the Published version of the following publication

Garofolini, Alessandro, Mickle, Karen, McLaughlin, Patrick and Taylor, Simon (2024) Assessing the effects of foot strike patterns and shoe types on the control of leg length and orientation in running. *Scientific Reports*, 14. ISSN 2045-2322

The publisher's official version can be found at
<https://www.nature.com/articles/s41598-024-52446-0>
Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/48403/>



OPEN

Assessing the effects of foot strike patterns and shoe types on the control of leg length and orientation in running

Alessandro Garofolini¹✉, Karen J. Mickle², Patrick McLaughlin¹ & Simon B. Taylor¹

This research investigates the stabilization of leg length and orientation during the landing phase of running, examining the effects of different footwear and foot strike patterns. Analyzing kinematic data from twenty male long-distance runners, both rearfoot and forefoot strikers, we utilized the Uncontrolled Manifold approach to assess stability. Findings reveal that both leg length and orientation are indeed stabilized during landing, challenging the hypothesis that rearfoot strikers exhibit less variance in deviations than forefoot strikers, and that increased footwear assistance would reduce these deviations. Surprisingly, footwear with a lower minimalist index enhanced post-landing stability, suggesting that cushioning contributes to both force dissipation and leg length stability. The study indicates that both foot strike patterns are capable of effectively reducing task-relevant variance, with no inherent restriction on flexibility for rearfoot strikers. However, there is an indication of potential reliance on footwear for stability. These insights advance our understanding of the biomechanics of running, highlighting the role of footwear in stabilizing leg length and orientation, which has significant implications for running efficiency and injury prevention.

Running involves thousands of repetitive jump-land sequences, with each landing exerting significant force on the limb¹. The external ground force during these landings is modulated by muscle-tendon stress and is influenced by changes in effective leg length and its stiffness^{2,3}. Despite its importance, the central nervous system's (CNS) coordination of the redundant motor elements, such as the multiple muscles and joints involved in the landing process, remains an area of intrigue. These elements are considered 'redundant' because there are more motor components available than are strictly necessary to perform the simple mechanical task of landing.

Leg posture during landing is a result of CNS coordination and can be represented by a kinematic vector, typically defined from the foot centre of pressure to the body centre of mass³. This vector's vertical and fore-aft components define effective leg length and orientation⁴, both of which are controlled by the CNS^{5,6}. The influence of footwear and landing style on these control variables, particularly during the landing phase of running, is yet to be fully understood.

The uncontrolled manifold theory (UCM) by Scholz and Schöner⁷ provides a framework to study the CNS's coordination strategy during a redundant movement task^{8,9}. This theory posits that the CNS exploits the redundancy in the motor system by allowing variability in motor commands that do not affect the overall task goal, thereby offering flexibility and adaptability in motor control. UCM analysis distinguishes between two crucial forms of variability in motor control: goal-irrelevant variability and goal-relevant variability. Goal-irrelevant variability represents the part of motor variability that doesn't affect task performance, while goal-relevant variability contributes to achieving task goals. The UCM's application to running has revealed insights into CNS coordination in novice and experienced runners¹⁰, and in runners under fatigue¹¹, but its application to the landing phase remains unexplored.

From a task performance perspective, goal-irrelevant variability represents the repertoire of body-state configurations that will converge toward the goal state (i.e. variability along the manifold space), which expresses *system flexibility*. In contrast, goal-relevant variability represents the neuromotor system's resistance to perturbations and disruptive external influences (i.e. minimal variability distributed perpendicular to the manifold space), which expresses *system stability*. The strength of the attraction for body states to map onto the manifold space (i.e. goal state) is expressed by the analogy of the potential energy of a well (Fig. 1A). Rooted in dynamical systems

¹Institute for Health and Sport (IHES), Victoria University, Melbourne, Australia. ²School of Environmental and Life Sciences, University of Newcastle, Ourimbah, NSW, Australia. ✉email: alessandro.garofolini@vu.edu.au




Figure 1. Schematic overview of variance structure of an effector system. Well analogy in 3D space, illustrating how the goal-relevant variability is dependent upon the central tendency of the system's state where the space that the effector state is attracted upon can be expressed by the analogy of the potential energy of a well (A). The well's basin width (r) represents the goal-irrelevant variability, which is the repertoire of configurations that will converge toward the goal state—i.e. flexibility. The well depth ($1-V_{\text{ORTH}}$) represents the magnitude of goal-relevant variability, where an increased depth indicates the strength of the convergent attraction for motor solutions to meet the goal state. The depth of the well represents a resistance to system perturbations and disruptive external influences—i.e. stability (B). When a task evolves across time, a sequence of manifolds across the slices of time will express the design of the control system (C). This will be observed from the structure of variance relative to the manifold, where changes both parallel and orthogonal will describe the characteristics of the well and hence the design of the system's controllers (D). This figure has been generated by researcher AG using Adobe Illustrator.

theory¹², this analogy serves as a metaphorical representation. The behavior of the system can be visualized as a ball moving in a landscape of hills and valleys, or "wells". The depth of the well symbolizes the stability of the system, indicating how strongly the system is attracted to a particular state. Conversely, the width of the well represents the flexibility of the system, showcasing the range of states the system can adopt while still achieving the task goal (Fig. 1B). This analogy, while metaphorical, offers a tangible lens to interpret the intricate behaviours of the system, bridging the gap between complex numerical data and the overarching narrative of how the locomotor system manages stability and flexibility. Assuming that the salient details of performance that map to the CNS is equivalently represented by the solution space of the UCM, then the ratio of orthogonal deviations relative to parallel deviations is evidence of a CNS plan for a motor synergy^{13,14}. Here, the term *synergy* applies to the situation when the CNS achieves a motor strategy that is both flexible (well radius) and stable (well depth). As a task unfolds temporally, the evolution of the control system's design can be discerned through a time-sequenced series of manifolds (Fig. 1C). The structural organization of variance with respect to these manifolds—comprising

both parallel and orthogonal fluctuations—serves as a descriptor for the control system's architecture. Specifically, variations that align with the manifold's orientation (parallel) reflect the system's adaptive strategies, while those that deviate (orthogonal) reveal the system's compensatory mechanisms. Together, these variations delineate the contours of the 'well' in our metaphor (Fig. 1D), providing a quantifiable measure of the control system's design parameters, namely its stability and flexibility.

The aforementioned theoretical framework sets the stage for our empirical investigation into the functional aspects of running biomechanics. Specifically, we aim to investigate the consistency of leg length and orientation during the landing phase of running and to understand how different foot landing styles and shoe types might influence this. We predict that both leg length and orientation will be controlled by the CNS, with leg length showing more consistent patterns due to the greater forces in the vertical direction. Additionally, we previously found that habitual rearfoot strike runners exhibit a more consistent pattern in controlling leg stiffness during key phases of running, implying a potentially less adaptable but more stable system¹⁵. This is further supported by findings from¹⁶, which suggest greater core stability in rearfoot strikers due to more balanced muscle activation patterns. These findings align with our hypothesis that rearfoot strikers might exhibit greater stability with less flexibility. Regarding the influence of footwear, it has been observed that shoes with more support can lead to adjustments in the overall stiffness of the leg. This adjustment is a response to the footwear's characteristics, aiming to maintain an optimal balance in leg stiffness, which is crucial for efficient running mechanics¹⁷. Consequently, we propose that more supportive footwear might enhance stability for rearfoot strikers by facilitating these adjustments, without necessarily increasing flexibility. Lastly, we hypothesize that forefoot strikers, particularly when wearing shoes with minimal support, will demonstrate increased flexibility, a hypothesis that requires further investigation to establish a direct comparison with rearfoot strikers in terms of adaptable control patterns.

Methods

Participants

Twenty male long-distance runners (age: 31.2 ± 6.9 years, height: 1.77 ± 0.07 m, weight: 73.4 ± 7.9 kg) gave their informed consent to take part in this study which was approved by the High Risk Ethics committee of Victoria University (No. HRE16-061). All research was performed in accordance with the Declaration of Helsinki. Participants were excluded if they had not been running for at least 5 years, with an average of at least 40 km/week, and had not been free of neurological, cardiovascular, or musculoskeletal problems within the previous six months. Participants were classified as rearfoot strikers (RFS, $n = 10$) or forefoot strikers (FFS, $n = 10$) based on their habitual mode of foot strike assessed by analysing the joint ankle moment from foot contact to the time of reaching 1 body weight on the vertical component of the ground reaction force. Runners who displayed a positive (dorsiflexor) moment for at least 90% of the analysed period were classified as RFS; conversely, runner who displayed a negative (plantarflexor) moment for at least 90% of the analysed period were classified as FFS. This classification is more representative of the working condition of the ankle compared to conventional methods¹⁸.

Experimental protocol

Tests were performed on an instrumented treadmill at a fixed speed of 11 km/h. Kinematics of the lower extremities were recorded with a fourteen VICON camera system (Oxford Metrics Ltd, UK) at a sampling rate of 250 Hz. After a standardized 7-min progressive warm-up, participants ran for five minutes in each of the three different kinds of footwear characterised by different minimalist indexes—MI¹⁹ ranging from 0% (maximum structural support) to 100% (less interaction with the foot). The order of presentation was pseudo-randomized. The shoes adopted in our experiments were classified as low MI (Mizuno® Wave Rider 21, MI = 18%), medium MI (Mizuno® Wave Sonic, MI = 56%), and high MI (Vibram® Five fingers, MI = 96%). A rest period of at least 3 min was given between testing conditions.

Data processing and analysis

Joint position was recorded from 21 retro-reflective markers (14 or 9 mm diameter) attached to the pelvis, thigh, shank and feet as per²⁰. For the UCM analysis, the body was represented as a planar system of 7 rigid segments (pelvis, thigh $\times 2$, shank $\times 2$, and feet) (Fig. 2). Raw data were exported to Visual 3D (C-motion) and filtered using a low-pass Butterworth filter (4th order, zero lag) with a cut-off frequency of 15 Hz. We set this frequency based on the point where 95% of the signal's content remained intact, similar to²¹ and Sinclair, Greenhalgh²².

Gait events were defined using the vertical component of the ground reaction force which was low pass filtered using a 35 Hz cut-off frequency²³. An ascending and descending threshold of 20 N identified foot contact (FC), and foot off respectively, as per Garofolini, Oppici²⁴. Two other events were created 40 ms before foot contact (FC - 10), and 40 ms after foot contact (FC + 10). These events were then used to cut the period of interest and normalize the data into 21 equally distant (in time) slices, where the first 10 slices represented the pre landing phase (PRE) and last 10 slices represented the post landing phase (POST). The latter, post landing phase, can be referred to as the impact phase (≈ 50 ms)¹⁵. Data were then exported into Matlab (The MathWorks Inc., Massachusetts, US) to evaluate the structure of variances within the UCM framework.

Uncontrolled manifold formulation

The UCM analysis was computed at each of the 21 slices, encompassing both the PRE and POST landing phases, as well as the FC event. Each time slice corresponded to a time period of 4 ms. According to our hypothesis, elemental variables were segment angles (i.e. θ_p , θ_t , θ_s and θ_f), while the control variables were the vertical (Leg_z) and horizontal (Leg_y) components of the leg effector (Fig. 2). Details of the UCM formulation are reported in Supplementary material. Variance of goal-irrelevant deviations are parallel to the UCM (V_{UCM}), while




Figure 2. Geometric model used to estimate performance variables leg length (leg_z) and leg orientation (leg_y) and joint angle for pelvis (θ_P), thigh (θ_T), shank (θ_S), and foot (θ_F) segments within the sagittal plane (y,z). The leg effector is constructed from the centre of pressure (CoP) to the pelvis mid-point. Note: thigh, shank, and foot segment angles were computed on both left and right sides; the left side is not shown in the figure. Trials were ranked using foot strike as first, where the average foot strike angle (35–64 percentile) was selected; then a second (nested) classification was done by selecting the trials with a moderate (35–64 percentile) change in leg length (defined as the change during the period [FC, FC + 10]). The UCM analysis was then performed on each time slice. For each slice a linearized manifold (M) and a normal vector (\vec{n}) were computed; then the sum of all distances parallel (V_{UCM}) and orthogonal (V_{ORTH}) and their ratio (V_{RATIO}) were calculated. Using V_{UCM} to represent flexibility and $1 - V_{ORTH}$ to represent stability the possible well structures are shown. This figure has been generated by researcher AG using Power Point 2016.

goal-relevant deviations are orthogonal to the UCM (V_{ORTH}). A third UCM parameter V_{RATIO} was computed from the ratio of V_{UCM} and V_{ORTH} ²⁵.

Although the UCM analysis does not require temporal order of trials, it does presume that the effector repeatedly attempts the same task-goal, from a similar initial configuration state and with similar response behaviour. In this case the task-goal was landing, configuration was the foot strike angle, and response behaviour was the change in limb length. Trials within each participant were therefore rank-ordered based on foot strike angle (three groups of trials), and then based on change in limb length (three sub-groups of trials). The mid sub-group of the average foot strike angle group (average initial conditions) was considered the most representative and therefore used for further analysis (Fig. 2).

Visualizing system flexibility and stability: the well analogy

To visually represent the balance between flexibility and stability in our system, we employed the well analogy (Figs. 1, 2). We constructed the well by fitting a second order polynomial curve through the V_{UCM} and $1 - V_{ORTH}$ pair of values and forcing the curve to pass through the 0,0 coordinates. Forcing the curve through the origin is consistent with the idea that at the initial state (zero perturbation), there's no deviation in the system's behaviour. In this representation, the well's width (V_{UCM}) symbolizes the system's flexibility, while its depth ($1 - V_{ORTH}$) indicates either the system's stability (for positive values) or instability (for negative values). This metaphorical approach provides an intuitive way to interpret the nuanced behaviours of the system.

Statistical analysis

Dependent variables of V_{UCM} , V_{ORTH} , V_{RATIO} were averaged across the 10 time-slices within both the pre-landing phase and the landing phase, and reported as group mean (group standard deviation) for each footwear condition. The vertical and fore-aft dimensions of the control variable were analysed separately. Cohen's d effect sizes

for each group comparison were calculated as the difference between two means divided by the pooled standard deviation. This metric is used to assess the practical significance of the observed differences. A mixed design 3-factor (*Shoe* × *Phase* × *Group*) repeated-measures ANOVA was used to examine the interaction and main effects of within-subject factors of *Shoe* (3 levels: low MI, medium MI, high MI) and task-dependent *Phase* (2 levels: pre, post), and between-subject factor of foot loading *Group* (2 levels: forefoot, rearfoot) on the three dependent variables of V_{UCM} , V_{ORTH} , V_{RATIO} . Tukey post-hoc analysis was used for multiple pairwise comparisons to explore all possible mean differences, given its suitability for comprehensive exploratory analysis and its ability to control the family-wise error rate effectively. A two-way repeated measures ANOVA with main factors *Group* (3 levels: rearfoot, forefoot, Zero control), and *Time* (21 levels: each time slice) was used to test whether the ratio values were statistically greater than zero ($V_{UCM} > V_{ORTH}$), thus accepting the hypothesis of vertical and fore-aft dimension stabilisation. The Dunnett's multiple comparison correction was applied here, given its efficiency in comparing multiple treatments against a single control condition, which in our case was the 'Zero control' group. Multiple two-way repeated measures ANOVA with between-subject factor of foot loading *Group* (2 levels: forefoot, rearfoot) and within-subject factors of *Time* (21 levels: each time slice) were used to test differences between groups within each shoe condition and leg component on dependent variables (V_{UCM} , V_{ORTH} , and V_{RATIO}). The Sidak method was employed for correcting p values for multiple comparisons in these cases, due to its balance between error control and maintaining statistical power, especially suitable for our moderate number of comparisons. Multiple two-way repeated measures ANOVA with between-subject factor of *Shoe* (3 levels: low MI, medium MI, high MI) and within-subject factors of *Time* (5 levels: FC - 10, FC - 5, FC, FC + 5, FC + 10) were used to test differences in well characteristics between shoes within each time slice and leg component on dependent variables (V_{UCM} , $1 - V_{ORTH}$). Here, the Tukey method was again utilized for multiple comparisons, considering its appropriateness for extensive pairwise comparison in this context. Significance was set at 0.05 for all tests. All statistics were performed using GraphPad Prism (version 9.1.1, GraphPad Software, San Diego, California, USA).

Results

V_{RATIO} values for both the Leg_Z and Leg_Y control variables were statistically different from zero for most of the landing phase for all shoe conditions (Fig. 3). Although not significant, FFS displayed higher V_{RATIO} values in low MI shoes at the beginning of the PRE phase, and in high MI shoes at the end of the POST phase on the Leg_Y component. On the Leg_Z component, FFS had an earlier peak in med MI shoes than RFS.

Figure 4 shows the V_{UCM} and V_{ORTH} along the time course of the landing phase. RFS and FFS groups show similar behaviours, V_{ORTH} decreases in the PRE phase and remains constant after foot contact, showing a statistical difference between PRE and POST phase for the Leg_Z ($p < 0.001$) but not for Leg_Y components ($p = 0.166$; Table 1). For both groups, the elbow in the curve happens earlier in low MI shoes than in med or high MI shoes. Although not significant, FFS tend to have higher values for V_{UCM} in low MI shoes in both Leg_Z and Leg_Y components.

Variance parallel to the UCM, V_{UCM}

There was a main effect of *Phase* ($p = 0.027$; Table 1) only for the horizontal (Leg_Y) control variable. Indicating that on average V_{UCM} was dependent on the phase of the landing task only in Leg_Y . Post-hoc analysis revealed that V_{UCM} pre landing is higher compared to post landing for Leg_Y , while it stays statistically unchanged for Leg_Z (Fig. 4, Table 2).

Variance orthogonal to the UCM, V_{ORTH}

Inverse results were found when testing the differences in variance orthogonal to the UCM (Table 1). There was a main effect of *Phase* ($p < 0.001$) for the vertical (Leg_Z) control variable. Post-hoc analysis revealed that V_{ORTH} pre landing is higher compared to post landing for Leg_Z ; while it stays statistically unchanged for Leg_Y . *Shoe* had a significant main effect on V_{ORTH} ($p = 0.007$) for the vertical component only. Post-hoc tests reveal that V_{ORTH} is higher in low MI shoes compared to med MI and high MI ($p = 0.003$; $p = 0.003$, respectively), indicating that more supportive shoes may induce V_{ORTH} to increase (Fig. 4, Table 2).

Ratio of variances perpendicular and orthogonal to the UCM, V_{RATIO}

There was a main effect of *Phase* ($p < 0.001$, Table 1) for the vertical (Leg_Z) control variable. Post-hoc analysis revealed that V_{RATIO} pre landing is lower compared to post landing for Leg_Z ; while it stays statistically unchanged for Leg_Y . *Shoe* had a significant main effect on V_{RATIO} ($p = 0.006$) for the vertical component only. Post-hoc tests reveal that V_{RATIO} is higher in high MI shoes compared to low MI shoes ($p = 0.063$), and med MI shoes compared to low MI shoes ($p = 0.043$) indicating that more supportive shoes may induce V_{RATIO} to decrease (Fig. 3, Table 2).

Flexibility and stability

Given there were no statistical differences between groups in the two control variables (Leg_Z , Leg_Y), we merged the results from RFS and FFS (Fig. 5, Table 3) to analyse flexibility and stability using the well analogy for state attraction to the task manifold (Fig. 1). There was a main effect of *Time* ($p < 0.0001$) for both the vertical (Leg_Z) and horizontal (Leg_Y) control variables. Post-hoc analysis revealed the following (Table 3): (i) at FC - 10 (~ 40 ms pre foot contact) both control variables are in an unstable and highly flexible state; and (ii) at FC - 5 (~ 20 ms pre foot contact) there is a significant change for both control variables ($p < 0.0001$), where their states can be considered significantly more stable but less flexible. Leg_Z control variable reaches highest stability at post landing, with values 3 times higher than Leg_Y . For both control variables flexibility decreases as a function of time (the well basin narrows as limb loading evolves).




Figure 3. Mean \pm SE ratio values for RFS and FFS groups. Time has been divided in two phases: PRE from 10 frames before foot contact (FC – 10) to foot contact (FC); and POST from FC to 10 frames after foot contact (FC + 10). Note: frames correspond to absolute time (ms); 1 frame = 4 ms. FC + 10 is \sim 15% of stance. * indicates statistically significant difference from the ‘Zero Control’ condition; grey colour refers to FFS, black colour refers to RFS group.

Discussion

In this study we sought to determine whether leg length and orientation are stabilized during the landing phase of running; and whether foot strike pattern and footwear affect such stability. To this aim, we asked two types of runners, based on their habitual foot strike posture at landing, to run in different footwear on an instrumented treadmill. We had three main hypotheses. (1) Both leg length and leg orientation will be stabilized during the landing phase; this was confirmed from the presence of a decrease in variance of goal-relevant deviations (i.e. V_{ORTH}). (2) RFS will have comparatively less variance of both goal-irrelevant deviations (V_{UCM}) and goal-relevant deviations (V_{ORTH}); this was not supported. (3) Increased footwear assistance would reduce the variance of goal-relevant deviations more than goal-irrelevant deviations for RFS, this was also not fully supported by the results.

Kinematic synergy in landing: leg length and orientation

Control variables leg length and orientation are stabilized through a kinematic synergy during landing, irrespective of landing type. The V_{RATIO} was significantly greater than zero (Fig. 3, Table 1), and a rapid reduction in V_{ORTH} occurred as the pre-landing phase evolved, demonstrating that control over limb length and orientation is a relevant goal of the locomotor control system. In contrast, goal-irrelevant variance (V_{UCM}) remained relatively constant, and remarkably larger than V_{ORTH} (i.e. $V_{UCM} > V_{ORTH}$), meaning that variance is structured to provide




Figure 4. Mean \pm SE of Variance components parallel (solid lines) and orthogonal (dashed lines) to the linearized UCM. Note: frames correspond to absolute time (ms); 1 frame = 4 ms. FC + 10 is \sim 15% of stance.

increased stability of leg length and orientation as the impact phase approached ($V_{\text{RATIO-POST}} > V_{\text{RATIO-PRE}}$). In the well analogy of stability (Fig. 1), this means that the depth of the well increases while its width remains relatively unchanged (Fig. 5). These trends are indications of a locomotor control system adhering to a minimal intervention principle²⁶. Our results support the idea that stabilisation of control variables (leg length and orientation) is under a hierarchical control system and subjected to a higher-level cost policy that is task-relevant. The idea of a strong synergy being responsible for stability of the kinematic leg effector during late swing and early stance of running is in agreement with previous studies^{5,27–29}. The general findings from Ivanenko, Cappellini⁵ (method of segment angle covariance) indicate that kinematic properties of leg length and orientation are important global parameters encoded within the CNS. The general concept of synergy is consistent, although the particular approach taken to this conclusion of synergy is different. The segment angle covariance from Ivanenko's principal component analysis (PCA) shows two major components. If one synergy (PCA component 1) relates to limb orientation, while the other synergy (component 2) relates to limb length, then each synergy should be evident from the UCM analysis if variance anisotropy relative to a solution manifold is revealed. While the PCA method was used to identify a synergy (i.e. limb axis length and orientation), the UCM in our study quantifies the hypothesized synergy somewhat more precisely (i.e. flexibility and stability of the synergy).

Variable	Group	Shoe	Phase	Group × Shoe	Group × Phase	Shoe × Phase	Group × Shoe × Phase	Tukey's Shoe effects	Phase effects
Leg_Z									
V _{UCM}	F _(1,18) = 0.88 p = .362	F _(2,36) = 0.33 p = .329	F _(1,18) = 1.33 p = .264	F _(2,36) = 1.47 p = .244	F _(1,18) = 0.15 p = .705	F _(2,36) = 0.28 p = .761	F _(2,36) = 0.22 p = .802		
V _{ORTH}	F _(1,18) = 0.13 p = .723	F_(2,36) = 6.36 p = .007	F_(1,18) = 104.6 p < .001	F _(2,36) = 3.335 p = .079	F _(1,18) = 0.02 p = .892	F _(2,36) = 0.53 p = .592	F _(2,36) = 1.36 p = .270	Low > med: p = .003 Low > high: p = .003 Med > high: p = .739	Pre > Post p < .001
V _{RATIO}	F _(1,18) = 0.43 p = .518	F_(2,36) = 7.95 p = .006	F_(1,18) = 221.8 p < .001	F _(2,36) = 0.47 p = .534	F _(1,18) = 0.11 p = .743	F _(2,36) = 0.37 p = .692	F _(2,36) = 0.41 p = .666	Low < med: p = .043 Low < high: p < .001 Med < high: p = .063	Pre < Post p < .001
Leg_Y									
V _{UCM}	F _(1,18) = 1.19 p = .290	F _(2,36) = 0.74 p = .432	F_(1,18) = 5.79 p = .027	F _(2,36) = 1.98 p = .173	F _(1,18) = 0.17 p = .689	F _(2,36) = 0.26 p = .772	F _(2,36) = 0.22 p = .803		Pre > Post p = .027
V _{ORTH}	F _(1,18) = 0.02 p = .896	F _(2,36) = 0.89 p = .409	F _(1,18) = 2.08 p = .166	F _(2,36) = 0.38 p = .623	F _(1,18) = 0.06 p = .808	F _(2,36) = 0.29 p = .750	F _(2,36) = 0.12 p = .890		
V _{RATIO}	F _(1,18) = 4.39 p = .051	F _(2,36) = 1.29 p = .287	F _(1,18) = 3.83 p = .066	F _(2,36) = 0.28 p = .723	F _(1,18) = 0.004 p = .949	F _(2,36) = 0.09 p = .909	F _(2,36) = 0.23 p = .799		

Table 1. Primary statistical results for differences between Groups, Shoes, and Phase for primary variance characteristics (Table 2): variance parallel to the UCM (V_{UCM}), variance orthogonal (V_{ORTH}), and ratio (V_{RATIO}) for the vertical component (Leg_Z) and horizontal component (Leg_Y). ANOVA results are given for main effects and interactions. Given that interactions effects were not significant, the Tukey's pairwise comparisons results are reported only for the main effect of shoe. Low = low minimal index shoe; med = medium minimal index shoe; high = high minimal index shoe. Statistically significant findings are in bold.

Differential stabilization: leg length vs. leg orientation

We found that both control variables demonstrated relatively similar V_{UCM} , however, they differed in the way that V_{ORTH} was reduced (Fig. 4, Table 2). The leg length (Leg_Z) was stabilized rapidly prior to foot contact, indicated by the rapid decrease in V_{ORTH} . This rapid reduction in V_{ORTH} may represent the convergence toward a stable passive attractor (a deep well in Figs. 1, 5) where higher level intervention is not necessary to provide task stability (passive attractor). The tools to appropriately reconcile the responsible source of this rapid change in V_{ORTH} would require extended analysis that combines surrogate data sets where the segment angle correlations are randomised to reduce their non-trivial covariance structure and then perform the UCM method³⁰. Nevertheless, we can reasonably conclude that a consistent leg length is a goal for the landing task of running. In contrast, the leg orientation (Leg_Y) was relatively less stable during the same period, indicated by a modest reduction in V_{ORTH} . However, the V_{RATIO} of leg orientation was significantly greater than 0, indicating that there is a significant non-trivial structuring of the covariance – a synergy. Control of leg orientation (i.e. fore-aft dimension) has implications with braking forces and it has different implications for RFS and FFS³¹. A RFS enables forwards migration of the centre of pressure, while a FFS opposes this motion. The consequence of a restricted forwards migration leads to an increase in the braking impulse³².

Influence of footwear on stability and flexibility

Leg posture (length and orientation) at landing determines stance goals, such as efficiency, stable trajectory of body centre of mass³³, and loading stresses³⁴. There is no direct evidence in the literature to indicate which dimension of leg posture contributes most to these goals; however, because Leg_Z stability continues to increase as landing evolves (Figs. 4, 5), we argue that Leg_Z is relatively more important than Leg_Y to meeting these essential goals of stance. Indeed, control of both leg length and orientation is important; however, a kinematic effector system cannot simultaneously optimize orthogonal dimensions when each dimension shares common elemental properties. From our results it appears that the locomotor control system is designed to prioritise the vertical dimension. By stabilizing leg length pre landing, the locomotor control system sets consistent initial conditions to control leg stiffness post landing^{17,35–38}. The transient foot–ground impact forces are easily perturbed, and the control system relies less upon feedback-driven corrections to the limb's dynamic state³⁹. Therefore, stability of leg length pre-landing is likely a pre-emptive control strategy to in-turn control the associated properties of leg stiffness and musculoskeletal forces upon landing. It is worth noticing that stability of leg length post landing seems affected by the shoe substrate (Fig. 4): low MI shoes provide more stability post landing (descending values of V_{ORTH}) compared to the Med MI and high MI shoes where V_{ORTH} reaches a minimum and stays constant afterwards, indicating that cushioning may not only help with dissipation of external forces⁴⁰ but also with stability of leg length. Our results offer a new insight in understanding running related injuries: shoe type seems to negatively affect the kinematic stability of the leg effector system, and while the associated effects are unknown, it is likely that a lack of leg length stability will have a negative effect on tissue load and energy efficiency. This instability can be particularly evident in real-world running scenarios, such as when a runner

Z component	PRE			POST		
	RFS	FFS	Cohen's d	RFS	FFS	Cohen's d
Low MI						
V _{UCM}	1.12 ± 0.31	1.28 ± 0.30	-0.52	1.03 ± 0.30	1.09 ± 0.44	-0.15
V _{ORTH}	0.98 ± 0.25	1.05 ± 0.26	-0.27	0.49 ± 0.14	0.52 ± 0.13	-0.21
V _{RATIO}	0.09 ± 0.10	0.12 ± 0.11	-0.27	0.37 ± 0.08	0.40 ± 0.08	-0.38
Med MI						
V _{UCM}	1.14 ± 0.29	1.15 ± 0.19	-0.04	1.08 ± 0.30	1.05 ± 0.28	0.11
V _{ORTH}	0.95 ± 0.26	0.88 ± 0.15	0.29	0.44 ± 0.12	0.45 ± 0.10	-0.09
V _{RATIO}	0.11 ± 0.11	0.15 ± 0.08	-0.36	0.44 ± 0.09	0.43 ± 0.06	0.13
High MI						
V _{UCM}	1.16 ± 0.23	1.23 ± 0.26	-0.27	1.05 ± 0.30	1.12 ± 0.30	-0.23
V _{ORTH}	0.87 ± 0.24	0.91 ± 0.24	-0.17	0.41 ± 0.10	0.46 ± 0.12	-0.42
V _{RATIO}	0.16 ± 0.10	0.17 ± 0.10	-0.10	0.45 ± 0.06	0.45 ± 0.07	0.00
Y component	PRE			POST		
	RFS	FFS		RFS	FFS	
Low MI						
V _{UCM}	1.15 ± 0.33	1.33 ± 0.36	-0.51	0.90 ± 0.33	0.97 ± 0.48	-0.16
V _{ORTH}	0.94 ± 0.17	0.93 ± 0.16	0.06	0.82 ± 0.21	0.87 ± 0.10	-0.26
V _{RATIO}	0.09 ± 0.09	0.16 ± 0.14	-0.53	0.06 ± 0.11	0.10 ± 0.11	-0.36
Med MI						
V _{UCM}	1.16 ± 0.29	1.15 ± 0.17	0.04	0.93 ± 0.30	0.90 ± 0.32	0.09
V _{ORTH}	0.92 ± 0.24	0.89 ± 0.21	0.13	0.87 ± 0.25	0.84 ± 0.21	0.13
V _{RATIO}	0.12 ± 0.10	0.13 ± 0.09	-0.10	0.06 ± 0.14	0.09 ± 0.07	-0.23
High MI						
V _{UCM}	1.15 ± 0.25	1.25 ± 0.27	-0.37	0.91 ± 0.33	0.99 ± 0.36	-0.22
V _{ORTH}	0.90 ± 0.20	0.88 ± 0.20	0.10	0.81 ± 0.20	0.82 ± 0.12	-0.06
V _{RATIO}	0.12 ± 0.09	0.17 ± 0.09	-0.56	0.08 ± 0.09	0.14 ± 0.09	-0.67

Table 2. Mean ± standard deviation and Cohen's d effect sizes for variance parallel (V_{UCM}), orthogonal (V_{ORTH}), and ratio (V_{RATIO}) across the three footwear conditions for the vertical (Z) component and horizontal (Y) component. The effect sizes were calculated to provide a measure of the magnitude of differences between groups.

encounters an unexpected increase in ground height, like a rock, curb, or slope. In such situations, if the leg is not in the desired position to accept body weight due to altered stability, it can lead to stumbling or increased difficulty in maintaining efficient running mechanics.

Rearfoot landing: implications for flexibility and shoe assistance

The hypothesis that habitual rearfoot landing restricts the flexibility of the two control variables was not statistically supported (Fig. 4, Table 2), although the results indicate a trend from two footwear conditions (low MI and high MI shoes) where RFS showed a lower V_{UCM} . Any decrease of the observed V_{UCM} may represent a reduction in the neuromotor repertoire of flexible solutions, and evidence of a system designed for minimal intervention that can default to the low-level allometric control process⁴¹. Possessing an abundant and flexible system could be important if the landing phase requires a rapid movement solution from low-level control mechanisms^{42,43}. The RFS group may have restricted their available degrees of freedom and the potential to find flexible motor solutions for landing, and in consequence they may become dependent upon the shoe for assistance⁴⁴. The implications for a RFS runner relate to situations when they are faced with alternate conditions. For example, many athletes habituate to a RFS technique but may perform in endurance competitions where they use high MI footwear. Future research will need to determine if RFS-related loss in landing flexibility exists and persists, leading to loss of energy efficiency and increased injury risk when faced with situations that impose alternate substrate conditions. Additionally, it would be valuable to explore what happens when habitual rearfoot strikers are presented with scenarios requiring a more adaptable control strategy, such as a change in foot striking pattern in response to certain perturbations or testing constraints. In these situations, the enhanced stability or reduced number of solutions typically experienced by rearfoot strikers could potentially lead to decreased energy efficiency^{45,46} and an increased injury risk⁴⁷. This consideration is particularly relevant in understanding how habitual movement patterns adapt to unexpected changes in running dynamics.

Study limitations and assumptions

A limitation of this study is related to the biomechanical model, as we assumed the central point of the pelvis to be the goal position of the effector endpoint, although the true goal of the effector endpoint is likely to be the




Figure 5. Stability ($1-V_{ORTH}$) and flexibility (V_{UCM}) using the well analogy. Groups and shoes have been merged. Five time slices were compared: Foot Contact (FC), ten time slices before foot contact (FC - 10), five time slices before foot contact (FC - 5), 5 time slices after foot contact (FC + 5), ten time slices after foot contact (FC + 10).

Tukey's multiple comparisons test	Leg _z		Leg _y	
	V_{UCM}	$1-V_{ORTH}$	V_{UCM}	$1-V_{ORTH}$
FC - 10 vs. FC - 5	0.12	-0.57**	0.29**	-0.22**
FC - 10 vs. FC	0.14*	-0.86**	0.39**	-0.22**
FC - 10 vs. FC + 5	0.15*	-0.97**	0.43**	-0.24**
FC - 10 vs. FC + 10	0.19**	-0.97**	0.46**	-0.26**
FC - 5 vs. FC	0.01	-0.29**	0.10	-0.01
FC - 5 vs. FC + 5	0.02	-0.40**	0.14	-0.02
FC - 5 vs. FC + 10	0.07	-0.40**	0.18**	-0.04
FC vs. FC + 5	0.01	-0.11	0.04	-0.01
FC vs. FC + 10	0.05	-0.11	0.07	-0.03
FC + 5 vs. FC + 10	0.04	0.00	0.04	-0.02

Table 3. Post-hoc analysis comparing each time slices for flexibility (V_{UCM}) and stability (V_{ORTH}) within each control variable (Leg_z and Leg_y). *Indicates significant differences at $p < 0.05$, **Indicates significant differences at $p < 0.001$.

body CoM. When interpreting the results of the UCM method we are unable to distinguish which joint is more responsible for effector endpoint change. Specifically, we did not test the sensitivity of the effector endpoint position to individual joints as we assume the effector endpoint position will change an equivalent displacement per change in unit radian of each elemental variable. It may be that the system becomes more sensitive to certain elemental variables in unfamiliar conditions⁴⁸. Another limitation of the study is the limited sample size. To address this, in addition to standard statistical analyses, we computed Cohen's d effect sizes for each comparison (Table 2). Reporting effect sizes allows us to provide a nuanced understanding of the data, complementing inferential statistics and offering insights into the potential real-world implications of our findings.

Conclusion

In conclusion, FFS and RFS are equally able to reduce task-relevant variance in order to stabilize the control variables leg length and leg orientation. The assistance provided by the shoe may affect the control variables (leg length and orientation) with possible implications for force control at landing.

Data availability

The datasets generated during and analysed during the current study are available from the corresponding author on reasonable request.

Received: 15 November 2023; Accepted: 18 January 2024

Published online: 26 January 2024

References

- Hoeger, W. W. *et al.* One-mile step count at walking and running speeds. *ACSMs Health Fit. J.* **12**(1), 14–19 (2008).
- Dorn, T. W., Schache, A. G. & Pandy, M. G. Muscular strategy shift in human running: Dependence of running speed on hip and ankle muscle performance. *J. Exp. Biol.* **215**(11), 1944–1956 (2012).
- Silder, A., Besier, T. & Delp, S. L. Running with a load increases leg stiffness. *J. Biomech.* **48**(6), 1003–1008 (2015).
- Auyang, A. G., Yen, J. T. & Chang, Y.-H. Neuromechanical stabilization of leg length and orientation through interjoint compensation during human hopping. *Exp. Brain Res.* **192**(2), 253–264 (2009).
- Ivanenko, Y. P. *et al.* Modular control of limb movements during human locomotion. *J. Neurosci.* **27**(41), 11149–11161 (2007).
- Ivanenko, Y. P. *et al.* On the origin of planar covariation of elevation angles during human locomotion. *J. Neurophysiol.* **99**(4), 1890–1898 (2008).
- Scholz, J. P. & Schöner, G. The uncontrolled manifold concept: Identifying control variables for a functional task. *Exp. Brain Res.* **126**(3), 289–306 (1999).
- Latash, M. L. Abundant degrees of freedom are not a problem. *Kinesiol. Rev.* **7**(1), 64–72 (2018).
- Sternad, D. It's not (only) the mean that matters: Variability, noise and exploration in skill learning. *Curr. Opin. Behav. Sci.* **20**, 183–195 (2018).
- Möhler, F. *et al.* Variability of running coordination in experts and novices: A 3D uncontrolled manifold analysis. *Eur. J. Sport Sci.* **20**(9), 1187–1196 (2020).
- Möhler, F. *et al.* Influence of fatigue on running coordination: A UCM analysis with a geometric 2D model and a subject-specific anthropometric 3D model. *Hum. Mov. Sci.* **66**, 133–141 (2019).
- Kelso, J. S. *Dynamic Patterns: The Self-organization of Brain and Behavior* (MIT Press, 1995).
- Latash, M. L., Scholz, J. P. & Schöner, G. Toward a new theory of motor synergies. *Motor Control* **11**(3), 276–308 (2007).
- Todorov, E., Li, W. & Pan, X. From task parameters to motor synergies: A hierarchical framework for approximately optimal control of redundant manipulators. *J. Robot. Syst.* **22**(11), 691–710 (2005).
- Garofolini, A. *et al.* Insight into the hierarchical control governing leg stiffness during the stance phase of running. *Sci. Rep.* **12**(1), 12123 (2022).
- Swager van Dok, J. & Cabri, J. Feedforward activity of the transversus abdominis and multifidus muscle in a forefoot versus rearfoot strike running pattern. *Man. Ther.* **13**(04), 156–166 (2009).
- Ferris, D. P., Louie, M. & Farley, C. T. Running in the real world: adjusting leg stiffness for different surfaces. *Proc. R. Soc. Lond. Ser. B Biol. Sci.* **265**(1400), 989–994 (1998).
- Garofolini, A. *et al.* Foot strike classification: a comparison of methodologies. *Footwear Sci.* **9**(sup1), S129–S130 (2017).
- Esculier, J.-F. *et al.* A consensus definition and rating scale for minimalist shoes. *J. Foot Ankle Res.* **8**(1), 1–9 (2015).
- Garofolini, A. *et al.* Ankle joint dynamic stiffness in long-distance runners: Effect of foot strike and shoes features. *Appl. Sci.* **9**(19), 4100 (2019).
- Fellin, R. E., Manal, K. & Davis, I. S. Comparison of lower extremity kinematic curves during overground and treadmill running. *J. Appl. Biomech.* **26**(4), 407–414 (2010).
- Sinclair, J. *et al.* The influence of barefoot and barefoot-inspired footwear on the kinetics and kinematics of running in comparison to conventional running shoes. *Footwear Sci.* **5**(1), 45–53 (2013).
- Garofolini, A., Taylor, S. & Lepine, J. Evaluating dynamic error of a treadmill and the effect on measured kinetic gait parameters: Implications and possible solutions. *J. Biomech.* **82**, 156–163 (2019).
- Garofolini, A., Oppici, L. & Taylor, S. A real-time feedback method to reduce loading rate during running: Effect of combining direct and indirect feedback. *J. Sports Sci.* **38**(21), 2446–2453 (2020).
- Papi, E., Rowe, P. J. & Pomeroy, V. M. Analysis of gait within the uncontrolled manifold hypothesis: Stabilisation of the centre of mass during gait. *J. Biomech.* **48**(2), 324–331 (2015).
- Todorov, E. & Jordan, M. I. Optimal feedback control as a theory of motor coordination. *Nat. Neurosci.* **5**(11), 1226–1235 (2002).
- Blum, Y. *et al.* Swing-leg trajectory of running guinea fowl suggests task-level priority of force regulation rather than disturbance rejection. *PLoS One* **9**(6), e100399 (2014).
- Blum, Y. *et al.* Swing leg control in human running. *Bioinspir. Biomim.* **5**(2), 026006 (2010).
- Daley, M. A. & Usherwood, J. R. Two explanations for the compliant running paradox: Reduced work of bouncing viscera and increased stability in uneven terrain. *Biol. Lett.* **6**(3), 418–421 (2010).
- Scholz, J. P. & Schöner, G. Use of the uncontrolled manifold (UCM) approach to understand motor variability, motor equivalence, and self-motion. In *Progress in Motor Control* 91–100 (Springer, 2014).
- Boyer, E. R., Rooney, B. D. & Derrick, T. R. Rearfoot and midfoot or forefoot impacts in habitually shod runners. *Med. Sci. Sports Exerc.* **46**(7), 1384–1391 (2014).
- Nordin, A. D., Dufek, J. S. & Mercer, J. A. Three-dimensional impact kinetics with foot-strike manipulations during running. *J. Sport Health Sci.* **6**(4), 489–497 (2017).
- Seyfarth, A., Geyer, H. & Herr, H. Swing-leg retraction: A simple control model for stable running. *J. Exp. Biol.* **206**(15), 2547–2555 (2003).
- Vejdani, H. *et al.* Bio-inspired swing leg control for spring-mass robots running on ground with unexpected height disturbance. *Bioinspir. Biomim.* **8**(4), 046006 (2013).
- Cusumano, J. P. & Cesari, P. Body-goal variability mapping in an aiming task. *Biol. Cybern.* **94**(5), 367–379 (2006).
- Birn-Jeffery, A. V. *et al.* Don't break a leg: Running birds from quail to ostrich prioritise leg safety and economy on uneven terrain. *J. Exp. Biol.* **217**(21), 3786–3796 (2014).
- Shen, Z. & Seipel, J. Effective leg stiffness of animal running and the co-optimization of energetic cost and stability. *J. Theor. Biol.* **451**, 57–66 (2018).
- Shen, Z. & Seipel, J. The leg stiffnesses animals use may improve the stability of locomotion. *J. Theor. Biol.* **377**, 66–74 (2015).
- Daley, M. A. & Biewener, A. A. Running over rough terrain reveals limb control for intrinsic stability. *Proc. Natl. Acad. Sci.* **103**(42), 15681–15686 (2006).
- Gruber, A. H. *et al.* Impact shock frequency components and attenuation in rearfoot and forefoot running. *J. Sport Health Sci.* **3**(2), 113–121 (2014).
- Dingwell, J. B., John, J. & Cusumano, J. P. Do humans optimally exploit redundancy to control step variability in walking?. *PLoS Comput. Biol.* **6**(7), e1000856 (2010).

42. Krogt, M. M. V. D. *et al.* Robust passive dynamics of the musculoskeletal system compensate for unexpected surface changes during human hopping. *J. Appl. Physiol.* **107**(3), 801–808 (2009).
43. Moritz, C. T. & Farley, C. T. Human hopping on very soft elastic surfaces: Implications for muscle pre-stretch and elastic energy storage in locomotion. *J. Exp. Biol.* **208**, 939–949 (2005).
44. Gruber, A. H. *et al.* Footfall patterns during barefoot running on harder and softer surfaces. *Footwear Sci.* **5**(1), 39–44 (2013).
45. Cavanagh, P. R. & Williams, K. R. The effect of stride length variation on oxygen uptake during distance running. *Med. Sci. Sports Exerc.* **14**(1), 30–35 (1982).
46. Yin, L. *et al.* Leg stiffness and vertical stiffness of habitual forefoot and rearfoot strikers during running. *Appl. Bionics Biomech.* **2020** (2020).
47. Zhou, W. *et al.* Surface effects on kinematics, kinetics and stiffness of habitual rearfoot strikers during running. *Plos One* **18**(3), e0283323 (2023).
48. Yen, J. T. & Chang, Y.-H. Rate-dependent control strategies stabilize limb forces during human locomotion. *J. R. Soc. Interface* **7**(46), 801 (2010).

Acknowledgements

The authors gratefully acknowledge Mizuno® Australia for donating the footwear used in the study.

Author contributions

A.G.: Conceptualization, Methodology, Software, Formal Analysis, Investigation, Resources, Data curation, Writing original draft, Writing review and editing, Visualisation. K.J.M.: Conceptualization, Methodology, Resources, Writing review and editing. P.M.: Writing review and editing. S.B.T.: Conceptualization, Methodology, Software, Formal Analysis, Supervision, Writing review and editing.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-024-52446-0>.

Correspondence and requests for materials should be addressed to A.G.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2024