



**VICTORIA UNIVERSITY**  
MELBOURNE AUSTRALIA

## *Fejer-type inequalities (I)*

This is the Published version of the following publication

Tseng, Kuei-Lin, Hwang, Shioh-Ru and Dragomir, Sever S (2010) Fejer-type inequalities (I). *Journal of Inequalities and Applications*, 2010. pp. 1-7. ISSN 1025-5834 (print) 1029-242X (online)

The publisher's official version can be found at  
<http://www.journalofinequalitiesandapplications.com/content/2010/1/531976>  
Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/7359/>

## Research Article

# Fejér-Type Inequalities (I)

**Kuei-Lin Tseng,<sup>1</sup> Shiow-Ru Hwang,<sup>2</sup> and S. S. Dragomir<sup>3,4</sup>**

<sup>1</sup> Department of Applied Mathematics, Aletheia University, Tamsui 25103, Taiwan

<sup>2</sup> China University of Science and Technology, Nankang, Taipei 11522, Taiwan

<sup>3</sup> School of Engineering Science, VIC University, P.O. Box 14428, Melbourne City MC, Victoria 8001, Australia

<sup>4</sup> School of Computational and Applied Mathematics, University of the Witwatersrand, Private Bag 3, Wits 2050, Johannesburg, South Africa

Correspondence should be addressed to S. S. Dragomir, sever.dragomir@vu.edu.au

Received 3 May 2010; Revised 26 August 2010; Accepted 3 December 2010

Academic Editor: Yeol J. E. Cho

Copyright © 2010 Kuei-Lin Tseng et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We establish some new Fejér-type inequalities for convex functions.

## 1. Introduction

Throughout this paper, let  $f : [a, b] \rightarrow \mathbb{R}$  be convex, and let  $g : [a, b] \rightarrow [0, \infty)$  be integrable and symmetric to  $(a + b)/2$ . We define the following functions on  $[0, 1]$  that are associated with the well-known Hermite-Hadamard inequality [1]

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2}, \quad (1.1)$$

namely

$$I(t) = \int_a^b \frac{1}{2} \left[ f\left(t \frac{x+a}{2} + (1-t) \frac{a+b}{2}\right) + f\left(t \frac{x+b}{2} + (1-t) \frac{a+b}{2}\right) \right] g(x) dx,$$

$$J(t) = \int_a^b \frac{1}{2} \left[ f\left(t \frac{x+a}{2} + (1-t) \frac{3a+b}{4}\right) + f\left(t \frac{x+b}{2} + (1-t) \frac{a+3b}{4}\right) \right] g(x) dx,$$

$$\begin{aligned}
M(t) &= \int_a^{(a+b)/2} \frac{1}{2} \left[ f\left(ta + (1-t)\frac{x+a}{2}\right) + f\left(t\frac{a+b}{2} + (1-t)\frac{x+b}{2}\right) \right] g(x) dx \\
&\quad + \int_{(a+b)/2}^b \frac{1}{2} \left[ f\left(t\frac{a+b}{2} + (1-t)\frac{x+a}{2}\right) + f\left(tb + (1-t)\frac{x+b}{2}\right) \right] g(x) dx, \\
N(t) &= \int_a^b \frac{1}{2} \left[ f\left(ta + (1-t)\frac{x+a}{2}\right) + f\left(tb + (1-t)\frac{x+b}{2}\right) \right] g(x) dx.
\end{aligned} \tag{1.2}$$

For some results which generalize, improve, and extend the famous integral inequality (1.1), see [2–6].

In [2], Dragomir established the following theorem which is a refinement of the first inequality of (1.1).

**Theorem A.** Let  $f$  be defined as above, and let  $H$  be defined on  $[0, 1]$  by

$$H(t) = \frac{1}{b-a} \int_a^b f\left(tx + (1-t)\frac{a+b}{2}\right) dx. \tag{1.3}$$

Then,  $H$  is convex, increasing on  $[0, 1]$ , and for all  $t \in [0, 1]$ , one has

$$f\left(\frac{a+b}{2}\right) = H(0) \leq H(t) \leq H(1) = \frac{1}{b-a} \int_a^b f(x) dx. \tag{1.4}$$

In [6], Yang and Hong established the following theorem which is a refinement of the second inequality in (1.1).

**Theorem B.** Let  $f$  be defined as above, and let  $P$  be defined on  $[0, 1]$  by

$$P(t) = \frac{1}{2(b-a)} \int_a^b \left[ f\left(\left(\frac{1+t}{2}\right)a + \left(\frac{1-t}{2}\right)x\right) + f\left(\left(\frac{1+t}{2}\right)b + \left(\frac{1-t}{2}\right)x\right) \right] dx. \tag{1.5}$$

Then,  $P$  is convex, increasing on  $[0, 1]$ , and for all  $t \in [0, 1]$ , one has

$$\frac{1}{b-a} \int_a^b f(x) dx = P(0) \leq P(t) \leq P(1) = \frac{f(a) + f(b)}{2}. \tag{1.6}$$

In [3], Fejér established the following weighted generalization of the Hermite-Hadamard inequality (1.1).

**Theorem C.** Let  $f, g$  be defined as above. Then,

$$f\left(\frac{a+b}{2}\right) \int_a^b g(x) dx \leq \int_a^b f(x) g(x) dx \leq \frac{f(a) + f(b)}{2} \int_a^b g(x) dx \quad (1.7)$$

is known as Fejér inequality.

In this paper, we establish some Fejér-type inequalities related to the functions  $I, J, M, N$  introduced above.

## 2. Main Results

In order to prove our main results, we need the following lemma.

**Lemma 2.1** (see [4]). Let  $f$  be defined as above, and let  $a \leq A \leq C \leq D \leq b$  with  $A+B = C+D$ . Then,

$$f(C) + f(D) \leq f(A) + f(B). \quad (2.1)$$

Now, we are ready to state and prove our results.

**Theorem 2.2.** Let  $f, g$ , and  $I$  be defined as above. Then  $I$  is convex, increasing on  $[0, 1]$ , and for all  $t \in [0, 1]$ , one has the following Fejér-type inequality:

$$f\left(\frac{a+b}{2}\right) \int_a^b g(x) dx = I(0) \leq I(t) \leq I(1) = \int_a^b \frac{1}{2} \left[ f\left(\frac{x+a}{2}\right) + f\left(\frac{x+b}{2}\right) \right] g(x) dx. \quad (2.2)$$

*Proof.* It is easily observed from the convexity of  $f$  that  $I$  is convex on  $[0, 1]$ . Using simple integration techniques and under the hypothesis of  $g$ , the following identity holds on  $[0, 1]$ :

$$\begin{aligned} I(t) &= \int_a^b \frac{1}{2} \left[ f\left(t \frac{x+a}{2} + (1-t) \frac{a+b}{2}\right) g(x) + f\left(t \frac{a+2b-x}{2} + (1-t) \frac{a+b}{2}\right) g(a+b-x) \right] dx \\ &= \int_a^b \frac{1}{2} \left[ f\left(t \frac{x+a}{2} + (1-t) \frac{a+b}{2}\right) + f\left(t \frac{a+2b-x}{2} + (1-t) \frac{a+b}{2}\right) \right] g(x) dx \\ &= \int_a^{(a+b)/2} \left[ f\left(tx + (1-t) \frac{a+b}{2}\right) + f\left(t(a+b-x) + (1-t) \frac{a+b}{2}\right) \right] g(2x-a) dx. \end{aligned} \quad (2.3)$$

Let  $t_1 < t_2$  in  $[0, 1]$ . By Lemma 2.1, the following inequality holds for all  $x \in [a, (a+b)/2]$ :

$$\begin{aligned} & f\left(t_1x + (1-t_1)\frac{a+b}{2}\right) + f\left(t_1(a+b-x) + (1-t_1)\frac{a+b}{2}\right) \\ & \leq f\left(t_2x + (1-t_2)\frac{a+b}{2}\right) + f\left(t_2(a+b-x) + (1-t_2)\frac{a+b}{2}\right). \end{aligned} \quad (2.4)$$

Indeed, it holds when we make the choice

$$\begin{aligned} A &= t_2x + (1-t_2)\frac{a+b}{2}, \\ C &= t_1x + (1-t_1)\frac{a+b}{2}, \\ D &= t_1(a+b-x) + (1-t_1)\frac{a+b}{2}, \\ B &= t_2(a+b-x) + (1-t_2)\frac{a+b}{2}, \end{aligned} \quad (2.5)$$

in Lemma 2.1.

Multiplying the inequality (2.4) by  $g(2x-a)$ , integrating both sides over  $x$  on  $[a, (a+b)/2]$  and using identity (2.3), we derive  $I(t_1) \leq I(t_2)$ . Thus  $I$  is increasing on  $[0, 1]$  and then the inequality (2.2) holds. This completes the proof.  $\square$

**Remark 2.3.** Let  $g(x) = 1/(b-a)$  ( $x \in [a, b]$ ) in Theorem 2.2. Then  $I(t) = H(t)$  ( $t \in [0, 1]$ ) and the inequality (2.2) reduces to the inequality (1.4), where  $H$  is defined as in Theorem A.

**Theorem 2.4.** Let  $f, g, J$  be defined as above. Then  $J$  is convex, increasing on  $[0, 1]$ , and for all  $t \in [0, 1]$ , one has the following Fejér-type inequality:

$$\begin{aligned} & \frac{f((3a+b)/4) + f((a+3b)/4)}{2} \int_a^b g(x)dx = J(0) \leq J(t) \leq J(1) \\ & = \frac{1}{2} \int_a^b \left[ f\left(\frac{x+a}{2}\right) + f\left(\frac{x+b}{2}\right) \right] g(x)dx. \end{aligned} \quad (2.6)$$

*Proof.* By using a similar method to that from Theorem 2.2, we can show that  $J$  is convex on  $[0, 1]$ , the identity

$$\begin{aligned} J(t) &= \int_a^{(3a+b)/4} \left[ f\left(tx + (1-t)\frac{3a+b}{4}\right) + f\left(t\left(\frac{3a+b}{2} - x\right) + (1-t)\frac{3a+b}{4}\right) \right. \\ & \quad \left. + f\left(t\left(x + \frac{b-a}{2}\right) + (1-t)\frac{a+3b}{4}\right) + f\left(t(a+b-x) + (1-t)\frac{a+3b}{4}\right) \right] \\ & \quad \times g(2x-a)dx \end{aligned} \quad (2.7)$$

holds on  $[0, 1]$ , and the inequalities

$$\begin{aligned} & f\left(t_1x + (1-t_1)\frac{3a+b}{4}\right) + f\left(t_1\left(\frac{3a+b}{2} - x\right) + (1-t_1)\frac{3a+b}{4}\right) \\ & \leq f\left(t_2x + (1-t_2)\frac{3a+b}{4}\right) + f\left(t_2\left(\frac{3a+b}{2} - x\right) + (1-t_2)\frac{3a+b}{4}\right), \end{aligned} \quad (2.8)$$

$$\begin{aligned} & f\left(t_1\left(x + \frac{b-a}{2}\right) + (1-t_1)\frac{a+3b}{4}\right) + f\left(t_1(a+b-x) + (1-t_1)\frac{a+3b}{4}\right) \\ & \leq f\left(t_2\left(x + \frac{b-a}{2}\right) + (1-t_2)\frac{a+3b}{4}\right) + f\left(t_2(a+b-x) + (1-t_2)\frac{a+3b}{4}\right) \end{aligned} \quad (2.9)$$

hold for all  $t_1 < t_2$  in  $[0, 1]$  and  $x \in [a, (3a+b)/4]$ .

By (2.7)–(2.9) and using a similar method to that from Theorem 2.2, we can show that  $J$  is increasing on  $[0, 1]$  and (2.6) holds. This completes the proof.  $\square$

The following result provides a comparison between the functions  $I$  and  $J$ .

**Theorem 2.5.** *Let  $f, g, I$ , and  $J$  be defined as above. Then  $I(t) \leq J(t)$  on  $[0, 1]$ .*

*Proof.* By the identity

$$J(t) = \int_a^{(a+b)/2} \left[ f\left(tx + (1-t)\frac{3a+b}{4}\right) + f\left(t(a+b-x) + (1-t)\frac{a+3b}{4}\right) \right] g(2x-a) dx, \quad (2.10)$$

on  $[0, 1]$ , (2.3) and using a similar method to that from Theorem 2.2, we can show that  $I(t) \leq J(t)$  on  $[0, 1]$ . The details are omitted.  $\square$

Further, the following result incorporates the properties of the function  $M$ .

**Theorem 2.6.** *Let  $f, g, M$  be defined as above. Then  $M$  is convex, increasing on  $[0, 1]$ , and for all  $t \in [0, 1]$ , one has the following Fejér-type inequality:*

$$\begin{aligned} & \int_a^b \frac{1}{2} \left[ f\left(\frac{x+a}{2}\right) + f\left(\frac{x+b}{2}\right) \right] g(x) dx \\ & = M(0) \leq M(t) \leq M(1) = \frac{1}{2} \left[ f\left(\frac{a+b}{2}\right) + \frac{f(a)+f(b)}{2} \right] \int_a^b g(x) dx. \end{aligned} \quad (2.11)$$

*Proof.* Follows by the identity

$$M(t) = \int_a^{(3a+b)/4} \left[ f(ta + (1-t)x) + f\left(t\frac{a+b}{2} + (1-t)\left(\frac{3a+b}{2} - x\right)\right) \right. \\ \left. + f\left(t\frac{a+b}{2} + (1-t)\left(x + \frac{b-a}{2}\right)\right) + f(tb + (1-t)(a+b-x)) \right] \\ \times g(2x-a)dx, \quad (2.12)$$

on  $[0, 1]$ . The details are left to the interested reader.  $\square$

We now present a result concerning the properties of the function  $N$ .

**Theorem 2.7.** *Let  $f, g, N$  be defined as above. Then  $N$  is convex, increasing on  $[0, 1]$ , and for all  $t \in [0, 1]$ , one has the following Fejér-type inequality:*

$$\int_a^b \frac{1}{2} \left[ f\left(\frac{x+a}{2}\right) + f\left(\frac{x+b}{2}\right) \right] g(x)dx = N(0) \leq N(t) \leq N(1) = \frac{f(a) + f(b)}{2} \int_a^b g(x)dx. \quad (2.13)$$

*Proof.* By the identity

$$N(t) = \int_a^{(a+b)/2} [f(ta + (1-t)x) + f(tb + (1-t)(a+b-x))] g(2x-a)dx \quad (2.14)$$

on  $[0, 1]$  and using a similar method to that for Theorem 2.2, we can show that  $N$  is convex, increasing on  $[0, 1]$  and (2.13) holds.  $\square$

**Remark 2.8.** Let  $g(x) = 1/(b-a)$  ( $x \in [a, b]$ ) in Theorem 2.7. Then  $N(t) = P(t)$  ( $t \in [0, 1]$ ) and the inequality (2.13) reduces to (1.6), where  $P$  is defined as in Theorem B.

**Theorem 2.9.** *Let  $f, g, M$ , and  $N$  be defined as above. Then  $M(t) \leq N(t)$  on  $[0, 1]$ .*

*Proof.* By the identity

$$N(t) = \int_a^{(3a+b)/4} \left[ f(ta + (1-t)x) + f\left(ta + (1-t)\left(\frac{3a+b}{2} - x\right)\right) \right. \\ \left. + f(tb + (1-t)(a+b-x)) + f\left(tb + (1-t)\left(x + \frac{b-a}{2}\right)\right) \right] g(2x-a)dx, \quad (2.15)$$

on  $[0, 1]$ , (2.12) and using a similar method to that for Theorem 2.2, we can show that  $M(t) \leq N(t)$  on  $[0, 1]$ . This completes the proof.  $\square$

The following Fejér-type inequality is a natural consequence of Theorems 2.2–2.9.

**Corollary 2.10.** Let  $f, g$  be defined as above. Then one has

$$\begin{aligned} f\left(\frac{a+b}{2}\right) \int_a^b g(x) dx &\leq \frac{f((3a+b)/4) + f((a+3b)/4)}{2} \int_a^b g(x) dx \\ &\leq \int_a^b \frac{1}{2} \left[ f\left(\frac{x+a}{2}\right) + f\left(\frac{x+b}{2}\right) \right] g(x) dx \\ &\leq \frac{1}{2} \left[ f\left(\frac{a+b}{2}\right) + \frac{f(a)+f(b)}{2} \right] \int_a^b g(x) dx \\ &\leq \frac{f(a)+f(b)}{2} \int_a^b g(x) dx. \end{aligned} \quad (2.16)$$

**Remark 2.11.** Let  $g(x) = 1/(b-a)$  ( $x \in [a, b]$ ) in Corollary 2.10. Then the inequality (2.16) reduces to

$$\begin{aligned} f\left(\frac{a+b}{2}\right) &\leq \frac{f((3a+b)/4) + f((a+3b)/4)}{2} \leq \frac{1}{b-a} \int_a^b f(x) dx \\ &\leq \frac{1}{2} \left[ f\left(\frac{a+b}{2}\right) + \frac{f(a)+f(b)}{2} \right] \leq \frac{f(a)+f(b)}{2}, \end{aligned} \quad (2.17)$$

which is a refinement of (1.1).

**Remark 2.12.** In Corollary 2.10, the third inequality in (2.16) is the weighted generalization of Bullen's inequality [5]

$$\frac{1}{b-a} \int_a^b f(x) dx \leq \frac{1}{2} \left[ f\left(\frac{a+b}{2}\right) + \frac{f(a)+f(b)}{2} \right]. \quad (2.18)$$

## Acknowledgment

This research was partially supported by Grant NSC 97-2115-M-156-002.

## References

- [1] J. Hadamard, "Étude sur les propriétés des fonctions entières en particulier d'une fonction considérée par Riemann," *Journal de Mathématiques Pures et Appliquées*, vol. 58, pp. 171–215, 1893.
- [2] S. S. Dragomir, "Two mappings in connection to Hadamard's inequalities," *Journal of Mathematical Analysis and Applications*, vol. 167, no. 1, pp. 49–56, 1992.
- [3] L. Fejér, "Über die Fourierreihen, II," *Math. Naturwiss. Anz Ungar. Akad. Wiss.*, vol. 24, pp. 369–390, 1906 (Hungarian).
- [4] D.-Y. Hwang, K.-L. Tseng, and G.-S. Yang, "Some Hadamard's inequalities for co-ordinated convex functions in a rectangle from the plane," *Taiwanese Journal of Mathematics*, vol. 11, no. 1, pp. 63–73, 2007.
- [5] J. E. Pečarić, F. Proschan, and Y. L. Tong, *Convex Functions, Partial Orderings, and Statistical Applications*, vol. 187 of *Mathematics in Science and Engineering*, Academic Press, Boston, Mass, USA, 1992.
- [6] G.-S. Yang and M.-C. Hong, "A note on Hadamard's inequality," *Tamkang Journal of Mathematics*, vol. 28, no. 1, pp. 33–37, 1997.